Lecture I : Liftings of Galois covers of smooth curves

by Michel Matignon

These two lectures are a report on the lifting problem of Galois covers of smooth curves from char. $p > 0$ to char. 0. The references are listed as [G-M 1], [G-M 2], [M].

The first lecture will focus on lifting problems, the main results are a local global principle and the positive answer to the lifting problem for p^2-cyclic covers generalizing a former result in p-cyclic case due to F.Oort, T.Sekiguchi, N.Suwa.

The second lecture will focuss on the geometry of order p-automorphisms of an open disc over a p-adic field.

I would like to thank Professors T.Sekiguchi and N.Suwa for inviting me to Japan and for organizing this symposium which gives me the opportunity for this report. I am very grateful to T.Ito and M.Yato for making the TeX version of my notes of the lectures.

Finally I would like to dedicate this work to Michel Raynaud who has so much influenced us.

Notations:

k is an algebraicaly closed field of char. $p > 0$.
R is a complete DVR finite over $W(k)$.
π is a uniformising parameter for R.
$\mathrm{Fr}R =: K \subset K^{\mathrm{alg}}$ is endowed with the unique polongation of the valuation v.

0. Introduction

Let C/k be a smooth irreducible complete curve of genus g.

We are interested in the following:

Global lifting problem. *Let $G \underset{\text{finite}}{\subset} \mathrm{Aut}_k C$. Is it possible to find an R as above and $\mathcal{C}/R$ a relative smooth R-curve such that $G \subset \mathrm{Aut}_R \mathcal{C}$ and $(\mathcal{C}, G)$ gives $(C, G) \mod \pi$; i.e. we have a commutative diagramm:*



It is possible to formulate this in terms of G-covers. Let



I_y = Inertia group at $y \in C$; then
$I_y \subset \mathrm{Aut}_k \hat{\mathcal{O}}_{C,y} \simeq \mathrm{Aut}_k k[[z]]$
In the lifting process then I_y is lifted as $I_y \subset \mathrm{Aut}_R \hat{\mathcal{O}}_{\mathcal{C},y} \simeq \mathrm{Aut}_R R[[Z]]$.

Results.

· If $(|G|, p) = 1$ the answer is yes by Grothendieck (SGA I). In fact he proves that the answer is yes under the condition that the ramification in $C \xrightarrow{f_s} C/G$ is tame; i.e. I_y has prime to p order for any $y \in C$.

So the main problem occurs when the ramification is wild. Recall that any p-group can occur as an inertia group moreover in infinitely many ways.

· If $|G| > 84(g(C)-1)$ the answer is no due to a trivial contradiction using Hurwitz bound for automorphism group in char 0.

· If G is cyclic of order pe(resp. $p^2 e$) with $(e,p) = 1$, the answer is yes for R large enough namely $W(k)[\zeta_{(1)}]$(resp. $W(k)[\zeta_{(2)}]$ miss a parenthesis before where $\zeta_{(n)} \in K^{\mathrm{alg}}$ is a primitive p^n-th root of 1. See [O-S-S] resp. [G-M1].

We have seen that the global lifting problem induces the:

Local lifting problem. *Let $G \underset{\text{finite}}{\subset} \mathrm{Aut}_k k[[z]]$; can we find R as above and a commutative diagram*



I. Local global principle

We prove the following

Theorem([G-M1]). *The global lifting problem over R for (C,G) is equivalent to the local lifting problem over R for any y $(\hat{\mathcal{O}}_{C,y}, I_y \subset \mathrm{Aut}_k \hat{\mathcal{O}}_{C,y})$ where y runs the branch locus of $C \xrightarrow{f_s} C/G$.*

Proof. Sketch

We use rigid analytic geometry

First Step: Lift D/k as $\mathcal{D}/R$ (SGA I). Let $B_r :=$ the branch locus for f_s.



lift $\bar{A} \hookrightarrow \bar{B}$
in $A \hookrightarrow B$
finite étale G-cover
of affinoid
(This is possible
via "Hensel" lemma)



The lifting is unique up to isomorphism. In fact the morphism $f_{(k)}$ extends (use Krasner-lemma) at the boundary of the formal fiber at x. And the germ of prolongation is unique so Galois I_y-cover.



The main problem is so to compactify in a Glois way.
The local lifting problem says that we have Galois cover of open disks lifting

$$\operatorname{Spec} k[[z]] \to \operatorname{Spec} k[[z]]^{I_y} = k[[t]]$$



Second Step: We prove a gllueing lemma using Newton's theorem (the main point is that the extension $k((z))/k((t))$ is separable !). Conclude to the algebraicaly using GAGA.

II. Local lifting for order p or p^2 automorphisms

It is possible to "lift" $k[[z]]/k[[z]]^G = k[[t]]$ in a Galois way as $A/R[[T]] = A^G$ for some A finite normal over $R[[T]]$(see Garuti [Ga]). The main problem is here to do this with a smooth A/R i.e. with good reduction over R.

For this purpose we need a numerical criteria for smoothness which is a particular case of a formula due to K.Kato (Duke M.J. 81 [Ka]).

Theorem(Kato). *Let $A/R[[T]]$ be a finite normal local ring such that A/π is reduced. We assume that $k((z)) = \mathrm{Fr}(A/\pi)/k((t))$ is separable and $\mathrm{rank}_{R[[T]]}A = \dim_{k((t))} k((z)) = n$. Let d_s be the degree of the different in the extension $k((z))/k((t))$. and d_η be the degree of the different in the extension $A \otimes_R K/R[[T]] \otimes_R K$. Then A is smooth over R i.e. $A \simeq R[[Z]]$ iff $d_\eta = d_s$.*

Application.

Order p^n case: We have Sekiguchi-Suwa [S-S] theory which shows the existence of a generic way to deform geometrically and in a Galois way a p^n-cyclic cover over $\mathrm{Spec}\, k((z))$ $\to \mathrm{Spec}\, k((z))^{<\sigma>}$. In order to deform in a smooth way we should be able to calculate differents; this is the main obstacle at the present to go further than $n = 2$.

Order p case([O-S-S]):

Let σ be an order p k-autmorphism of $k[[z]]$. By Artin-Schreier theory (abbreviation A-S) there exists $x \in k((z))$ such that $\sigma(x) = x + 1$ and $x^p - x = f(t)$. After a translation on x and a change of parameter we can take

$$\begin{cases} \sigma(x) = x + 1 \\ x^p - x = \frac{1}{t^m} \end{cases}$$

for some m, $(m, p) = 1$; we call $m + 1$ the Hasse-conductor or Weierstrass degree for σ.

Note that $z := x^{-\frac{1}{m}}$ is a parameter and that $\sigma(z) = z(1 + z^m)^{-\frac{1}{m}} = z(1 - \frac{1}{m}z^m + \cdots)$. By O-S-S theory we deform the A-S isogeny

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/p\mathbb{Z} & \longrightarrow & \mathbb{G}_a & \longrightarrow & \mathbb{G}_a & \longrightarrow & 0 \\ & & & & x & \mapsto & x^p - x & & \end{array}$$

to

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/p\mathbb{Z} & \longrightarrow & \mathcal{G}^{(\lambda)} & \longrightarrow & \mathcal{G}^{(\lambda^p)} & \longrightarrow & 0 \\ & & & & x & \mapsto & \frac{(\lambda x+1)^p-1}{\lambda^p} & & \end{array}.$$

Let $R = W(k)[\zeta_{(1)}]$, $\lambda = \zeta_{(1)} - 1$. Let A be the integral closure of $R[[T]]$ in $F := \mathrm{Fr}R[[T]](X)$ where $\frac{(\lambda X+1)^p-1}{\lambda^p} = \frac{1}{T^m}$ (this is a p-cyclic cover of $\mathbb{P}^1$ ramified in $T = 0$ and $T^m = -\lambda^p$ inside the open disc $|T| > 1$. So the generic different is

$$d_\eta = (m + 1)(p - 1)$$

which is known to be the different in the extension $k((z))/k((t))$. It is possible here to show that $Z := X^{-\frac{1}{m}}$ is a parameter for the open disc over R then $\sigma(Z) := \zeta Z(1 + (\zeta Z)^m)^{-\frac{1}{m}}$ lifts σ.

Remarks. 1. We could consider a lifting

$$\frac{(\lambda X+1)^p-1}{\lambda^p}=\frac{1}{(T-t_1)\cdots(T-t_m)}$$

with $t_i \in \pi R$, 2 by 2 distinct then $d_\eta = 2m(p-1) > d_s$. This cover has bad reduction and induces at the special fiber a cover with a cusp.



2. We will see in the next lecture that there are other non equivalent ways to lift an order p-automorphism.

Order p^2 case: Artin-Schreier-Witt theory (abbreviation A-S-W) gives

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}/p^2\mathbb{Z} & \longrightarrow & W_2 & \xrightarrow{F-1} & W_2 & \longrightarrow & 0 \\ & & & & (x_1,x_2) & \mapsto & (x_1^p-x_1, x_2^p-x_2-c(x_1^p,-x_1) & & \end{array}$$

where $c(x_1^p,-x_1) := \frac{(x_1^p-x_1)^p-(x_1^{p^2}+(-x_1)^p)}{p} \in \mathbb{Z}[x_1]$. After some translation we can find an Artin-Schreier representant

$$\begin{cases} x_1^p - x_1 = \frac{1}{t^{m_1}} & (m_1,p)=1 \\ x_2^p - x_2 - c(x_1^p,-x_1) = f_2\left(\frac{1}{t}\right) & \end{cases}$$

in such a way that $f_2\left(\frac{1}{t}\right) \in k\left[\frac{1}{t}, x_1^p - x_1\right]$ is written in a way which gives the different of the extension(see [G-M1] lemma 5.1).

Our main contribution was first to give an explicit formula for deforming A-S-W as the existence was proved in [S-S]; and secondly to provide in each case a lifting with the good different. We prove

Theorem([G-M1]). *Let $\lambda := \zeta_{(1)} - 1$, $\pi := \zeta_{(2)} - 1$, $\mu = \mathrm{Log}_p(1+\pi)$ the truncated logarithm in degree p (ζ_n is a compatible system of p^n th roots of 1).*

$$(y_1,y_2) = \left(\frac{(\lambda x_1+1)^p-1}{\lambda^p}, \frac{1}{\lambda^p}\{(\lambda x_2+\mathrm{Exp}_p\mu x_1)^p - (\lambda x_1+1)\mathrm{Exp}_p\mu^p y_1\}\right)$$

lifts A-S-W isogeny. (*)

Application:

$$\begin{cases} y_1 = \frac{1}{T^m} \\ y_2 = 0 \end{cases}$$

lifts the cover above with $f_2\left(\frac{1}{t}\right) = 0$.

In order to prove the smoothness it is sufficient to proceed by stage in the p^2-cyclic extension. We have only to look at the second stage above the open disc $|x_1| > 1$ ($x_1^{-\frac{1}{m_1}}$ is a parameter for the open disk in the first stage). It is easy to bound the generic different

(*)In particular $[(\mathrm{Exp}_p\mu x_1)^p - (\lambda x_1+1)\mathrm{Exp}_p\mu^p y_1]/\lambda^p$ lifts the cocycle $c(x_1^p,-x_1)$. In proving this we have to prove the congruence $p\mu - \lambda - p\frac{\mu^p}{\lambda^{p-1}} \equiv 0 \mod \pi^{p^2+1}$.

$d_\eta \leq [1 \text{ (for } \infty \text{ pt)} + 1 \text{ (for } x_1 = -\frac{1}{\lambda}) + d^0_{x_1}\mathrm{Exp}_p(\mu^p y_1)](p-1) = (2+p(p-1))(p-1)$. The special differnt is $d_s = (m_2+1)(p-1)$ where $m_2 = d^0 c(x_1^p, -x_1) = p(p-1)+1$. So $d_\eta \leq d_s$ and as always $d_\eta \geq d_s$ we conclude.

The general p^2-cyclic extension needs a good choice of f_2 in such a way we can read the different and for the lifting other involved formulas.

III. Other p-groups and Galois Inverse type conjecture

a) Obstructions to local lifting for $G = (\mathbb{Z}/p\mathbb{Z})^2$.

Theorem([G-M1]). *Let $G =< \sigma_1, \sigma_2 > \simeq (\mathbb{Z}/p\mathbb{Z})^2 \subset \mathrm{Aut}_k k[[z]]$ then we have the picture with $p+1$ intermediate p-cyclic extensions with Hasse conductors in the lower level $m_1+1 \leq m_2+1 = m_3+1 = ... = m_{p+1}+1$ and in the upper level $m_1' = m_2 p - m_1(p-1)$ and $m_i' = m_1$ for $i > 1$.*



If there is a local lifting then $p|m_1+1$ (this is stronger than Hasse-Arf congruence). Moreover, if



are liftings then the branch locus for $f_{1(k)}$ and $f_{2(k)}$ meet in $\frac{m_1+1}{p}(p-1)$ and this condition is an iff condition.

The proof is elementary, the congruence can be seen as follows. We know that $m_1'+1$ is the number of fix points for a lifting for σ_1, then the lifting of σ_2 acts freely on the $m_1'+1$ points (because the inertia groups are at most cyclic of order p) and so $p|m_1'+1$ and so $p|m_1+1$. The end is an application of Kato's criterion.

Remark The congruences above have been generalised by J.Bertin [Be] for general groups. Moreover he proves that the Artin representation can be expressed in term of the regular representation and a representation of permutation (given by the action on the ramification locus) in particular it is rational.

b) We are ended this lecture by what we call :

"Inverse Galois type conjecture".

Let G a group then there is an R as above and a commutative diagram



This is in some respect a strong form of Abhyankar's conjecture ([Ra]) and we prove

Theorem([M]). *The Inverse Galois type conjecture is true for abelian group of type* $(p,\dots,p)$.

Proof. In case $p=2$, $G=(\mathbb{Z}/2\mathbb{Z})^3$. The general case will be consider in next lecture.

Lemma. *The elliptic curve*

$$y^2=(1+\alpha_1 x)(1+\alpha_2 x)(1+(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2 x)$$

where $\alpha_i\in\mathbb{Z}_2^{\mathrm{ur}}$ *are such that* $\alpha_1\alpha_2(\alpha_1+\alpha_2)\neq 0 \mod 2$, *has potentially good reduction at* 2, *an equation for the special fiber is*

$$z^2-z=\bar{\alpha_1}\bar{\alpha_2}(\bar{\alpha_1}+\bar{\alpha_2})s^3$$

(a supersinguler elliptic curve as an étale 2*-cyclic cover of* $\mathbb{A}^1$*).*

Proof. Write

$$y^2=1+(\alpha_1+\alpha_2+(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2)x \\ +(\alpha_1\alpha_2+(\alpha_1+\alpha_2)(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2)x^2+\alpha_1\alpha_2(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2x^3.$$

Let $\gamma=\alpha_1+\alpha_2+\sqrt{\alpha_1}\sqrt{\alpha_2}$ then

$$y^2=(1+\gamma x)^2+\alpha_1\alpha_2(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2x^3.$$

Call

$$\begin{cases} x=2^{\frac{2}{3}}s \\ y=1+\gamma x-2z \end{cases}$$

///

For the theorem consider the 3 elliptic curves

$$y_1^2=(1+\alpha_1 x)(1+\alpha_2 x)(1+(\sqrt{\alpha_1}+\sqrt{\alpha_2})^2x)$$
$$y_2^2=(1+\alpha_2 x)(1+\alpha_3 x)(1+(\sqrt{\alpha_2}+\sqrt{\alpha_3})^2x)$$
$$y_3^2=(1+\alpha_3 x)(1+\alpha_1 x)(1+(\sqrt{\alpha_3}+\sqrt{\alpha_1})^2x)$$

where $\alpha_i\in\mathbb{Z}_2^{\mathrm{ur}}$ and if $A_1=\overline{\alpha_1\alpha_2}(\overline{\alpha_1+\alpha_2})$, $A_2=\overline{\alpha_2\alpha_3}(\overline{\alpha_2+\alpha_3})$, $A_3=\overline{\alpha_3\alpha_1}(\overline{\alpha_3+\alpha_1})$ then if

$$\prod(\varepsilon_1A_1+\varepsilon_2A_2+\varepsilon_3A_3)\neq 0 \qquad (\varepsilon_1,\varepsilon_2,\varepsilon_3)\in(\mathbb{Z}/2\mathbb{Z})^3-(0,0,0)$$

they have simultanously good reduction and the normalisation of the composition over $\mathbb{P}^1$ $(\mathbb{Q}_2^{\mathrm{ur}}(x))$ is a $(\mathbb{Z}/2\mathbb{Z})^3$-cover with good reduction and gives mod 2 the $(\mathbb{Z}/2\mathbb{Z})^3$ étale cover of the affine line generated by the three equations

$$\begin{cases} z_1^2+z_1=A_1s^3 \\ z_2^2+z_2=A_2s^3 \\ z_3^2+z_3=A_3s^3 \end{cases}.$$

Calculate $d_s=4(2-1)(1+2+2^2), d_\eta=\sharp\{\text{branch points}\}2^2(2-1)$
and the brach locusis $\{\infty,\alpha_1,\alpha_2,\alpha_3,(\sqrt{\alpha_i}+\sqrt{\alpha_j})_{i\neq j}\}$ so $d_s=d_\eta$. ///

Remark. (see [G-M2] II. 3.3.3) The "Inverse Galois type conjecture" is true for p^n-cyclic groups. The proof uses Lubin-Tate formal groups, we will indicate the method in the next lecture.

IV. References

[Be] J. Bertin, *Obstructions locales au relèvement de revêtements galoisiens de courbes lisses*, C.R. Acad. Sci. Paris, 326, Série I, (1998), 55–58.

[Ga] M. Garuti, *Prolongement de revêtement galoisiens en géométrie rigide*, Compositio Math., **104** (1996), 305-331.

[G-M 1] B. Green, M. Matignon, *Liftings of Galois Covers of Smooth Curves*, Compositio Math., to appear.

[G-M 2] B. Green, M. Matignon, *Order p automorphisms of the open disc of a p-adic field*, Journal of AMS, to appear.

[K] K. Kato (with collaboration of T. Saito), *Vanishing cycles, ramification of valuations, and class field theory*, Duke Math. J. **55**, 3 (1987), 629–659.

[M] M. Matignon,*p-groupes abéliens de type* $(p,...,p)$ *et disques ouverts p-adiques*, Prépublication 83 (1998), Laboratoire de Mathématiques pures de Bordeaux.

[O-S-S] F. Oort, T. Sekiguchi, N. Suwa, *On the deformation of Artin-Schreier to Kummer*, Ann. scient. Éc. Norm. Sup., 4^e série, t. **22** (1989), 345–375.

[Ra 1] M. Raynaud, *Revêtements de la droite affine en caractéristique* $p > 0$ *et conjecture d'Abhyankar*, Invent. Math. **116** (1994), 425–462.

[S-S 1] T. Sekiguchi, N. Suwa, *On the unified Kummer-Artin-Schreier-Witt theory*, (Preprint series), CHUO MATH NO.41, (1994).

[S-S 2] T. Sekiguchi, N. Suwa *Théories de Kummer-Artin-Schreier-Witt*, C. R. Acad. Sci. Paris, t. **319**, Série I (1994), 105–110.

M. Matignon
UPRS-A 5467 CNRS
351, cours de la Libération
France

Mathématiques Pures de Bordeaux
Université de Bordeaux I
33405 – Talence, Cedex
email: matignon@math.u-bordeaux.fr