

TD Elliptic Curves 4, 5 and 6

Damien Robert

29 October 2020

1 Rational functions

Exercise 1.1.

- Show that if $y_P \neq 0$, a uniformiser at P is $x - x_P$;
- Show that if $y_P = 0$, a uniformiser at P is y ;
- Show that a uniformiser at 0_E is x/y . Deduce that $v_{0_E}(g) = -\deg(g)$ where $\deg(x) = 2$ and $\deg(y) = 3$.

Exercise 1.2.

- Let E be the elliptic curve $y^2 = x^3 - x$ over \mathbb{Q} .
- Compute the order and the value of the rational function x/y at $P = (0, 0)$.
 - Compute the order and the value of the rational function $\frac{y+x-1}{x-1}$ at $P = (1, 0)$.
 - Compute the order and the value of the rational function $\frac{x^3}{2y^2}$ at 0_E .
 - Compute the order and the value of the rational function $\frac{x^2+y}{xy}$ at 0_E .

Exercise 1.3.

- Let E be the elliptic curve $y^2 = x^3 + 6$ over \mathbb{F}_{11} .
- Compute the order and the value of the rational function $-2x - y + 4$ at $P = (-2, 8)$.
 - Compute the order and the value of the rational function $x + 3y$ at $P = (-2, 8)$.

Exercise 1.4.

- Let E be an elliptic curve and P a point of E which is not a Weierstrass point. Let $r(x, y) = \frac{y-y_P}{x-x_P}$. Compute the order and the value of r at P .
- Let $P = (x_P, 0)$ a Weierstrass point of E . Compute the order and the value of $x - x_P$ at P .

Exercise 1.5.

- Let E be the elliptic curve $y^2 = x^3 - 7x + 6$ over \mathbb{Q} .
- Compute the order and the value of the rational function $x^2 + y - 1$ at $P = (1, 0)$.
 - Compute the order and the value of the rational function $x^2 + y^2 - 1$ at $P = (1, 0)$.
 - Compute the order and the value of the rational function $x^3 - (x^2 - 1)y - 1$ at $P = (1, 0)$.

2 Divisors

Exercise 2.1.

- Let E be the elliptic curve $y^2 = x^3 + x + 3$ over \mathbb{F}_{11} , $P = (1, 4)$, $Q = (3, 0)$, $R = (0, 6)$, $S = (1, 7)$. Compute the equation and the associated divisor of
- the line going through S and $-S$;

3 Pairings

- the tangent at R ;
- the line going through P and Q ;
- the tangent at P ;
- the tangent at Q .

Exercise 2.2. Let E be the elliptic curve $y^2 = x^3 + x + 1$ over \mathbb{F}_7 and $P = (0, 1)$. Check that $D = 5[P] - [5P] - 4[0_E]$ is principal and compute a function whose associated divisor is D .

Exercise 2.3.

- Write a function which takes for input three points $P_1, P_2, Q \in E$ and outputs $\mu_{P_1, P_2}(Q)$ where μ_{P_1, P_2} is a function with divisor $[P_1] + [P_2] - [P_1 + P_2] - [0_E]$.
- Write a function which takes for input $\ell \in \mathbb{N}$ and two points $P, Q \in E$ and outputs $f_{\ell, P}(Q)$ where $f_{\ell, P}$ is a function with divisor $\ell[P] - [\ell P] - (\ell - 1)[0_E]$.

3 Pairings

Exercise 3.1. The Weil pairing is a \mathbb{Z} -bilinear application, alternate and non degenerate :

$$e_m : E(\overline{\mathbb{F}_p})[m] \times E(\overline{\mathbb{F}_p})[m] \longrightarrow \mu_m(\overline{\mathbb{F}_p})$$

where $\mu_m(\overline{\mathbb{F}_p})$ is the multiplicative group of m -th roots of unity in $\overline{\mathbb{F}_p}$.

The pairings e_m are compatibles between each others : let m' be another integer prime to p , and $P \in E[m]$, $Q \in E[mm']$. Then

$$e_{mm'}(P, Q) = e_m(P, m'Q).$$

1. What does the bilinearity of e_m means?
2. What does alternate means for e_m ?
3. What does non degenerate means for e_m ?
4. Show that e_m is antisymmetric, which means that

$$e_m(Q, P) = e_m(P, Q)^{-1}$$

for all tuple $(P, Q) \in E[m]^2$.

5. To which group is $\mu_m(\overline{\mathbb{F}_p})$ isomorphic to? Show that there exists an integer k such that

$$\mu_m(\overline{\mathbb{F}_p}) = \mu_m(\mathbb{F}_{p^k}).$$

What arithmetic condition does k satisfy? What is the smallest k possible?

6. Let P and Q two points de m -torsion. Determine a relation between the order of $e_m(P, Q)$, and the orders of P and Q .
7. Let $P \in E(\overline{\mathbb{F}_p})$ be a (primitive) point of order m . Show that there exists another (primitive) point Q of order m such that $e_m(P, Q)$ is a primitive m -root of unity.

8. Let R be a multiple of P , and Q as in the previous question. We try to determinate the discrete logarithm, meaning an integer ℓ such that $R = \ell P$. Let ℓ_2 be an integer such that $e_{W,m}(R, Q) = e_{W,m}(P, Q)^{\ell_2}$. Show that $R = \ell_2 \cdot P$. Deduce a procedure that determines the discrete logarithm over E from the discrete logarithm over a finite field.

Exercise 3.2. Let r be a prime number. Compute the smallest k (the embedding degree) such that \mathbb{F}_{q^k} contains the r -th roots of unity.

Let E be an elliptic curve over \mathbb{F}_q . Recall that $\#E = q + 1 - t$ where t is the trace of the Frobenius. Let $r \mid \#E$, and k the corresponding embedding degree. Show that k is the order of $t - 1$ modulo r .

- Compute the embedding degree of $y^2 = x^3 - x$ over several prime numbers. (Meaning the embedding degree of the group $E(\mathbb{F}_p)$ over several p).
- Compute the embedding degree of $y^2 = x^3 + 1$ over several prime numbers.

Exercise 3.3. Let E be the elliptic curve defined by the long Weierstrass coefficients $[1, 0, 0, -4, -1]$ over \mathbb{F}_{23} . E is not given by a short Weierstrass equation, but the Sage function `weil_pairing` still allows to compute the Weil pairing.

- Let $P = (5, 8)$ and $Q = (-2, 1)$ in $E(\mathbb{F}_{23})$. Check that P is of order 4 and Q of order 2. Compute $e_4(P, Q)$ and deduce that Q is not a multiple of P .
- What is the smallest integer k such that \mathbb{F}_{23^k} contains the 4-th roots of unity?
- Check the help of the function `E.division_polynomial` which allows to compute the ψ_n , the n -th division polynomial of E . Compute the polynomial ψ_4 of 4-division of E .
- Looking at the factorisation of ψ_4 , check that all points of 4-torsion are defined over \mathbb{F}_{23^2} . (Hint : also use the functions `E.is_x_coord` and `E.lift_x`).
- Find a point R of order 4 in $E(\mathbb{F}_{23^2})$ such that $e_4(P, R)$ is a primitive 4-th root of unity.

Exercise 3.4.

- Recall the formulae to compute the Weil and Tate pairings from the functions $f_{\ell,P}(Q)$ defined in exercise 2.3.
- Write a function that computes the Tate and Weil pairings. Compare to `tate_pairing` and `weil_pairing`
- Test some examples on the curve $y^2 = x^3 + 5$ over

$$\mathbb{F}_{16030569034403128277756688287498649515636838101184337499778392980116222246913}.$$

(Check your result with `tate_pairing` and `weil_pairing`.) What is the embedding degree of this curve?