

Computation of the maximal degree of the inverse of a cubic automorphism of the affine plane with Jacobian 1 via Gröbner Bases

M. FOURNIÉ[†], J.-Ph. FURTER[‡] AND D. PINCHON[†]

[†]UMR CNRS 5640, University of Paul Sabatier, 118 route de Narbonne, 31062 Toulouse, France.

[‡]UMR CNRS 128, Ecole Normale Supérieure de Lyon, 46 allée d'Italie, 69007 Lyon, France.

(Received 10 October 1997)

In this paper we propose to compute the maximal degree of the inverse of a cubic automorphism of the affine plane with Jacobian 1 via Gröbner Bases. This degree is equal to 9 and we give coefficients of the inverse.

1. Introduction

If k is any commutative ring, $k[X, Y]$ will denote the algebra of polynomials with coefficients in k in the indeterminates X, Y and $\mathbb{A}_k^2 = \text{Spec } k[X, Y]$ the affine plane over k . A k -endomorphism f of \mathbb{A}_k^2 will be identified with its coordinate functions $f = (f_1, f_2)$ where f_i ($i = 1, 2$) belongs to $k[X, Y]$. We define the Jacobian of f by $\text{Jac}(f) = \frac{\partial f_1}{\partial X} \frac{\partial f_2}{\partial Y} - \frac{\partial f_1}{\partial Y} \frac{\partial f_2}{\partial X}$ and the degree of f by $\deg(f) = \max_{1 \leq i \leq 2} \deg(f_i)$.

Let d be a nonnegative integer and f an endomorphism of $\mathbb{A}_{\mathbb{C}}^2$ whose degree is less than or equal to d . The Jacobian Conjecture in degree d (CJ(d)) states that f is invertible if and only if its Jacobian is a nonzero constant.

Let C_d be the smallest integer C such that if k is a \mathbb{Q} -algebra and f a k -automorphism of \mathbb{A}_k^2 satisfying $\text{Jac}(f) = 1$ and $\deg(f) \leq d$, then we have $\deg(f^{-1}) \leq C$.

H. Bass has proven the following result in (Bass, 1983):

THEOREM 1.1. *The three following assertions are equivalent :*

- (i) *CJ(d) is true,*
- (ii) *if k is any \mathbb{Q} -algebra and f any k -endomorphism of \mathbb{A}_k^2 whose degree is less than or equal to d , then f is invertible if and only if $\text{Jac}(f)$ is an invertible element of $k[X, Y]$,*
- (iii) *$C_d < \infty$.*

If k is a reduced \mathbb{Q} -algebra and f a k -automorphism of \mathbb{A}_k^2 satisfying $\text{Jac}(f) = 1$ and

$\deg(f) \leq d$, it follows from a formula of O. Gabber (see (Bass, Connel, Wright, 1982) and (Cheng, Wang, Yu, 1994)) that $\deg f^{-1} = \deg f$. What happens if k is not reduced ? Is it true that $C_d = d$ (see Question 2.19 of the paper (van den Essen, 1991)) ?

A negative answer to this question is given in (Furter, to appear) where it is proven that $C_d \geq d + 1$ as soon as $d \geq 3$. Also, T. T. Moh has proven that $CJ(d)$ is true when $d \leq 100$ (see (Moh, 1983)). It then follows from Theorem 1.1 that C_d is finite for $d \leq 100$.

We could easily check that $C_1 = 1$. Theorem 2 of (Furter, to appear) shows us that $C_2 = 2$. The purpose of this paper is to establish the following result :

THEOREM 1.2. $C_3 = 9$.

As far as we know, there is no explicit upper bound for C_d when $d \geq 4$ and there is even no conjectured upper bound. An investigation of C_4 seems rather important to us in order to get some insight in the behaviour of C_d in general.

2. Computation of C_3

Let k be the algebra of polynomials with coefficients in \mathbb{Q} in the indeterminates $a_1, a_2, a_3, b_1, b_2, b_3, b_4, c_1, c_2, c_3, d_1, d_2, d_3, d_4$ and let $f = (f_1, f_2)$ be the k -endomorphism of \mathbb{A}_k^2 whose coordinate functions are

$$\begin{cases} f_1 = X + a_3X^2 + a_2XY + a_1Y^2 + b_4X^3 + b_3X^2Y + b_2XY^2 + b_1Y^3, \\ f_2 = Y + c_3X^2 + c_2XY + c_1Y^2 + d_4X^3 + d_3X^2Y + d_2XY^2 + d_1Y^3. \end{cases}$$

Let $g = (g_1, g_2)$ be the formal inverse of f . The formal series g_1 and g_2 have expressions of the form

$$\begin{cases} g_1 = X + \sum_{(i,j) \in \mathbb{N}^2, i+j \geq 2} x_{i,j} X^i Y^j, \\ g_2 = Y + \sum_{(i,j) \in \mathbb{N}^2, i+j \geq 2} y_{i,j} X^i Y^j, \end{cases}$$

where $x_{i,j}, y_{i,j}$ belong to k .

The Jacobian of f is a polynomial with coefficients in k in the indeterminates X, Y . Its constant term is equal to 1 and we could check that its other nontrivial coefficients

are equal to

$$\left\{ \begin{array}{l} -3 b_3 d_4 + 3 b_4 d_3, \\ -6 b_2 d_4 + 6 b_4 d_2, \\ -9 b_1 d_4 - 3 b_2 d_3 + 3 b_3 d_2 + 9 b_4 d_1, \\ -6 b_1 d_3 + 6 b_3 d_1, \\ -3 b_1 d_2 + 3 b_2 d_1, \\ -3 a_2 d_4 + 2 a_3 d_3 - 2 b_3 c_3 + 3 b_4 c_2, \\ -6 a_1 d_4 - a_2 d_3 + 4 a_3 d_2 - 4 b_2 c_3 + b_3 c_2 + 6 b_4 c_1, \\ -4 a_1 d_3 + a_2 d_2 + 6 a_3 d_1 - 6 b_1 c_3 - b_2 c_2 + 4 b_3 c_1, \\ -2 a_1 d_2 + 3 a_2 d_1 - 3 b_1 c_2 + 2 b_2 c_1, \\ d_3 - 2 a_2 c_3 + 2 a_3 c_2 + 3 b_4, \\ 2 d_2 - 4 a_1 c_3 + 4 a_3 c_1 + 2 b_3, \\ 3 d_1 - 2 a_1 c_2 + 2 a_2 c_1 + b_2, \\ c_2 + 2 a_3, \\ 2 c_1 + a_2. \end{array} \right.$$

Let I be the ideal of k generated by the 14 polynomials given above.

Let us set $\bar{k} = k/I$. By reducing all the coefficients of f modulo I , we obtain a \bar{k} -endomorphism of $\mathbb{A}_{\bar{k}}^2$ which we will denote by \bar{f} . Clearly, \bar{f} is the generic cubic endomorphism with Jacobian 1 of the affine plane with the following meaning. Let A be any \mathbb{Q} -algebra and α be any cubic A -endomorphism of \mathbb{A}_A^2 with Jacobian 1. Up to an affine change of coordinates, we can always suppose that $\alpha(0) = 0$ and $\alpha'(0) = \text{Id}$. Therefore, there exists a canonical algebra-homomorphism $\phi : \bar{k} \rightarrow A$ such that the A -endomorphism of \mathbb{A}_A^2 obtained by replacing the coefficients of \bar{f} by their image under ϕ will be equal to α . As CJ(3) is true, the endomorphism \bar{f} is an automorphism and we have clearly $C_3 = \deg(\bar{f})^{-1}$. Hence, the integer C_3 is the smallest integer C such that $x_{i,j}, y_{i,j}$ belong to I as soon as $i + j > C$.

Using a computer, we found that the smallest integer N such that $x_{i,j}, y_{i,j}$ belong to I as soon as $i + j = N$ is equal to 10. This encouraged us to believe that $C_3 = 9$ (and this already proved that $C_3 \geq 9$). Let h denote the k -endomorphism obtained from g by truncating its terms of degree bigger than or equal to 10. Then, to show that $C_3 = 9$, we only had to check that all coefficients of the endomorphism $f \circ h - \text{Id}$ of \mathbb{A}_k^2 (whose degree is $9^3 = 729$) belong to I . Indeed, denoting by $\bar{h} = (\bar{h}_1, \bar{h}_2)$ the \bar{k} -endomorphism of $\mathbb{A}_{\bar{k}}^2$ obtained by reducing the coefficients of h modulo I , the latter fact is equivalent to saying that the endomorphism $\bar{f} \circ \bar{h} - \text{Id}$ of $\mathbb{A}_{\bar{k}}^2$ is identically zero, which is well known to ensure that $\bar{h} = (\bar{f})^{-1}$.

All computations were done using computer algebra system AXIOM (see (Jenks, Sutor, 1983)).

3. Inversion formula

Let us endow $k = \mathbb{Q}[a_1, \dots, d_4]$ with the total degree-inverse lexicographical order (see (Davenport, Siret, Tournier, 1993)) for the following order of the indeterminates :

$$a_1 < a_2 < a_3 < c_1 < c_2 < c_3 < b_1 < b_2 < b_3 < b_4 < d_1 < d_2 < d_3 < d_4.$$

Considering the automorphism $(Y, X) \circ \bar{f} \circ (Y, X)$, one could easily show that the coefficient of $X^i Y^j$ in \bar{h}_2 is obtained from the coefficient of $X^j Y^i$ in \bar{h}_1 by replacing $a_1, a_2, a_3, c_1, c_2, c_3, b_1, b_2, b_3, b_4, d_1, d_2, d_3, d_4$ by $c_3, c_2, c_1, a_3, a_2, a_1, d_4, d_3, d_2, d_1, b_4, b_3, b_2, b_1$ respectively. Now we give the coefficients of \bar{h}_1 , or, to be more precise, the coefficients of h_1 reduced modulo the Gröbner basis of I (see (Davenport, Siret, Tournier, 1993)).

coefficient of X^2	$\frac{1}{2}c_2$
coefficient of XY	$2c_1$
coefficient of Y^2	$-a_1$

Coefficients of degree 2

coefficient of X^3	$\frac{1}{2}b_4 + \frac{1}{2}d_3$
coefficient of X^2Y	d_2
coefficient of XY^2	$-\frac{1}{2}b_2 + \frac{3}{2}d_1$
coefficient of Y^3	$-b_1$

Coefficients of degree 3

coefficient of X^4	$\frac{1}{8}c_2b_4 - \frac{1}{2}c_3d_2 + \frac{3}{8}c_2d_3 - \frac{1}{2}c_1d_4$
coefficient of X^3Y	$c_1b_4 - 2c_3d_1 + c_1d_3$
coefficient of X^2Y^2	$-\frac{3}{2}(a_1b_4 + c_2d_1 + a_1d_3)$
coefficient of XY^3	$\frac{1}{3}c_1b_2 - 3c_1d_1 - \frac{4}{3}a_1d_2$
coefficient of Y^4	$c_1b_1 + \frac{1}{4}a_1b_2 - \frac{1}{4}a_1d_1$

Coefficients of degree 4

coefficient of X^5	$\frac{3}{4}b_4^2 + \frac{1}{4}d_3^2 + \frac{3}{4}b_3d_4 - \frac{1}{4}d_2d_4$
coefficient of X^4Y	$\frac{3}{4}b_3b_4 + \frac{1}{4}b_3d_3 + \frac{3}{4}d_2d_3 + 2b_2d_4 - \frac{3}{4}d_1d_4$
coefficient of X^3Y^2	$\frac{1}{2}b_2b_4 - \frac{3}{2}b_3d_2 + \frac{1}{2}d_2^2 + 2b_2d_3 + \frac{3}{2}d_1d_3 + 6b_1d_4$
coefficient of X^2Y^3	$-\frac{3}{2}b_1b_4 + 2d_1d_2 + \frac{3}{2}b_1d_3$
coefficient of XY^4	$-\frac{3}{4}b_1b_3 + \frac{3}{2}d_1^2 - \frac{1}{4}b_1d_2$
coefficient of Y^5	$-\frac{1}{4}b_1b_2 - \frac{3}{4}b_1d_1$

Coefficients of degree 5

coefficient of X^6	$\frac{1}{8}c_2b_4^2 - \frac{1}{4}c_3d_2d_3 + \frac{1}{4}c_2d_3^2 + \frac{1}{2}c_1b_4d_4 + \frac{7}{4}c_3d_1d_4 - \frac{5}{8}c_2d_2d_4$
coefficient of X^5Y	$\frac{3}{2}c_1b_4^2 - 5c_3d_1d_3 + \frac{7}{4}c_2d_2d_3 - 2c_1d_3^2 + 12c_1b_3d_4 + 33a_1b_4d_4$ $-\frac{51}{4}c_2d_1d_4 + \frac{57}{2}c_1d_2d_4 + 19a_1d_3d_4$
coefficient of X^4Y^2	$-\frac{15}{4}a_1b_4^2 - \frac{5}{2}c_1b_3d_3 - \frac{25}{4}c_2d_1d_3 + \frac{35}{4}c_1d_2d_3 + \frac{15}{2}a_1d_3^2$ $+15c_1b_2d_4 + 5a_1b_3d_4 + \frac{75}{4}c_1d_1d_4 - \frac{15}{4}a_1d_2d_4$
coefficient of X^3Y^3	$-\frac{5}{3}a_1b_3b_4 - \frac{85}{21}c_1d_2^2 - \frac{145}{63}c_1b_2d_3 - \frac{5}{7}a_1b_3d_3 + \frac{115}{21}c_1d_1d_3$ $-\frac{215}{63}a_1d_2d_3 + \frac{5}{7}c_1b_1d_4 - \frac{55}{7}a_1b_2d_4 + \frac{40}{7}a_1d_1d_4$
coefficient of X^2Y^4	$-\frac{5}{4}a_1b_2b_4 - \frac{5}{6}c_1b_2d_2 + \frac{10}{3}a_1b_3d_2 - 5c_1d_1d_2 - \frac{5}{12}a_1d_2^2$ $-\frac{5}{2}c_1b_1d_3 - \frac{55}{12}a_1b_2d_3 - 5a_1d_1d_3 - \frac{45}{4}a_1b_1d_4$
coefficient of XY^5	$\frac{3}{23}c_1b_1b_3 + 3a_1b_1b_4 - \frac{3}{2}c_1d_1^2 + c_1b_1d_2 - a_1d_1d_2$
coefficient of Y^6	$\frac{1}{3}c_1b_1b_2 + \frac{1}{4}a_1b_1b_3 + c_1b_1d_1 - \frac{1}{4}a_1d_1^2 + \frac{1}{6}a_1b_1d_2$

Coefficients of degree 6

coefficient of X^7	$\frac{5}{72}d_3^3 + \frac{3}{8}b_3b_4d_4 - \frac{5}{12}b_3d_3d_4 - \frac{11}{24}d_2d_3d_4 + 2b_2d_4^2 + \frac{27}{8}d_1d_4^2$
coefficient of X^6Y	$\frac{7}{24}d_2d_3^2 + \frac{21}{8}b_2b_4d_4 - 4d_2^2d_4 - \frac{1}{4}b_2d_3d_4 + 12d_1d_3d_4 + 18b_1d_4^2$
coefficient of X^5Y^2	$\frac{49}{24}d_1d_3^2 + \frac{113}{24}b_2b_3d_4 - \frac{75}{4}b_1b_4d_4 + \frac{53}{6}b_2d_2d_4 + \frac{7}{4}d_1d_2d_4 - \frac{43}{4}b_1d_3d_4$
coefficient of X^4Y^3	$\frac{5}{36}d_1d_2d_3 - \frac{65}{18}b_1d_3^2 + \frac{35}{72}b_2^2d_4 + \frac{35}{3}b_1b_3d_4 + \frac{305}{8}d_1^2d_4 + \frac{445}{12}b_1d_2d_4$
coefficient of X^3Y^4	$\frac{5}{4}b_1b_3d_3 + \frac{105}{8}d_1^2d_3 + 5b_1d_2d_3 + \frac{75}{8}b_1b_2d_4 + \frac{135}{4}b_1d_1d_4$
coefficient of X^2Y^5	$-\frac{7}{24}b_2^2d_2 + \frac{63}{8}d_1^2d_2 - \frac{5}{4}b_1d_2^2 + 2b_1b_2d_3 + \frac{39}{2}b_1d_1d_3 + \frac{27}{2}b_1^2d_4$
coefficient of XY^6	$-\frac{5}{8}b_1b_2b_3 + \frac{3}{8}b_1^2b_4 + \frac{21}{8}d_1^3 - \frac{3}{8}b_1b_2d_2 - \frac{3}{2}b_1^2d_3$
coefficient of Y^7	$-\frac{1}{8}b_1b_2^2 - \frac{9}{8}b_1d_1^2 - \frac{3}{4}b_1^2d_2$

Coefficients of degree 7

coefficient of X^8	$-\frac{1}{144}c_2d_3^3 - \frac{23}{192}c_2d_2d_3d_4 + \frac{29}{48}c_1d_3^2d_4 - \frac{11}{16}c_1b_3d_4^2 - \frac{33}{16}a_1b_4d_4^2$ $+ \frac{147}{64}c_2d_1d_4^2 - \frac{31}{8}c_1d_2d_4^2 - \frac{11}{8}a_1d_3d_4^2$
coefficient of X^7Y	$-\frac{1}{9}c_1d_3^3 - \frac{11}{6}c_1b_3d_3d_4 - \frac{23}{12}c_1d_2d_3d_4 - 3a_1d_3^2d_4 + \frac{11}{2}c_1b_2d_4^2$ $+ \frac{81}{4}c_1d_1d_4^2 + 9a_1d_2d_4^2$
coefficient of X^6Y^2	$\frac{7}{18}a_1d_3^3 + \frac{21}{4}c_1b_2d_3d_4 + \frac{203}{12}a_1b_3d_3d_4 + \frac{35}{24}a_1d_2d_3d_4 - \frac{189}{4}c_1b_1d_4^2$ $- \frac{203}{4}a_1b_2d_4^2 - \frac{189}{8}a_1d_1d_4^2$
coefficient of X^5Y^3	$\frac{7}{18}a_1d_2d_3^2 + 21c_1b_2d_2d_4 + \frac{21}{2}a_1d_2^2d_4 - 63c_1b_1d_3d_4 + \frac{203}{12}a_1b_2d_3d_4$ $- 35a_1d_1d_3d_4 - \frac{609}{4}a_1b_1d_4^2$
coefficient of X^4Y^4	$\frac{35}{36}a_1d_1d_3^2 - \frac{315}{4}c_1b_1b_3d_4 - \frac{2975}{144}a_1b_2b_3d_4 - \frac{805}{16}a_1b_1b_4d_4 - \frac{945}{4}c_1d_1^2d_4$ $- \frac{315}{2}c_1b_1d_2d_4 - \frac{2905}{72}a_1b_2d_2d_4 - \frac{245}{3}a_1d_1d_2d_4 - \frac{875}{24}a_1b_1d_3d_4$
coefficient of X^3Y^5	$\frac{21}{2}c_1b_1d_2d_3 + \frac{217}{36}a_1d_1d_2d_3 + \frac{581}{18}a_1b_1d_3^2 - \frac{7}{9}a_1b_2^2d_4 + \frac{7}{3}a_1b_1b_3d_4$ $- \frac{189}{2}c_1b_1d_1d_4 - \frac{217}{4}a_1d_1^2d_4 - \frac{581}{6}a_1b_1d_2d_4$
coefficient of X^2Y^6	$-\frac{21}{4}c_1b_1b_2d_3 - \frac{7}{36}a_1b_2^2d_3 - \frac{63}{4}a_1b_1b_3d_3 - \frac{77}{24}a_1b_1d_2d_3 + \frac{189}{4}c_1b_1^2d_4$ $+ 49a_1b_1b_2d_4 + \frac{231}{8}a_1b_1d_1d_4$
coefficient of XY^7	$-3c_1b_1b_2d_2 - \frac{1}{9}a_1b_2^2d_2 - \frac{11}{6}a_1b_1d_2^2 + 9c_1b_1^2d_3 - \frac{23}{12}a_1b_1b_2d_3$ $+ \frac{11}{2}a_1b_1d_1d_3 + \frac{81}{4}a_1b_1^2d_4$
coefficient of Y^8	$\frac{9}{8}c_1b_1^2b_3 + \frac{5}{32}a_1b_1b_2b_3 + \frac{51}{32}a_1b_1^2b_4 + \frac{27}{8}c_1b_1d_1^2 - \frac{3}{8}a_1d_1^3 + \frac{9}{4}c_1b_1^2d_2$ $+ \frac{5}{16}a_1b_1b_2d_2 + \frac{3}{4}a_1b_1d_1d_2 + \frac{15}{16}a_1b_1^2d_3$

Coefficients of degree 8

coefficient of X^9	$-\frac{131}{144}d_2^2d_4^2 - \frac{131}{288}b_2d_3d_4^2 + \frac{131}{48}d_1d_3d_4^2 + \frac{131}{32}b_1d_4^3$
coefficient of X^8Y	$\frac{131}{32}b_2b_3d_4^2 - \frac{1179}{32}b_1b_4d_4^2 + \frac{131}{16}b_2d_2d_4^4 - \frac{393}{16}b_1d_3d_4^2$
coefficient of X^7Y^2	$-\frac{131}{32}b_1d_3^2d_4 - \frac{131}{32}b_2^2d_4^2 + \frac{393}{32}b_1b_3d_4^2 + \frac{393}{32}b_1d_2d_4^2$
coefficient of X^6Y^3	$\frac{917}{12}b_1b_3d_3d_4 + \frac{917}{24}b_1d_2d_3d_4 - \frac{917}{4}b_1b_2d_4^2 - \frac{2751}{8}b_1d_1d_4^2$
coefficient of X^5Y^4	$\frac{917}{8}b_1d_2^2d_4 + \frac{917}{16}b_1b_2d_3d_4 - \frac{2751}{8}b_1d_1d_3d_4 - \frac{8253}{16}b_1^2d_4^2$
coefficient of X^4Y^5	$\frac{8253}{32}b_1^2b_4d_4 + \frac{8253}{32}d_1^3d_4 + \frac{8253}{32}b_1d_1d_2d_4 + \frac{8253}{32}b_1^2d_3d_4$
coefficient of X^3Y^6	$\frac{917}{96}b_1^2d_3^2 + \frac{917}{96}b_1b_2^2d_4 - \frac{917}{32}b_1^2b_3d_4 - \frac{917}{32}b_1^2d_2d_4$
coefficient of X^2Y^7	$-\frac{131}{4}b_1^2b_3d_3 - \frac{131}{8}b_1^2d_2d_3 + \frac{393}{4}b_1^2b_2d_4 + \frac{1179}{8}b_1^2d_1d_4$
coefficient of XY^8	$-\frac{131}{16}b_1^2d_2^2 - \frac{131}{32}b_1^2b_2d_3 + \frac{393}{16}b_1^2d_1d_3 + \frac{1179}{32}b_1^3d_4$
coefficient of Y^9	$-\frac{131}{64}(b_1^3b_4 + b_1d_1^3 + b_1^2d_1d_2 + b_1^3d_3)$

Coefficients of degree 9

References

- Bass, H. (1983). The Jacobian Conjecture and inverse degree. *Arithmetic and Geometry*. Vol. II, Prog. Math., **36**, Birkhäuser, Boston, Mass., 65-75.
- Bass, H., Connel, E.H., Wright, D. (1982). The Jacobian Conjecture : reduction of degree, and formal expansion of the inverse. *Bull. Amer. Math. Soc.* 287-330.
- Cheng, C.C.-A., Wang, S.S.-S., Yu, J.-T. (1994). Degree Bounds for Inverses of Polynomial Automorphisms, *Proc. of the Amer. Math. Soc.* **120**, N° **3**, 705-707.
- Davenport, J., Siret, Y., Tournier, E. (1993). Calcul Formel - Systèmes et algorithmes de manipulations algébriques. Masson, Paris.
- van den Essen, A. (1991). Polynomial maps and the Jacobian Conjecture, Computational Aspects of Lie Group Representations and Related Topics. *Proc. of the 1990 Computational Algebra Seminar*, **84**, 29-44.
- Furter, J.-Ph. On the Degree of the Inverse of an Automorphism of the Affine Plane. *J. Pure and Applied Algebra*. accepted.
- Jenks, R.D., Sutor, R.S. (1983). Axiom - The scientific computation system. The numerical algorithms group limited, Springer Verlag.
- Moh, T.T. (1983). On the global Jacobian Conjecture and the configuration of roots. *J. Reine Angew. Math.* **340**, 140-212.