

INTRODUCTION TO FATOU COMPONENTS IN HOLOMORPHIC DYNAMICS

XAVIER BUFF AND JASMIN RAISSY

ABSTRACT. This survey is an introduction to the classification of Fatou components in holomorphic dynamics. We start with the description of the Fatou and Julia sets for rational maps of the Riemann sphere, and finish with an updated account of the recent results on Fatou components for polynomial skew-products in complex dimension two, where we focus on the key steps in the construction giving the existence of a wandering domain for a polynomial endomorphism of \mathbb{C}^2 .

1. INTRODUCTION

Discrete dynamics concerns maps $f : X \rightarrow X$ from a space to itself. One of the main goals is to understand sequences of the form

$$x_0 \in X, \quad x_1 := f(x_0), \quad x_2 := f(x_1) = f^{\circ 2}(x_0), \quad x_3 := f(x_2) = f^{\circ 3}(x_0), \dots$$

obtained by *iterating* f . The *forward orbit* of x_0 under f is

$$\mathcal{O}^+(x_0) := \bigcup_{n \geq 0} \{f^{\circ n}(x_0)\}.$$

The main problem is to understand the long term behaviour of the sequence $(x_n)_{n \geq 0}$. The answer will depend on the assumptions we make on the space X and the map $f : X \rightarrow X$.

If X is a topological space, we may wonder whether the sequence converges. If this is the case, and in addition $f : X \rightarrow X$ is a continuous map, then the limit ℓ is a fixed point of f , i.e., $f(\ell) = \ell$.

If X is compact, then thanks to the Bolzano-Weierstrass Theorem, we may extract converging subsequences. We shall denote by $\omega(x_0)$ the ω -*limit set* of x_0 , that is the set of possible limit values:

$$\omega(x_0) := \bigcap_{n \geq 0} \overline{\mathcal{O}^+(x_n)}.$$

If $f : X \rightarrow X$ is continuous, then for any $x \in X$, the ω -limit set $\omega(x)$ is f -invariant.

Exercise 1.1. Prove it.

In the rest of these notes, we will study the case where X is a complex manifold and $f : X \rightarrow X$ is a holomorphic map. We shall first consider the case where $X = \widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ is the Riemann sphere and $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ is a polynomial or a rational map.

This work is part of the C.I.M.E. Lecture Notes “Modern Aspects of Dynamical Systems” to be published in the Springer Lecture Notes in Mathematics – C.I.M.E. subseries, corresponding to the course taught by the second author in Cetraro in August 2021.

This subject has a fairly long history, with contributions by Koenigs [43], Schröder [57], Böttcher [18] in the late 19th century, and the great memoirs of Fatou [30, 31, 32] and Julia [41] around 1920. For the history of this early period we refer the reader to the book of Alexander [1]. Followed a dormant period, with notable contributions by Cremer [22, 23] (1936) and Siegel [58] (1942), and a rebirth in the 1960's (Brolin, Guckenheimer, Jakobson). Since the early 1980's, partly under the impetus of computer graphics, the subject has grown vigorously, with major contributions by Douady, Hubbard, Sullivan, Thurston, and more recently Lyubich, McMullen, Milnor, Shishikura, Yoccoz . . . Although the subject still presents many open problems, there is now a substantial body of knowledge and several books have appeared: Beardon [5], Carleson and Gamelin [21], Milnor [50] and Steinmetz [59] and the more specialized books by McMullen [48] [49]. Surveys by Blanchard [12], Devaney [25], Douady [26], Keen [42] and Lyubich [46] are also highly recommended.

There is a small collection of polynomials, for instance

$$f(z) = az + b, \quad f(z) = z^d \quad \text{and} \quad f(z) = z^2 - 2,$$

whose dynamics can be fairly easily understood.

Exercise 1.2. Assume $f(z) = az + b$ with $a \in \mathbb{C}$ and $b \in \mathbb{C}$.

- (1) Show that when $|a| \neq 1$ or when $a = 1$ and $b \neq 0$, the map

$$\mathbb{C} \ni z \mapsto \omega(z) \in \widehat{\mathbb{C}}$$

is constant.

- (2) Show that in all other cases, $\omega(z)$ depends on z and is either a finite set, or a euclidean circle.

Exercise 1.3. Assume $f(z) = z^d$ with $d \geq 2$.

- (1) Show that $\omega(z) = \{0\}$ if z belongs to the unit disk \mathbb{D} and $\omega(z) = \{\infty\}$ if $z \in \widehat{\mathbb{C}} \setminus \mathbb{D}$.
- (2) Show that
- $\omega(z) \subset S^1$ if z belongs to the unit circle S^1 ,
 - $\omega(z)$ can be equal to S^1 and
 - $\omega(z)$ can be a finite set of arbitrary cardinality.

Exercise 1.4. Assume $f(z) = z^2 - 2$.

- (1) Show that every $z \in \mathbb{C} \setminus \{0\}$ may be written $z = w + 1/w$ for some $w \in \mathbb{C}$ with $|w| = 1$ if $z \in [-2, 2]$ and $|w| > 1$ if $z \in \mathbb{C} \setminus [-2, 2]$.
- (2) Show that
- $\omega(z) = \{\infty\}$ if $z \in \widehat{\mathbb{C}} \setminus [-2, 2]$ and
 - $\omega(z) \subset [-2, 2]$ if $z \in [-2, 2]$.

Exercise 1.5. Assume $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ is a Tchebychev polynomial of degree d , i.e., such that $\cos(d\theta) = f(\cos(\theta))$. Show that $\omega(z) = \{\infty\}$ if $z \in \widehat{\mathbb{C}} \setminus [-1, 1]$ and $\omega(z) \subset [-1, 1]$ if $z \in [-1, 1]$.

Those examples are really atypical examples, since most cases exhibit *fractal* and *chaotic* behaviour, the analysis of which requires tools from complex analysis, dynamical systems, topology, combinatorics, . . .

2. FATOU SETS AND JULIA SETS

Both Fatou's and Julia's work have as a major theme that the Riemann sphere $\widehat{\mathbb{C}}$ breaks up sharply into two complementary subsets: the open Fatou set \mathcal{F}_f with orderly dynamics, and the Julia set \mathcal{J}_f on which the dynamical behaviour of f is chaotic. In practice, understanding the dynamics of a rational map has come to mean understanding the topology of the Julia set, its geometry if possible, and classifying the components of the Fatou set.

2.1. Definition. There are several possible definitions of the Fatou and Julia sets. We will adopt the one based on normal families, which seems very natural to us: a normal family of analytic maps is "well-behaved". But the very fact that it has become standard reflects the origins of holomorphic dynamics in complex analysis, and may explain why the subject has never quite entered the main stream of dynamical systems.

Definition 2.1 (Normal families). *A family $(f_\alpha : U \rightarrow \widehat{\mathbb{C}})_{\alpha \in \mathcal{A}}$ of holomorphic maps on an open subset $U \subset \widehat{\mathbb{C}}$ is normal if from any sequence $(f_{\alpha_n})_{n \geq 0}$ one can extract a subsequence converging uniformly (for the spherical metric in the range) on compact subsets of U .*

A criterion that will enable us to deal with normality is due to Montel.

Theorem 2.2 (Montel). *A family $(f_\alpha : U \rightarrow \widehat{\mathbb{C}})_{\alpha \in \mathcal{A}}$ of holomorphic maps on an open subset $U \subset \widehat{\mathbb{C}}$ which omits at least three values in $\widehat{\mathbb{C}}$ is normal.*

In particular, a bounded sequence of holomorphic maps $(f_n : U \rightarrow \mathbb{C})_{n \in \mathbb{N}}$ is normal.

Definition 2.3 (Fatou sets and Julia sets). *The Fatou set \mathcal{F}_f of a rational map $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ is the largest open subset of $\widehat{\mathbb{C}}$ on which the family of iterates $(f^{on})_{n \in \mathbb{N}}$ is normal. The Julia set \mathcal{J}_f is the complement of the Fatou set.*

Example 2.4. Example For $f(z) = z^d$, with $d \geq 2$ or $d \leq -2$, the Julia set \mathcal{J}_f is the unit circle

$$S^1 := \{z \in \mathbb{C} \mid |z| = 1\}.$$

Indeed, the family of iterates $(f^{on} : \widehat{\mathbb{C}} \setminus S^1 \rightarrow \widehat{\mathbb{C}})_{n \in \mathbb{N}}$ takes its values in $\widehat{\mathbb{C}} \setminus S^1$, hence it omits more than three values in $\widehat{\mathbb{C}}$. By Montel's theorem, it is normal and so, $\mathcal{J}_f \subseteq S^1$. To prove that $S^1 \subseteq \mathcal{J}_f$, one may argue that the sequence $(f^{o(2n)})_{n \in \mathbb{N}}$ converges uniformly on any compact subset of \mathbb{D} to the constant map 0 and on any compact subset of $\widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}$ to the constant map ∞ . As any neighborhood of a point $z \in S^1$ contains points in \mathbb{D} and points in $\widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}$, there is no neighborhood of z on which the sequence $(f^{on})_{n \in \mathbb{N}}$ is normal.

2.2. The polynomial case. In the case of a polynomial, there is a slightly different way of understanding things. In this case, ∞ is a fixed point and this fixed point has no preimage other than itself. It is an exceptional point.

Definition 2.5. *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a polynomial of degree $d \geq 2$. The filled-in Julia set of f is the set \mathcal{K}_f of points with bounded orbits:*

$$\mathcal{K}_f = \left\{ z \in \mathbb{C} \mid (f^{on}(z))_{n \in \mathbb{N}} \text{ is bounded} \right\}.$$




FIGURE 1. The Julia set of the quadratic polynomial $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ defined by $f(z) = z^2 - 1$.

This set is not only very natural for a dynamical system, but fits in very nicely with the general theory.

Proposition 2.6. *The filled-in Julia set \mathcal{K}_f is a non-empty compact set. The Julia set is the topological boundary of \mathcal{K}_f .*

Proof. Since f is of degree $d \geq 2$, $f(z)/z \rightarrow \infty$ as $z \rightarrow \infty$, and so, there exists a constant $R > 0$ such that $|f(z)| > 2|z|$ whenever $|z| > R$. Set $D_R := \{z \in \mathbb{C} \mid |z| > R\}$. Then, the nested intersection of compact sets

$$\mathcal{K}_f = \bigcap_{n \in \mathbb{N}} f^{-n}(\overline{D}_R)$$

is a non-empty compact set.

Furthermore, define U_R to be the open set $U_R := \widehat{\mathbb{C}} \setminus \overline{D}_R$. Then, the sequence $(f^{o_n})_{n \geq 0}$ converges uniformly to infinity on U_R . Clearly, the same property holds on the open set

$$\Omega_\infty = \bigcup_{n \in \mathbb{N}} f^{-n}(U_R)$$

Observe that $\Omega_\infty = \widehat{\mathbb{C}} \setminus \mathcal{K}_f$ and $\partial\Omega_\infty = \partial\mathcal{K}_f$.

We will first show that $\partial\mathcal{K}_f \subseteq \mathcal{J}_f$. Choose a point $z_0 \in \partial\Omega_\infty = \partial\mathcal{K}_f$ and assume that there is a neighborhood U of z_0 on which the family of iterates of f is normal. Then, U intersects Ω_∞ and the sequence $(f^{o_n})_{n \geq 0}$ must converge to infinity on the whole set U . This contradicts the fact that the sequence $(f^{o_n}(z_0))_{n \geq 0}$ is bounded.

We will now show that $\mathcal{J}_f \subseteq \partial\mathcal{K}_f$. As mentioned previously, the sequence (f^{o_n}) converges locally uniformly to a map which is constant and equal to ∞ outside \mathcal{K}_f . Thus, the complement of \mathcal{K}_f is contained in the Fatou set \mathcal{F}_f and $\mathcal{J}_f \subseteq \mathcal{K}_f$. Similarly, on the interior of \mathcal{K}_f , the sequence (f^{o_n}) is bounded, thus normal by Montel's theorem. As a consequence, the interior of \mathcal{K}_f is contained in \mathcal{F}_f and $\mathcal{J}_f \subseteq \partial\mathcal{K}_f$. \square

The reason why \mathcal{K}_f is called a filled-in Julia set is the following.

Definition 2.7. *A compact subset $K \subset \mathbb{C}$ is called full if it is connected and its complement is connected.*

Proposition 2.8. *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a polynomial. Each connected component of \mathcal{K}_f is full.*

Proof. Of course, any component K_0 is compact and connected. Any component U of the complement must satisfy $\partial U \subset K_0$, and if the complement is not connected at least one such component must be bounded. Then, by the maximum principle

$$\sup_U |f^{\circ n}(z)| = \sup_{\partial U} |f^{\circ n}(z)| \leq \sup_{K_0} |f^{\circ n}(z)|.$$

Hence, it is uniformly bounded, so $U \subset K_0$. □

2.3. Invariance of Fatou sets and Julia sets.

Proposition 2.9. *Both the Julia set and the Fatou set are completely invariant:*

$$f(\mathcal{J}_f) = \mathcal{J}_f, \quad f^{-1}(\mathcal{J}_f) = \mathcal{J}_f, \quad f(\mathcal{F}_f) = \mathcal{F}_f, \quad f^{-1}(\mathcal{F}_f) = \mathcal{F}_f.$$

Proof. As the Julia set is the complement of the Fatou set, it is enough to prove the statements about the Fatou set. This is an immediate consequence of the following lemma.

Lemma 2.10. *Let U be an open subset of $\widehat{\mathbb{C}}$, and let $\phi : U \rightarrow \widehat{\mathbb{C}}$ be holomorphic and non-constant. Then the family $(g_\alpha : \phi(U) \rightarrow \widehat{\mathbb{C}})_{\alpha \in \mathcal{A}}$ is normal if and only if $(f_\alpha := g_\alpha \circ \phi : U \rightarrow \widehat{\mathbb{C}})_{\alpha \in \mathcal{A}}$ is normal.*

Remark 2.11. Since ϕ is a non-constant analytic map, it is an open map. Thus $\phi(U)$ is open and the family g_α is defined on an open set.

Proof. (\Rightarrow) This is obvious. If the sequence g_{α_n} converges to g , then $f_{\alpha_n} := g_{\alpha_n} \circ \phi$ converges to $g \circ \phi$.

(\Leftarrow) Assume that the sequence $f_{\alpha_k} := g_{\alpha_k} \circ \phi$ is uniformly convergent on every compact subset of U . Let $K \subset \phi(U)$ be compact, and choose a cover of U by open subsets U_i with compact closure. Since ϕ is open, $\{\phi(U_i)\}$ is an open cover of K , and there exists a finite subcover $\{\phi(U_{i_j})\}$. The set

$$K' := \phi^{-1}(K) \cap \bigcup_j \overline{U_{i_j}}$$

is a compact subset of U such that $\phi(K') = K$. Then,

$$\sup_{z \in K} \text{dist}_{\widehat{\mathbb{C}}}(g_{\alpha_n}(z), g_{\alpha_m}(z)) = \sup_{w \in K'} \text{dist}_{\widehat{\mathbb{C}}}(f_{\alpha_n}(w), f_{\alpha_m}(w)) \rightarrow 0$$

as $n, m \rightarrow \infty$. The lemma follows. □

To prove $\mathcal{F}_f = f^{-1}(\mathcal{F}_f)$, apply this lemma to the families

$$\left\{ f^{\circ(m+1)} : f^{-1}(\mathcal{F}_f) \rightarrow \widehat{\mathbb{C}} \right\} \quad \text{and} \quad \left\{ f^{\circ m} : \mathcal{F}_f \rightarrow \widehat{\mathbb{C}} \right\}$$

and to the holomorphic map $f : f^{-1}(\mathcal{F}_f) \rightarrow \mathcal{F}_f$. Applying f to both sides of the equality $\mathcal{F}_f = f^{-1}(\mathcal{F}_f)$, we deduce that $f(\mathcal{F}_f) = \mathcal{F}_f$. □

Proposition 2.12. *The Julia and Fatou sets of the k -th iterate $f^{\circ k}$ are equal to the Julia and Fatou sets of f :*

$$\mathcal{J}_{f^{\circ k}} = \mathcal{J}_f \quad \text{and} \quad \mathcal{F}_{f^{\circ k}} = \mathcal{F}_f$$

Proof. It is enough to prove this proposition for the Fatou sets. If the family of iterates $f^{\circ n}$ is normal on an open set U , then the subfamily $f^{\circ(kn)}$ is also normal. Hence, $\mathcal{F}_f \subseteq \mathcal{F}_{f^{\circ k}}$.

Now, assume that $z \in \mathcal{F}_{f^{\circ k}}$ and let U be a neighborhood of z on which the family of iterates $f^{\circ(kn)}$ is normal. For any sequence $(n_i)_{i \in \mathbb{N}}$, there exists an integer N such that $n_i = N \pmod{k}$ for infinitely many i 's. Let us extract this subsequence which, with an abuse of notations, we still denote by n_i . By assumption, the sequence $f^{\circ(n_i - N)} : U \rightarrow \widehat{\mathbb{C}}$ is normal and we can extract a subsequence converging to a map $g : U \rightarrow \widehat{\mathbb{C}}$. As $f^{\circ n_i} = f^{\circ N} \circ f^{\circ(n_i - N)}$, we can extract a subsequence of the sequence $f^{\circ n_i} : U \rightarrow \widehat{\mathbb{C}}$ converging to $f^{\circ N} \circ g : U \rightarrow \widehat{\mathbb{C}}$. This shows that the family $(f^{\circ n} : U \rightarrow \widehat{\mathbb{C}})_{n \in \mathbb{N}}$ is normal and that $\mathcal{F}_{f^{\circ k}} \subseteq \mathcal{F}_f$. \square

3. CONJUGACY

In every part of mathematics, the *morphisms* are the maps that preserve the structure. Thus the morphisms of a dynamical system $f : X \rightarrow X$ to a dynamical system $g : Y \rightarrow Y$ are the maps sending sequences of iterates of f to sequences of iterates of g . This is just what maps $\phi : X \rightarrow Y$ such that $\phi \circ f = g \circ \phi$ do; such maps are called *semi-conjugacies* between f and g . Indeed, we see by induction that then $\phi \circ f^{\circ n} = g^{\circ n} \circ \phi$ for all $n \geq 1$: by hypothesis it is true for $n = 1$, and if it is true for $n - 1$, then

$$\phi \circ f^{\circ n} = \phi \circ f \circ f^{\circ(n-1)} = g \circ \phi \circ f^{\circ(n-1)} = g \circ g^{\circ(n-1)} \circ \phi = g^{\circ n} \circ \phi.$$

When ϕ is invertible, $\phi \circ f = g \circ \phi$ can be rewritten $f = \phi^{-1} \circ g \circ \phi$, so f and g are *conjugate*: conjugacies are the isomorphisms of dynamical systems. When studying dynamical systems, we constantly try to construct conjugacies or semi-conjugacies to simpler dynamical systems, linear ones in particular.

Definition 3.1 (Conjugate rational maps). *Two rational maps f and g are topologically (respectively analytically) conjugate if there exists a homeomorphism $\phi : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ (respectively an analytic isomorphism) such that $\phi \circ f = g \circ \phi$.*

One should think of ϕ as a change of coordinates on $\widehat{\mathbb{C}}$. The only analytic isomorphisms of $\widehat{\mathbb{C}}$ are the Möbius transformations $z \mapsto (az + b)/(cz + d)$ and the only isomorphisms of \mathbb{C} are the affine maps $z \mapsto az + b$. Being a zero or a pole of a rational map depends on the coordinates, whereas being a fixed point or a critical point of a rational map does not depend on the choice of coordinates.

Proposition 3.2. *If f and g are conjugate (topologically or analytically) by $\phi : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$, then $\phi(\mathcal{F}_f) = \mathcal{F}_g$ and $\phi(\mathcal{J}_f) = \mathcal{J}_g$.*

Proof. The family $(f^{\circ n} : U \rightarrow \widehat{\mathbb{C}})_{n \in \mathbb{N}}$ is normal if and only if the family $(\phi \circ f^{\circ n} \circ \phi^{-1} : \phi(U) \rightarrow \widehat{\mathbb{C}})_{n \in \mathbb{N}}$ is normal. \square

Example 3.3. Example We will see below that the Newton's methods

$$N_P : z \mapsto z - \frac{P(z)}{P'(z)} \quad \text{and} \quad N_Q : z \mapsto z - \frac{Q(z)}{Q'(z)}$$

are analytically conjugate as soon as $Q(z) = \lambda P(\alpha z + \beta)$.

Exercise 3.4. Let $a \neq b$ be complex numbers and set $P(z) := (z - a)(z - b)$.

- (1) Show that the Newton's method N_P is conjugate to $f : w \mapsto w^2$ via the isomorphism $z \mapsto w := (z - a)/(z - b)$.
- (2) Determine the Julia set of N_P and the set $\omega(z)$ for $z \in \widehat{\mathbb{C}} \setminus \mathcal{J}_{N_P}$.

4. PERIODIC POINT AND CRITICAL POINTS

Periodic points and critical points are important objects in holomorphic dynamics as we shall see in this section.

4.1. Periodic points.

Definition 4.1 (Multiplier at a fixed point). *A fixed point of f is a point α such that $f(\alpha) = \alpha$. The derivative of f at α is an endomorphism of the tangent line $D_\alpha f : T_\alpha \widehat{\mathbb{C}} \rightarrow T_\alpha \widehat{\mathbb{C}}$, thus a multiplication by a number $\lambda \in \mathbb{C}$. The eigenvalue λ of the endomorphism $D_\alpha f : T_\alpha \widehat{\mathbb{C}} \rightarrow T_\alpha \widehat{\mathbb{C}}$ is called the multiplier of f at α . The fixed point is called*

- superattracting if $\lambda = 0$;
- attracting if $0 < |\lambda| < 1$;
- repelling if $|\lambda| > 1$;
- indifferent if $|\lambda| = 1$, and more precisely
 - parabolic if λ is a root of 1,
 - elliptic otherwise, that is if $\lambda = e^{2\pi i\theta}$ with $\theta \in \mathbb{R} \setminus \mathbb{Q}$.

Exercise 4.2. Let α be a fixed point of a holomorphic map f .

- (1) Show that if $\alpha \neq \infty$, then the multiplier of f at α is $\lambda = f'(\alpha)$.
- (2) Show that if $\alpha = \infty$ and if $f(z) \sim az$ as $z \rightarrow \infty$, then the multiplier is $\lambda = 1/a$, whereas if $f(z) \sim az^k$ with $k \geq 2$, the multiplier is $\lambda = 0$.
- (3) Show that if f is a polynomial of degree $d \geq 2$, then ∞ is a superattracting fixed point of f .

The multiplier at a fixed point is invariant under analytic conjugacy. In other words, if $\phi : (\widehat{\mathbb{C}}, \alpha) \rightarrow (\widehat{\mathbb{C}}, \beta)$ is a local isomorphism conjugating $f : (\widehat{\mathbb{C}}, \alpha) \rightarrow (\widehat{\mathbb{C}}, \alpha)$ to $g : (\widehat{\mathbb{C}}, \beta) \rightarrow (\widehat{\mathbb{C}}, \beta)$, then the multiplier of f at α is equal to the multiplier of g at β . Indeed, the derivative of $\phi \circ f = g \circ \phi$ at α is

$$D_\alpha \phi \circ D_\alpha f = D_\beta g \circ D_\alpha \phi.$$

In other words, $D_\alpha \phi$ conjugates $D_\alpha f : T_\alpha \widehat{\mathbb{C}} \rightarrow T_\alpha \widehat{\mathbb{C}}$ to $D_\beta g : T_\beta \widehat{\mathbb{C}} \rightarrow T_\beta \widehat{\mathbb{C}}$ and the two endomorphisms have the same eigenvalue.

Definition 4.3 (Periodic points). *A periodic point of f is a fixed point of $f^{\circ p}$ for some $p \geq 1$. The smallest such integer $p \geq 1$ is called the period of α . In this case, we say that $\langle \alpha, f(\alpha), \dots, f^{\circ(p-1)}(\alpha) \rangle$ is a cycle of f .*

Proposition 4.4. *If $\langle \alpha_1, \alpha_2, \dots, \alpha_p \rangle$ is a cycle, then the multiplier of $f^{\circ p}$ at all points of the cycle is the same.*

Proof. According to the Chain Rule, the derivative of $f^{\circ p}$ at any point of the cycle is the composition of the derivatives of f , as linear transformations, along the cycle. If the derivative of f vanishes at a point of the cycle, then the multiplier is 0 at all points of the cycle. If the derivative never vanishes, then f is locally invertible at each point of the cycle and f conjugates $f^{\circ p}$ at α_i to $f^{\circ p}$ at α_{i+1} . Thus, the multiplier of $f^{\circ p}$ at α_i is equal to the multiplier of $f^{\circ p}$ at α_{i+1} . \square

Definition 4.5 (Multiplier along a cycle). *The multiplier of f along a cycle of period p is the multiplier of $f^{\circ p}$ at any point of the cycle. The cycle is superattracting (respectively attracting, indifferent or repelling) for f if the points of the cycle are superattracting (respectively attracting, indifferent or repelling) fixed points of $f^{\circ p}$.*

If $\langle \alpha_1, \alpha_2, \dots, \alpha_p \rangle$ is a cycle of period p avoiding ∞ , then its multiplier is given by the product

$$\lambda = (f^{\circ p})'(\alpha_1) = \prod_{i=1}^p f'(\alpha_i).$$

Proposition 4.6. *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a rational map, and let α be a periodic point of f . If α is superattracting or attracting, then $\alpha \in \mathcal{F}_f$. If α is repelling, then $\alpha \in \mathcal{J}_f$.*

Proof. Replacing f by $f^{\circ p}$ if necessary, we may assume that α is a fixed point of f . Conjugating by a Möbius transformation if necessary, we may assume that $\alpha = 0$. If α is (super)attracting, i.e., $|f'(\alpha)| < 1$, then there exists a bounded neighborhood U of α which is mapped into itself. In particular, for all $n \in \mathbb{N}$, $f^{\circ n}(U) \subset U$. Hence, the family $(f^{\circ n} : U \rightarrow \widehat{\mathbb{C}})_{n \in \mathbb{N}}$ is normal, which implies that $\alpha \in \mathcal{F}_f$. If α is repelling, then

$$(f^{\circ n})'(\alpha) = [f'(\alpha)]^n \xrightarrow{n \rightarrow +\infty} \infty.$$

So, the family $(f^{\circ n})_{n \in \mathbb{N}}$ cannot be normal in a neighborhood of α . \square

A cycle is superattracting if and only if it contains a point at which the derivative of f vanishes, i.e., a *critical point* of f .

4.2. Critical points and critical values. We will see that the dynamical properties of a rational map $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ are strongly related to the dynamical behaviour of the critical points of f . For example, the Julia set of a polynomial f is connected if and only if the orbit of every critical point of f is bounded.

Definition 4.7 (Critical points and critical values). *Let U and V be two Riemann surfaces and $f : (U, x) \rightarrow (V, y)$ be an analytic map which is not locally constant at x . The point x is a critical point if the derivative $D_x f : T_x U \rightarrow T_y V$ is zero. In that case, y is a critical value.*

More precisely, in local coordinates z vanishing at x and w vanishing at $y = f(x)$, the expression of f is of the form

$$z \mapsto aw^k + O(w^{k+1}) \quad \text{for some integer } k \geq 1 \text{ and some complex number } a \neq 0.$$

The integer k does not depend on the choice of coordinates and is called the *local degree* of f at x . The point x is a critical point if and only if $k \geq 2$. In that case, x is called a *critical point of f of multiplicity $k - 1$* .

Example 4.8. Example The map $z \mapsto z^3$ has a critical point at 0. The local degree of the map at 0 is 3, and the multiplicity of this critical point is 2.

We shall use the following result which is a consequence of the Riemann-Hurwitz Formula (which will not be proved in those notes).

Definition 4.9 (Euler characteristic). *If U is an open subset of $\widehat{\mathbb{C}}$ with m boundary components, the Euler characteristic of U is*

$$\chi(U) := 2 - m.$$

Theorem 4.10. *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a rational map of degree d , $V \subseteq \widehat{\mathbb{C}}$ be an open set and $U := f^{-1}(V)$. Then,*

$$\chi(U) = d \cdot \chi(V) - N$$

where N is the number of critical points of f in U , counting multiplicities.

Exercise 4.11. Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a rational map of degree d .

- (1) Prove that f has $2d - 2$ critical points counting multiplicities.
- (2) Prove that f has at least 2 distinct critical values.
- (3) Prove that if f is a polynomial, then ∞ is a critical point of multiplicity $d - 1$ and that there are $d - 1$ critical points in \mathbb{C} counting multiplicities.

5. DESCRIPTION OF THE JULIA SET

5.1. Topology of the Julia set.

Proposition 5.1. *The Julia set \mathcal{J}_f is non empty and compact.*

Proof. If the Julia set were empty, there would be a subsequence of f^{o_m} converging uniformly on $\widehat{\mathbb{C}}$; let f_0 be the limit. The rational map f_0 must have some finite degree. Since the degree is an invariant of homotopy, the approximating iterates $f^{o_{m_i}}$ must have the same degree. But the degree of $f^{o_{m_i}}$ is d^{m_i} , which leads to a contradiction.

Evidently the Fatou set \mathcal{F}_f is open, hence the Julia set \mathcal{J}_f is closed. Since $\widehat{\mathbb{C}}$ is compact, the Julia set \mathcal{J}_f is also compact. □

Proposition 5.2. *Either the Julia set \mathcal{J}_f has empty interior, or it is the entire Riemann sphere.*

Proof. Suppose $U \subset \mathcal{J}_f$ is an open subset of $\widehat{\mathbb{C}}$. The family $\{f^{o_m} : U \rightarrow \widehat{\mathbb{C}}\}$ must avoid \mathcal{F}_f , which is open and will contain more than two points if it is not empty. But this would make $\{f^{o_m} : U \rightarrow \widehat{\mathbb{C}}\}$ normal, a contradiction. Thus if the interior of \mathcal{J}_f is non empty, \mathcal{J}_f is the entire Riemann sphere. □

We have seen that the Julia set \mathcal{J}_f is completely invariant. We will now determine whether there are any other completely invariant closed sets than \mathcal{J}_f . For this purpose, we will use the Riemann Hurwitz formula.

Proposition 5.3. *Let E be a completely invariant closed set. If E is finite, then E contains at most two points.*

Proof. If E is finite, then its complement U is connected and completely invariant. Its Euler characteristic $\chi(U) = 2 - \text{card}(E)$ is finite. Let us denote by N the number of critical points of f in U counted with multiplicity. The Riemann-Hurwitz formula applied to $f : U \rightarrow U$ gives

$$\chi(U) = d \cdot \chi(U) - N.$$

Thus

$$\chi(U) = \frac{N}{d-1} \geq 0,$$

which leads to $\text{card}(E) \leq 2$. □

A union of completely invariant sets is still completely invariant, therefore the following definition makes sense.

Definition 5.4. *The exceptional set E_f is the largest finite completely invariant set.*

Remark 5.5. The grand-orbit of a point is a totally invariant set. If it is a finite set, then, by definition of the exceptional set E_f , we have $z \in E_f$.

Observe that when two rational maps $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ and $g : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ are conjugate by a homeomorphism ϕ , then the homeomorphism ϕ sends the forward (resp. backward) orbit of a point $z \in \widehat{\mathbb{C}}$ under the action of f to the forward (resp. backward) orbit of $\phi(z)$ under the action of g . Hence, f -invariant sets are mapped by ϕ to g -invariant sets. Thus, the largest finite completely f -invariant set is mapped by ϕ to the largest finite completely g -invariant set. In other words, when it is not empty, the exceptional set E_f is mapped by ϕ to the exceptional set E_g .

The exceptional set E_f is a minor irritant, which will come back to annoy us on several occasions. The following description should remove any doubts about its lack of significance.

Proposition 5.6. *Up to conjugacy, the exceptional set E_f is not empty in precisely two cases:*

- when $f(z) = z^d$ with $|d| \geq 2$, $E_f = \{0, \infty\}$ and
- when f is a polynomial but is not conjugate to $z \mapsto z^d$, $E_f = \{\infty\}$.

Proof. It is clear from the construction that E_f is completely invariant; so if it has only one point, this point must be fixed, and if it consists of two points, these must be fixed or exchanged by f . We will examine these three cases one at a time.

If E_f has just one point a , move it to ∞ , for example by making the change of coordinates $w = 1/(z - a)$. Then $f = P/Q$ is a rational map such that infinity is its only inverse image. Hence, the polynomial Q has no roots, it is constant, and f is a polynomial.

If E_f has two points a and b , both fixed, then we can put one at ∞ and the other at 0 (make the change of coordinates $w = (z - b)/(z - a)$). We see that f is a polynomial and that 0 is the only root of the polynomial equation $f(w) = 0$, so $f(w) = \lambda w^d$ for some λ . Choose a number ν such that $\nu^{d-1} = \lambda$, and make the change of coordinates $z = \nu w$. Then, it is easy to see that the expression of f becomes $z \mapsto z^d$.

If E_f has two points exchanged by f , put one at ∞ and the other at 0, and write $f = P/Q$. Since ∞ is the only inverse image of 0, we see that P has no roots, so it is a constant, which we may take to be 1. Since 0 is the only inverse image of ∞ , we see that $Q(w) = \lambda w^d$ for some $\lambda \neq 0$, so that $f(w) = 1/(\lambda w^d)$. We can do a change of coordinate $z = \nu w$, where $\nu^{d+1} = \lambda$. Again it is easy to check that the expression of f becomes $z \mapsto z^{-d}$. \square

Proposition 5.7. *The exceptional set E_f is always contained in the Fatou set \mathcal{F}_f .*

Proof. In all three cases, E_f is a union of superattracting cycles. Proposition 4.6 shows that such cycles are always contained in the Fatou set \mathcal{F}_f . \square

We can now give a new characterization of the Julia set.

Proposition 5.8. *The Julia set of a rational map is the smallest completely invariant closed set containing at least three points.*

Proof. The Julia set is a completely invariant closed set. It contains at least three points, since otherwise it would be contained in the exceptional set $E_f \subset \mathcal{F}_f$. Furthermore, if E is a completely invariant closed set containing at least 3 points, the complement $\Omega = \widehat{\mathbb{C}} \setminus E$ of E is a completely invariant open set omitting more than three points. Hence, by Montel’s theorem, the family of iterates $f^{\circ n}$ is normal on Ω . Thus $\Omega \subset \mathcal{F}_f$ and $\mathcal{J}_f \subset E$. \square

The following proposition asserts that if z does not belong to the exceptional set E_f , then the closure of the backward orbit of z ,

$$\mathcal{O}^-(z) := \bigcup_{m \geq 0} f^{-m}\{z\},$$

contains the Julia set \mathcal{J}_f .

Proposition 5.9. *For any $z \notin E_f$, we have*

$$\mathcal{J}_f \subseteq \overline{\mathcal{O}^-(z)}.$$

Proof. Take a point $z_0 \in \mathcal{J}_f$ and let U be an arbitrary neighborhood of z_0 . We must show that the set $\Omega = \bigcup_{n \in \mathbb{N}} f^{\circ n}(U)$ contains z . We will show that its complement $E = \widehat{\mathbb{C}} \setminus \Omega$ is contained in the exceptional set E_f . Indeed, Ω is a forward invariant open set which omits at most two points, since otherwise the family $f^{\circ n}$ would be normal on U . Thus, $E = \widehat{\mathbb{C}} \setminus \Omega$ is a backward invariant set. Since E is finite, it is permuted by f , so it is forward invariant. Hence, by definition of the exceptional set, it is contained in E_f . \square

Proposition 5.10. *The Julia set \mathcal{J}_f is perfect.*

Proof. Since the Julia set is closed, it suffices to prove that it has no isolated points. The Julia set \mathcal{J}_f is not finite. Thus, it contains a non-isolated point z_0 . Since the backward orbit of z_0 is dense in \mathcal{J}_f , any point in \mathcal{J}_f is non-isolated. This proves that \mathcal{J}_f is perfect, hence uncountable. \square

5.2. Julia sets and periodic points. Fatou and Julia both proved that the Julia set is the closure of the set of repelling periodic points. This “dynamical” definition of the Julia set is probably more natural to specialists of dynamical systems, and relates holomorphic dynamics to “Axiom A attractors” and hyperbolic dynamics. Fatou’s and Julia’s proofs are different but both involve results which will be proved later. Fatou’s proof is based on the fact that there are only finitely many non-repelling cycles. Julia’s proof is based on the existence of a repelling cycle.

In these notes, we will only prove the following weaker result which follows directly from Montel’s Theorem. This result is the first step in Fatou’s proof. The density of repelling cycles in the Julia set then follows from the finiteness of non-repelling cycles.

Proposition 5.11. *The closure of the set of periodic points contains the Julia set.*

Proof. Let $z_0 \in \mathcal{J}_f$ be arbitrary. Since \mathcal{J}_f is perfect, we may assume that z_0 is not a critical value of $f^{\circ 2}$, so there exist a neighborhood U of z_0 and three branches g_1, g_2, g_3 of f^{-2} with mutually disjoint images. We are considering the second iterate because if $d = 2$, there would only be two branches of f^{-1} .

Now consider the family of maps

$$h_m(z) := \frac{f^{\circ m}(z) - g_1(z)}{f^{\circ m}(z) - g_2(z)} \bigg/ \frac{g_3(z) - g_1(z)}{g_3(z) - g_2(z)}.$$

If $f^{\circ m}(z) \neq g_i(z)$, $i = 1, 2, 3$ for all z in some neighborhood U of z , then the maps $h_m(z)$ are never 0, 1 or ∞ in U , hence they form a normal family. But then the family $f^{\circ m}$ is also normal, since we can express $f^{\circ m}$ in terms of h_m :

$$f^{\circ m} = \frac{g_1(g_3 - g_2) - g_2(g_3 - g_1)h_m}{(g_3 - g_2) - (g_3 - g_1)h_m}.$$

This is a contradiction, so in every neighborhood of z there are roots of $f^{\circ m}(z) = g_i(z)$ for some i and some m , i.e., roots of $f^{\circ(m+2)}(z) = z$. \square

5.3. Connectivity of Julia sets of polynomials. Here is a first relation between the behaviour of critical orbits and the topology of the Julia set.

Theorem 5.12 (Fatou). *The filled-in Julia set \mathcal{K}_f of a polynomial f is connected if and only if all critical points of f belong to \mathcal{K}_f .*

Remark 5.13. Note that since the Julia set \mathcal{J}_f is the boundary of \mathcal{K}_f , \mathcal{J}_f is connected if and only if \mathcal{K}_f is connected.

Proof. Since f is a polynomial of degree $d \geq 2$, we have that $f(z)/z$ tends to infinity as z tends to infinity, hence there exists $R > 0$ so that $|f(z)| \geq 2|z|$ for $|z| > R$. Set $U_0 := \widehat{\mathbb{C}} \setminus \overline{D}_R$ and define inductively $U_{n+1} := f^{-1}(U_n)$. By the Riemann-Hurwitz formula, we have

$$\chi(U_{n+1}) = d\chi(U_n) - (d-1),$$

where the number $d-1$ is the multiplicity of the critical point at ∞ . Since U_0 is simply connected, we have $\chi(U_0) = 1$, and by induction, we see that when all the critical points of f belong to \mathcal{K}_f , then $\chi(U_n) = 1$ for all $n \geq 0$. It follows that

$$\Omega_\infty := \bigcup_{n \geq 0} U_n$$

is simply connected and its complement \mathcal{K}_f is connected.

Conversely, let $\Omega_\infty \subset \widehat{\mathbb{C}}$ be the connected component of the Fatou set \mathcal{F}_f which contains ∞ . Since the only inverse image of ∞ is ∞ itself, $f^{-1}(\Omega_\infty) = \Omega_\infty$. Applying the Riemann-Hurwitz formula to $f : \Omega_\infty \rightarrow \Omega_\infty$, we get

$$\chi(\Omega_\infty) = d\chi(\Omega_\infty) - (d-1) - k,$$

where the number $d-1$ is contributed by the critical point at ∞ , and the number k is the number of other critical points in Ω_∞ (i.e., not in \mathcal{K}_f) counted with multiplicity. If $\chi(\Omega_\infty)$ is finite, then $\chi(\Omega_\infty) = 1 + k/(d-1)$. But the Euler characteristic of a connected plane domain is at most 1. So $\chi(\Omega_\infty) = 1$ if no critical point is attracted to ∞ ; otherwise $\chi(\Omega_\infty)$ is infinite. \square

6. DESCRIPTION OF THE FATOU SET

We now return to the case where f is a rational map, not necessarily a polynomial. Our goal is to give a classification of periodic Fatou components.

Definition 6.1. *A map $f : U \rightarrow V$ is a proper map if for any compact set $K \subseteq V$, the preimage $f^{-1}(K)$ is compact.*

Proposition 6.2. *If U is a connected component of \mathcal{F}_f , then $f(U)$ is also a connected component of \mathcal{F}_f and $f : U \rightarrow f(U)$ is a proper map.*

Proof. The rational map $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ is a proper map. Hence, for any open set $V \subset \widehat{\mathbb{C}}$, $f : f^{-1}(V) \rightarrow V$ is a proper map. In particular, this holds for $U = \mathcal{F}_f$. But since $f^{-1}(\mathcal{F}_f) = \mathcal{F}_f$ by Proposition 2.9, we get that $f : \mathcal{F}_f \rightarrow \mathcal{F}_f$ is a proper map. Then the proposition follows, since the restriction of a proper map to a connected component of the domain is still proper. \square

Definition 6.3. *A connected component U of \mathcal{F}_f is called a Fatou component of f .*

Definition 6.4. *A Fatou component is periodic if there exists an integer $k \geq 1$ such that*

$$f^{\circ k}(U) = U.$$

Note that if U is periodic of period k for f , then it is invariant for $f^{\circ k}$.

Definition 6.5. *An invariant Fatou component U of f is called*

- a (super)attracting domain if there is a (super)attracting fixed point $\alpha \in U$ ($f(\alpha) = \alpha$ and $0 \leq |f'(\alpha)| < 1$) and the sequence $f^{\circ n}$ converges uniformly to α on every compact subset of U ;
- a parabolic domain if there is a fixed point $\alpha \in \partial U$ with $f'(\alpha) = 1$, and the sequence $f^{\circ n}$ converges uniformly to α on every compact subset of U ;
- a Siegel disk if it is simply connected, and if there exists an isomorphism $h : U \rightarrow \mathbb{D}$ such that

$$h \circ f \circ h^{-1}(z) = e^{2i\pi\theta} z$$

with $\theta \in \mathbb{R} \setminus \mathbb{Q}$;

- a Herman ring if it is doubly connected, and if there exists a radius r and an isomorphism

$$h : U \rightarrow \mathcal{A}_r := \{z \in \mathbb{C} ; r < |z| < 1\},$$

such that

$$h \circ f \circ h^{-1}(z) = e^{2i\pi\theta} z$$

with $\theta \in \mathbb{R} \setminus \mathbb{Q}$.

Remark 6.6. Clearly, the possibilities are mutually exclusive. Theorem 6.9 below asserts that these are the only possibilities and it is known that they all occur. The existence of Siegel disks was proved in 1942 by Siegel, and the existence of Herman rings was proved by Herman in 1981.

Figure 2 shows the Julia set of a polynomial having a Siegel disk and the Julia set of a rational map having a Herman ring.

Proposition 6.7. *If U is an invariant Fatou component of a rational map f of degree $d \geq 2$, then it is either simply connected, or doubly connected or its complement has infinitely many connected components. It is doubly connected if and only if it is a Herman ring.*

Proof. We can apply the Riemann-Hurwitz formula to $f : U \rightarrow U$. It gives

$$\chi(U) = d\chi(U) - n$$




FIGURE 2. Left: the polynomial $P : z \mapsto e^{i\pi(\sqrt{5}-1)z} + z^2$ has a Siegel disk (colored grey). We have drawn the orbits of some points in the Siegel disk. Each orbit accumulates on a \mathbb{R} -analytic circle. Right: if $t \in \mathbb{R}/\mathbb{Z}$ is chosen carefully enough, the rational map $f(z) = e^{2i\pi t} z^2(z-4)/(1-4z)$ has a Herman ring. It leaves the circle S^1 invariant and is conjugate to an irrational rotation on S^1 . The picture is drawn for $t = 0.61517321588\dots$

where d is the degree of $f : U \rightarrow U$ and n is the number of critical points of f in U , counting multiplicities. If $\chi(U)$ is finite, it follows that $\chi(U) = n/(d-1)$ is non negative, and since U is not the entire Riemann sphere, we have $\chi(U) = 0$ or 1 . Moreover, if $\chi(U) = 0$, i.e., if U is doubly connected, then $n = 0$ and $f : U \rightarrow U$ is a covering map. In addition, the degree of $f : U \rightarrow U$ is 1 since it preserves the modulus of U . Thus, $f : U \rightarrow U$ is an isomorphism. This isomorphism cannot be of finite order, since otherwise an iterate of f would have infinitely many fixed points. Therefore U is a Herman ring. \square

Remark 6.8. If f is a polynomial and U is a bounded Fatou component, then thanks to the maximum principle U is simply connected. In particular, there are no Herman rings.

The following result is due to Fatou. We will not give the detailed proof.

Theorem 6.9 (Classification of invariant Fatou components). *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a rational function of degree $d \geq 2$. An invariant Fatou component U of f is a (super)attracting domain, a parabolic domain, a Siegel disk or a Herman ring.*

Sketch of the proof. The classification starts by studying what are the possible limit values of the sequence of iterates $(f^{o_n} : U \rightarrow U)$.

Case 1. There is a non constant limit value.

Case 1.1. There is a sequence (n_k) such that $f^{o_{n_k}} \rightarrow \text{Id}$. This is a first place where we do not provide the details. One proves that

- either f has finite order (which is not the case since if $f^{\circ p} = \text{Id}$ on U for some $p \geq 1$, then $f^{\circ p} = \text{Id}$ on $\widehat{\mathbb{C}}$ by analytic continuation, which is not possible since $f^{\circ p}$ is a rational map of degree $d^p > 1$);
- or U is simply connected, in which case it is a Siegel disk;
- or U is doubly connected, in which case it is a Herman ring.

Case 1.2. There is a sequence (n_k) such that $f^{\circ n_k} \rightarrow \phi$ with ϕ non constant. By continuity, ϕ takes its values in \overline{U} . According to the Hurwitz Theorem, ϕ takes its values in U . Extracting a subsequence, we may assume that $n_{k+1} - n_k \rightarrow +\infty$ and that $f^{\circ(n_{k+1} - n_k)} \rightarrow \psi$. Then, $\psi \circ \phi = \phi$, so that ψ is the identity on the image of ϕ , thus on the whole component U by analytic continuation. In other words, we are in the previous situation : U is a Siegel disk or a Herman ring.

Case 2. Every limit value of the sequence $(f^{\circ n})$ is a constant. Note that if the sequence $(f^{\circ n_k})$ converges to α , then α is a fixed point of f . Indeed, since $f(z) \in U$ for all $z \in U$, we have that

$$\alpha \xleftarrow[k \rightarrow +\infty]{} f^{\circ n_k}(f(z)) = f(f^{\circ n_k}(z)) \xrightarrow[k \rightarrow +\infty]{} f(\alpha).$$

Note also that any point $z \in U$ can be joined to its image $f(z) \in U$ by a path compactly contained in U . It follows that the set of limit values of the sequence $(f^{\circ n}(z))$ is a continuum. Since f has finitely many fixed points, the whole sequence $(f^{\circ n})$ converges to a fixed point α of f .

Case 2.1. If $\alpha \in U$, the derivative of $f^{\circ n}$ at α tends to 0 as $n \rightarrow \infty$. As a consequence, α is a (super)attracting fixed point and U is a (super)attracting domain.

Case 2.2. If $\alpha \in \partial U$, then α cannot be (super)attracting since it belongs to the Julia set \mathcal{J}_f . It cannot be repelling since the sequence $(f^{\circ n})$ converges to α . So, it is indifferent. We claim that $D_\alpha f = 1$ and so, U is a parabolic domain.

Fix $z_0 \in U$ and set $z_n = f^{\circ n}(z_0)$. Let $\tilde{\rho} : [0, 1] \rightarrow U$ be a continuous path with $\tilde{\rho}(0) = z_0$ and $\tilde{\rho}(1) = z_1$. Then we can extend $\tilde{\rho}$ to a continuous path $\rho : [0, +\infty) \rightarrow U$ by setting $\rho(n+t) = f^{\circ n} \circ \tilde{\rho}(t)$ for all $n \in \mathbb{N}$ and all $t \in [0, 1)$. Then $\lim_{t \rightarrow +\infty} \rho(t) = \alpha$ and ρ satisfies $\rho(t+1) = f \circ \rho(t)$ for all $t \in [0, +\infty)$ and we conclude using the following lemma.

Lemma 6.10 (Snail lemma). *Let V be a neighborhood of the origin in \mathbb{C} and let $f : V \rightarrow f(V)$ be a holomorphic function such that $f(\alpha) = \alpha$ and $f'(\alpha) \neq 0$. If there is a continuous path $\rho : [0, +\infty) \rightarrow V \setminus \{\alpha\}$ such that $\lim_{t \rightarrow +\infty} \rho(t) = \alpha$ and $f \circ \rho(t) = \rho(t+1)$ for all $t \in [0, +\infty)$ then either $|f'(\alpha)| < 1$ or $f'(\alpha) = 1$.*

This is a second place where we do not provide the details. For a proof of this result see for example [50, Lemma 16.2]. \square

The following theorem, due to Fatou (1905), is probably the result which started the entire field of holomorphic dynamics.

Theorem 6.11. *A (super)attracting domain always contains at least one critical point.*

Proof. If U is a superattracting domain, then it contains a fixed critical point, and the result is obvious. If U is an attracting domain, it contains an attracting fixed point α with multiplier λ . We can show that there is a map $\phi : (\mathbb{C}, 0) \rightarrow (\widehat{\mathbb{C}}, \alpha)$ which conjugates the multiplication by λ to f . If U does not contain any critical

point, then ϕ extends to an entire map $\phi : \mathbb{C} \rightarrow U$ via the relation $\phi(\lambda z) = f \circ \phi(z)$. This contradicts the Liouville Theorem. \square

Remark 6.12. The same is true for parabolic domains, but the proof is more difficult and requires a detailed study of the local theory near a fixed point with multiplier 1. A proof can be found for example in [50, Theorem 10.15].

Fatou's classification of invariant Fatou components clearly provides also a classification of *periodic* Fatou components since they are invariant for an iterate of the rational map, and even *pre-periodic* Fatou components, that is components U so that there exist $m \geq 0$ and $p \geq 1$ such that $f^{o(m+p)}(U) = f^{o m}(U)$. The following fundamental result due to Sullivan completes the description for rational maps of degree $d \geq 2$.

Theorem 6.13 (Sullivan [60]). *Let $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ be a rational map of degree $d \geq 2$. Then every Fatou component of f is pre-periodic.*

The proof of Sullivan's non-wandering Theorem 6.13 strongly relies on the Ahlfors-Bers measurable mapping Theorem for quasi-conformal functions and we refer to the original paper of Sullivan [60] for it. The recent notes [19] present a proof due to Adam Epstein and based on a density result of Bers for quadratic differentials. Such results are strongly one-dimensional and do not have an analogue in higher dimension, making impossible to mimic Sullivan's proof there. Besides this observation, little was known about this problem until recently. In Section 8 we will present some recent results on Fatou components in dimension 2.

7. PARABOLIC IMPLOSION IN DIMENSION 1

In this section, we recall the main ingredients in the proof that the Julia set \mathcal{J}_f does not depend continuously on f for the Hausdorff topology on the space of compact subsets of \mathbb{C} . Lavaurs proved the following result (see also [27]).

Theorem 7.1 (Lavaurs [44]). *Assume $f : \mathbb{C} \rightarrow \mathbb{C}$ is a polynomial fixing 0 with $f(z) = z + z^2 + O(z^3)$. Then,*

$$\mathcal{J}_f \subsetneq \limsup_{\delta \rightarrow 0} \mathcal{J}_{f+\delta}.$$

The proof is based on the description of limits of iterates of $f + \delta$ in terms of maps called *Fatou coordinates* that will be introduced in subsection 7.2. Those limits are called *Lavaurs maps*.

7.1. Parabolic basin. In the rest of this section, we assume that $f : \mathbb{C} \rightarrow \mathbb{C}$ is a polynomial whose expansion near 0 is of the form

$$f(z) = z + z^2 + az^3 + O(z^4) \quad \text{with } a \in \mathbb{C}.$$

To understand the local dynamics of f near 0, it is convenient to consider the change of coordinates $Z := -1/z$. In the Z -coordinate, the expression of f becomes

$$F(Z) = Z + 1 + \frac{b}{Z} + O\left(\frac{1}{Z^2}\right) \quad \text{with } b := 1 - a.$$

In particular, if $R > 0$ is large enough, F maps the right half-plane $\{\Re(Z) > R\}$ into itself, so that if $r > 0$ is close enough to 0, f maps the disk $D(-r, r)$ into itself. In addition, the orbit under f of any point $z \in D(-r, r)$ converges to 0 tangentially to the real axis. Similarly, if $r > 0$ is sufficiently close to 0, there is a branch of f^{-1}

which maps the disk $D(r, r)$ into itself and the orbit under that branch of f^{-1} of any point $z \in D(r, r)$ converges to 0 tangentially to the real axis.

Definition 7.2. *The basin \mathcal{B}_f is the open set of points whose orbit under iteration of f intersects the disks $D(-r, r)$ for all $r > 0$.*

7.2. Fatou coordinates. In order to understand further the local dynamics of f near 0, it is customary to use local *attracting and repelling Fatou coordinates*. In the case of a polynomial, those Fatou coordinates have global properties.

Proposition 7.3. *There exists a (unique) attracting Fatou coordinate $\Phi_f : \mathcal{B}_f \rightarrow \mathbb{C}$ which semi-conjugates $f : \mathcal{B}_f \rightarrow \mathcal{B}_f$ to the translation $T_1 : \mathbb{C} \ni Z \mapsto Z + 1 \in \mathbb{C}$:*

$$\Phi_f \circ f = T_1 \circ \Phi_f.$$

and satisfies the normalization:

$$\Phi_f(z) = -\frac{1}{z} - b \log\left(-\frac{1}{z}\right) + o(1) \quad \text{as} \quad \Re\left(-\frac{1}{z}\right) \rightarrow +\infty.$$

Proof. For $z \in \mathcal{B}_f$, set

$$Z := -\frac{1}{z}, \quad Z_n := -\frac{1}{f^{\circ n}(z)} \quad \text{and} \quad \Phi_n(z) := Z_n - n - b \log Z_n.$$

We have that

$$\Re(Z_n) \rightarrow +\infty \quad \text{and} \quad Z_{n+1} = Z_n + 1 + \frac{b}{Z_n} + O\left(\frac{1}{Z_n^2}\right),$$

so that

$$\frac{1}{Z_n} = O\left(\frac{1}{Z+n}\right) \quad \text{as} \quad n \rightarrow +\infty.$$

As a consequence,

$$\Phi_{n+1}(z) - \Phi_n(z) = Z_{n+1} - Z_n - 1 - b \log \frac{Z_{n+1}}{Z_n} = O\left(\frac{1}{Z_n^2}\right) = O\left(\frac{1}{(Z+n)^2}\right).$$

So, the sequence (Φ_n) converges to a limit $\Phi_f : \mathcal{B}_f \rightarrow \mathbb{C}$ which satisfies

$$\Phi_f(z) = \Phi_0(z) + O\left(\frac{1}{Z}\right) = -\frac{1}{z} - b \log\left(-\frac{1}{z}\right) + o(1) \quad \text{as} \quad \Re\left(-\frac{1}{z}\right) \rightarrow +\infty.$$

Passing to the limit on the relation $\Phi_n \circ f = T_1 \circ \Phi_{n+1}$ yields the required result. \square

Figure ?? gives a rough idea of the behaviour of the Fatou coordinate Φ_f for the cubic polynomial $f(z) := z + z^2 + 0.95z^3$. The basin \mathcal{B}_f contains the two critical points of f . Those points and their iterated preimages form the critical points of Φ_f . Denote by c^+ the critical point with positive imaginary part and by c^- its complex conjugate. Set $v^\pm := \Phi_f(c^\pm)$. Points $z \in \mathcal{B}_f$ are colored according to the location $\Phi_f(z)$: dark grey when $\Im(\Phi_f(z)) < \Im(v^-)$, light grey when $\Im(v^-) < \Im(\Phi_f(z)) < \Im(v^+)$ and medium grey when $\Im(v^+) < \Im(\Phi_f(z))$.

Proposition 7.4. *There exists a (unique) repelling Fatou parametrization $\Psi_f : \mathbb{C} \rightarrow \mathbb{C}$ which semi-conjugates $T_1 : \mathbb{C} \rightarrow \mathbb{C}$ to $f : \mathbb{C} \rightarrow \mathbb{C}$:*

$$\Psi_f \circ T_1 = f \circ \Psi_f.$$

and satisfies the normalization:

$$\Psi_f(Z) = -\frac{1}{Z + b \log(-Z) + o(1)} \quad \text{as} \quad \Re(Z) \rightarrow -\infty.$$

Proof. Choose $r > 0$ sufficiently close to 0 so that there is a branch g of f^{-1} which maps the disk $D(r, r)$ into itself. For $z \in D(r, r)$, set

$$Z := -\frac{1}{z}, \quad Z_n := -\frac{1}{g^{\circ n}(z)} \quad \text{and} \quad \Phi_n(z) := Z_n + n - b \log(-Z_n).$$

Note that

$$Z_n = Z_{n+1} + 1 + \frac{b}{Z_{n+1}} + O\left(\frac{1}{Z_{n+1}^2}\right).$$

So, as in the previous proof

$$\Phi_{n+1}(z) - \Phi_n(z) = Z_{n+1} - Z_n + 1 + b \log \frac{Z_{n+1}}{Z_n} = O\left(\frac{1}{Z_{n+1}^2}\right) = O\left(\frac{1}{(Z-n)^2}\right).$$

The sequence Φ_n converges in the left half-plane $\{\Re(Z) < -1/(2r)\}$ to a limit Φ_g which satisfies

$$\Phi_g(z) = \Phi_0(z) + O\left(\frac{1}{Z}\right) = Z - b \log(-Z) + o(1) \quad \text{as} \quad \Re\left(-\frac{1}{z}\right) \rightarrow -\infty.$$

Passing to the limit on the equation $\Phi_{n+1} \circ f = T_1 \circ \Phi_n$ shows that Φ_g conjugates f to T_1 . The inverse Ψ_f of Φ_g conjugates T_1 to f and

$$Z = \Phi_g \circ \Psi_f(Z) = -\frac{1}{\Psi_f(Z)} - b \log\left(-\frac{1}{\Psi_f(Z)}\right) + o(1) = -\frac{1}{\Psi_f(Z)} - b \log(-Z) + o(1)$$

as $\Re(Z) \rightarrow -\infty$. □

7.3. Lavaurs maps.

Definition 7.5. *The Lavaurs map $\mathcal{L}_f : \mathcal{B}_f \rightarrow \mathbb{C}$ is the map*

$$\mathcal{L}_f := \Psi_f \circ \Phi_f : \mathcal{B}_f \rightarrow \mathbb{C}.$$

Note that the Lavaurs map \mathcal{L}_f commutes with f . Indeed, $\Phi_f \circ f = T_1 \circ \Phi_f$ and $\Psi_f \circ T_1 = f \circ \Psi_f$, so that

$$\mathcal{L}_f \circ f = \Psi_f \circ \Phi_f \circ f = \Psi_f \circ T_1 \circ \Phi_f = f \circ \Psi_f \circ \Phi_f = f \circ \mathcal{L}_f.$$

Figure 3 gives a rough idea of the behaviour of the Lavaurs map \mathcal{L}_f for the cubic polynomial $f(z) := z + z^2 + 0.95z^3$. Points in the basin \mathcal{B}_f are colored according to the location of their image by \mathcal{L}_f : grey when $\mathcal{L}_f(z) \in \mathcal{B}_f$, white otherwise. The restriction of \mathcal{L}_f to each bounded white domain is a covering of $\mathbb{C} \setminus \overline{\mathcal{B}_f}$.

7.4. Discontinuity of the Julia set. Set

$$\mathcal{K}(\mathcal{L}_f) := \bigcap_{n \geq 0} \mathcal{L}_f^{-n}(\mathcal{K}_f) \quad \text{and} \quad \mathcal{J}(\mathcal{L}_f) := \partial \mathcal{K}(\mathcal{L}_f).$$

Figure 4 shows $\mathcal{K}(\mathcal{L}_f)$ for the cubic polynomial $f(z) = z + z^2 + 0.95z^3$. The Lavaurs map \mathcal{L}_f has two complex conjugate sets of attracting fixed points. The fixed points of \mathcal{L}_f are indicated and their basins of attraction are colored (dark grey for one of the fixed points, and light grey for the others). Those basins form the interior of $\mathcal{K}(\mathcal{L}_f)$. The black set $\mathcal{J}(\mathcal{L}_f)$ is the topological boundary of $\mathcal{K}(\mathcal{L}_f)$.

The following result may be considered as the main reason why Lavaurs maps are studied in holomorphic dynamics.




FIGURE 3. Behavior of \mathcal{L}_f for $f(z) := z + z^2 + 0.95z^3$.




FIGURE 4. The set $\mathcal{K}(\mathcal{L}_f)$ for $f(z) = z + z^2 + 0.95z^3$.

Proposition 7.6 (Lavaurs [44]). *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a polynomial whose expansion at 0 is $f(z) = z + z^2 + O(z^3)$. Let (N_n) be a sequence of integers tending to $+\infty$ and (ε_n) be a sequence of complex numbers tending to 0, such that*

$$N_n - \frac{\pi}{\varepsilon_n} \rightarrow 0.$$

Then,

$$(f + \varepsilon_n^2)^{\circ N_n} \rightarrow \mathcal{L}_f \quad \text{locally uniformly on } \mathcal{B}_f.$$

In addition,

$$\mathcal{J}_f \subsetneq \mathcal{J}(\mathcal{L}_f) \subseteq \liminf \mathcal{J}_{f+\varepsilon_n^2} \quad \text{and} \quad \limsup \mathcal{K}_{f+\varepsilon_n^2} \subseteq \mathcal{K}(\mathcal{L}_f) \subsetneq \mathcal{K}_f.$$

We do not present the proof of this result which is rather technical, and can be found in [27] or [56] for example. Instead, we will show that the result holds for the Möbius transformation

$$g(z) = \frac{z}{1-z} = z + z^2 + O(z^3).$$

In that case, all maps involved are Möbius transformations and the computations are explicit. If we perform the change of coordinates $Z = -1/z$, the Möbius transformation g gets conjugated to T_1 . Thus,

$$\Phi_g(z) = -\frac{1}{z} \quad \text{and} \quad \Psi_g(Z) = -\frac{1}{Z}, \quad \text{so that} \quad \mathcal{L}_g = \text{Id}.$$

Note that $g + \varepsilon^2$ is also a Möbius transformation. It has two fixed points

$$\alpha^\pm = \pm i\varepsilon + O(\varepsilon^2) \quad \text{with multipliers} \quad \lambda^\pm = \exp(\pm 2i\varepsilon + O(\varepsilon^3)).$$

So, if $N \rightarrow +\infty$ and

$$N - \frac{\pi}{\varepsilon} \rightarrow 0, \quad \text{so that} \quad \varepsilon = \frac{\pi}{N + o(1)} = \frac{\pi}{N} + o\left(\frac{1}{N^2}\right),$$

then $(g + \varepsilon^2)^{\circ N}$ is a Möbius transformation fixing α^\pm with multipliers

$$\mu^\pm := (\lambda^\pm)^N = \exp(\pm 2\pi i + o(1/N)) = 1 + o\left(\frac{1}{N}\right).$$

This Möbius transformation is

$$z \mapsto \alpha^+ + \frac{\mu^+ \cdot (z - \alpha^+)}{1 - \frac{\mu^+ - 1}{\alpha^+ - \alpha^-} (z - \alpha^+)}.$$

Since

$$\mu^+ - 1 = o\left(\frac{1}{N}\right) = o(\alpha^+ - \alpha^-),$$

we see that indeed,

$$(g + \varepsilon^2)^{\circ N} \xrightarrow{N \rightarrow +\infty} \mathcal{L}_g.$$

Recently, in [63], Vivas studied a non-autonomous parabolic implosion in dimension 1 for the map $f(z) = \frac{z}{1-z}$.

In higher dimension, (semi-)parabolic implosion was recently studied for dissipative polynomial automorphisms of \mathbb{C}^2 by Bedford, Smillie and Ueda in [9] (see also [29]) and their strategy was recently adapted by Bianchi in [11] for a perturbation of a class of holomorphic endomorphisms tangent to identity, establishing a two-dimensional Lavaurs theorem for such a class.

8. FATOU COMPONENTS FOR POLYNOMIAL MAPS IN DIMENSION 2

We end these notes by giving an updated account of the recent results on Fatou components for polynomial skew-products in complex dimension two in a neighbourhood of a periodic fiber. We divide our discussion according to the different possible kinds of periodic fibers.

8.1. Preliminaries. Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a holomorphic endomorphism of \mathbb{C}^2 , and consider the discrete holomorphic dynamical system given by the iteration of F . In the investigation of the global behaviour of such a system it is natural to give the generalization of the definition of the *Fatou set of F* as the largest open set $\mathcal{F}(F)$ where the family of iterates $(F^{\circ n})_{n \in \mathbb{N}}$ of F is normal. A connected component of the Fatou set is called a *Fatou component*.

We have seen in the previous sections that in complex dimension one, Fatou components of rational maps of degree at least 2 on the Riemann sphere are well understood, thanks to Theorem 6.9 and Theorem 6.13.

In complex dimension two, the understanding of Fatou components is far less complete. A considerable progress in the classification of periodic Fatou components has been achieved thanks to Bedford and Smillie [6] [7] [8], Fornæss and Sibony [33], Lyubich and Peters [47] and Ueda [61].

The question of the existence of wandering (i.e., not pre-periodic) Fatou components in higher dimension was put forward by several authors since the 1990's (see e.g., [35]). Higher dimensional *transcendental* (i.e., non polynomial) maps with wandering domains can be constructed from one-dimensional examples by taking direct products. An example of a transcendental *biholomorphism* of \mathbb{C}^2 with a wandering Fatou component oscillating to infinity was constructed by Fornæss and Sibony in [34]. Nonetheless, until recently very little was known about the existence of wandering Fatou components for holomorphic endomorphisms of $\mathbb{P}^2(\mathbb{C})$ or for polynomial endomorphisms of \mathbb{C}^2 .

A first natural class of polynomial endomorphisms of \mathbb{C}^2 to consider are *direct product*, that is maps $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ of the form

$$F(z, w) = (f(z), g(w)),$$

where f and g are complex polynomials in one variable. This allows us to recover the generalizations of one-dimensional dynamical behaviours in dimension two. However, direct products are a very particular class and they do not give us a complete understanding of all possible behaviours of polynomial endomorphisms in \mathbb{C}^2 .

A more interesting class to consider is given by polynomial *skew-products* in \mathbb{C}^2 , namely polynomial maps $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ of the form

$$(1) \quad F(z, w) = (f(z, w), g(w)),$$

where g is a complex polynomial in one variable and f is a complex polynomial in two variables. Since they leave invariant the fibration $\{w = \text{const.}\}$, skew-products allow us to *build on one-dimensional dynamics* and to get a first flavour of the richness of the higher dimension setting we are working in. This idea has been used by several authors to construct maps with particular dynamical properties. Dujardin, for example, used in [28] specific skew-products to construct a non-laminar Green current. Boc-Thaler, Fornæss and Peters constructed in [17] a map having a Fatou component with a punctured limit set. Last but not least, as we will explain in subsection 8.2, skew-products are one of the key ingredients in the construction we recently obtained in [4] with Astorg, Dujardin, and Peters of holomorphic endomorphisms of $\mathbb{P}^2(\mathbb{C})$ having a wandering Fatou component.

The investigation of the holomorphic dynamics of polynomial skew-products was started by Heinemann [37] and then continued by Jonsson [40]. The topology of Fatou components of skew-products has been studied by Roeder in [54].

Given a Fatou component Ω of a polynomial skew-product F in \mathbb{C}^2 , its projection on the second coordinate $\Omega_2 = \pi_2(\Omega)$ is a Fatou components for g and hence thanks to Sullivan's non-wandering Theorem 6.13, up to considering an iterate of F , it has to fall into one of the three cases given by Theorem 6.9, and moreover, since we are considering polynomials, Herman rings cannot occur. Therefore, since (pre-)periodic points for g correspond to (pre-)periodic fibers for F , up to considering an iterate of F , we can restrict ourselves to study what happens in neighbourhoods of invariant fibers of the form $\{w = w_0\}$. One-dimensional theory also describes the dynamics on the invariant fiber, which is given by the action of the one-dimensional polynomial $f(z, w_0) := f_{w_0}(z)$, and hence the Fatou components of f_c will be again all pre-periodic and, up to consider an iterate, we can assume that they are either attracting basins, or parabolic basins or Siegel disks. This structure leads us to two immediate questions.

- (1) Do all Fatou components of f_c bulge to two-dimensional Fatou components of F ?
- (2) Is it possible to have wandering Fatou components for F in a neighbourhood of an invariant fiber?

In the following we shall call an invariant fiber $\{w = w_0\}$ *attracting*, *parabolic* or *elliptic* according to whether w_0 is an attracting, parabolic or elliptic fixed point for g . A *bulging* Fatou component will be a Fatou component Ω of F such that $\Omega \cap \{w = w_0\}$ is a one-dimensional Fatou component of f_{w_0} on the invariant fiber $\{w = w_0\}$. We shall say that a Fatou component Ω_{w_0} of f_{w_0} on the invariant fiber $\{w = w_0\}$ is *bulging* if there exists a bulging Fatou component Ω of F so that $\Omega_{w_0} = \Omega \cap \{w = w_0\}$.

8.2. Attracting invariant fiber. Let us consider a polynomial skew-product $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ of degree $d \geq 2$

$$F(z, w) = (f(z, w), g(w)),$$

with an *attracting invariant fiber*. We can assume without loss of generality that the invariant fiber is $\{w = 0\}$. Therefore we have $g(0) = 0$ and $|g'(0)| < 1$. In this case we know that there exists an *attracting basin*, containing the origin, of points whose iterates converge to the origin. The rate of convergence to the fixed point depends on whether the fixed point is *superattracting*, i.e., $g'(0) = 0$, or *attracting* or *geometrically attracting*, i.e., $g'(0) \neq 0$.

8.2.1. Superattracting case. This setting was studied by Lilov in [45] who was able to answer both questions stated in the introduction. He first proved the following result giving a positive answer to Question 1.

Theorem 8.1 (Lilov [45]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1) of degree $d \geq 2$. Let $\{w = w_0\}$ be a superattracting invariant fiber for F . Then all one-dimensional Fatou components of f_{w_0} bulge to Fatou components of F .*

Idea of the proof. We can assume $w_0 = 0$ without loss of generality. Thanks to Theorem 6.9 and Theorem 6.13, all Fatou components of the restriction $f_0(z) = f(z, 0)$ of $f(z, w)$ to the invariant fiber are (pre-)periodic and are either attracting basins, or parabolic basins or Siegel domains. The strategy of the proof is to prove separately for each of these cases that the corresponding component is contained in

a two-dimensional Fatou component of F . The bulging of one-dimensional Fatou components of attracting periodic points of $f_0(z)$ is well-known and follows for instance from the results of Rosay and Rudin [55]. For the remaining cases, by [45, Theorem 3.17] there exists a strong stable manifold through all point in the one-dimensional Fatou components of parabolic or elliptic periodic points of $f_0(z)$, and so the corresponding bulging Fatou components simply consist of the union of such manifolds. \square

Then Lilov proved the following result implying the non-existence of wandering Fatou components in a neighbourhood of a superattracting invariant fiber.

Theorem 8.2 (Lilov [45]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1) of degree $d \geq 2$. Let $\{w = w_0\}$ be a superattracting invariant fiber for F and let \mathcal{B} be the immediate basin of the superattracting fixed point w_0 . Take $w \in \mathcal{B}$ and let D_w be a one-dimensional open disk lying in the fiber over w . Then the forward orbit of D_w must intersect one of the bulging Fatou components of f_{w_0} .*

The proof relies on the repeated use of [45, Lemma 3.2.4] applied to the orbit of a disk lying in a fiber over a point in the attracting basin, in order to obtain estimates from below for the radii of the images. Thanks to [45, Proposition 3.2.8], by studying the geometry of the bulging Fatou components, it is also possible to obtain an upper bound on the largest possible disk lying in a fiber over a point in the attracting basin that can lie in the complement of a bulging Fatou component, depending on the distance to the invariant fiber. The conclusion then follows combining these two estimates.

All bulging Fatou components are (pre-)periodic, therefore all Fatou components for F in a neighbourhood of a superattracting invariant fiber are (pre-)periodic, and then the non-existence of wandering Fatou components in a neighbourhood of a superattracting invariant fiber follows immediately.

Corollary 8.3 (Lilov [45]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1) of degree $d \geq 2$. Let $\{w = w_0\}$ be a superattracting invariant fiber for F and let \mathcal{B} be the immediate basin of the superattracting fixed point w_0 . Then there are no wandering Fatou components in $\mathcal{B} \times \mathbb{C}$.*

8.2.2. Geometrically attracting case. The geometrically attracting case was first partially addressed by Lilov in [45] even if not stated explicitly. In fact, the proof of Theorem 8.1 can be readily adjusted to this case obtaining the following statement answering Question 1.

Theorem 8.4 (Lilov [45]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1) of degree $d \geq 2$. Let $\{w = w_0\}$ be an attracting invariant fiber for F . Then all one-dimensional Fatou components of f_{w_0} bulge to Fatou components of F .*

On the other hand, the proof of Theorem 8.2 cannot be generalized to this setting, which is indeed more complicated than the superattracting case. In fact, Theorem 8.2 does not hold in general, as showed by Peters and Vivas with the following result.

Theorem 8.5 (Peters-Vivas [53]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form*

$$(2) \quad F(z, w) = (p(z) + q(w), \lambda w),$$

with $0 < |\lambda| < 1$ and p and q complex polynomials. Then there exists a triple (λ, p, q) and a holomorphic disk $D \subset \{w = w_0\}$ whose forward orbit accumulates at a point $(z_0, 0)$, where z_0 is a repelling fixed point in the Julia set of f_0 .

As a consequence, the forward orbits of D cannot intersect the bulging Fatou components of f_0 .

The family $(F|_D^{\circ n})_{n \in \mathbb{N}}$ is normal on the disks D , and so these are Fatou disks. However such disks are completely contained in the Julia set of F , which is the complement in \mathbb{C}^2 of the Fatou set (see [53, Theorem 6.1]).

The geometrically attracting case has been further investigated by Peters and Smit in [52]. They focused their investigation on polynomial skew-products such that the action on the invariant attracting fiber is *subhyperbolic*, that is the polynomial does not have parabolic periodic points and all critical points lying on the Julia set are pre-periodic. They proved the following result.

Proposition 8.6 (Peters-Smit [52]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1). Assume that the origin is an attracting, not superattracting, fixed point for g with corresponding basin B_g , and the polynomial $f_0(z) := f(z, 0)$ is subhyperbolic. Then there exists a set $E \subset \mathbb{C}$ of full measure, such that for every $\tilde{w}_0 \in E$ the forward orbit of every disk in the fiber $\{w = \tilde{w}_0\}$ must intersect a bulging Fatou component of f_0 .*

Idea of the proof. Notice that it suffices to prove the proposition in a neighbourhood of the attracting fiber $\{w = 0\}$. Therefore, up to considering a smaller neighbourhood, we can assume without loss of generality that $g(w) = \lambda w$, and

$$f(z, w) = a_0(w) + a_1(w)z + \dots + a_d(w)z^d$$

where $a_0(w), \dots, a_d(w)$ are holomorphic functions in w . The subhyperbolicity of the polynomial f_0 implies that its Fatou set is the union of finitely many attracting basins, and the orbits of the critical points contained in the Fatou set converge to one of these attracting cycles. The proof can be divided into 5 main steps.

Step 1. Fix $R > 0$ large enough so that for all z such that $|z| > R$ we have $|f_0(z)| > 2|z|$ and set

$$W_0 = \{|z| > R\} \cup \bigcup_{y \in \text{Att}(f_0)} W_y$$

where $\text{Att}(f_0)$ is the set of all attracting periodic points of f_0 , and for each $y \in \text{Att}(f_0)$ the set W_y is an open neighbourhood of the orbit of y such that $f_0(\overline{W_y}) \subset W_y$. Fix a neighbourhood U of the post-critical set of f_0 . Then by [52, Proposition 15], there exists a set $E \subset \mathbb{C}$ of full measure in a neighbourhood of the origin such that for all $\tilde{w}_0 \in E$ there exists a constant $C = C(\tilde{w}_0, U)$ such that for all $n \in \mathbb{N}$ we have

$$(3) \quad \text{Card} \left\{ z : \frac{\partial F_1^{\circ n}}{\partial z}(z, \tilde{w}_0) = 0 \text{ and } F_1^{\circ n}(z, \tilde{w}_0) \notin W_0 \times U \right\} \leq C\sqrt{n},$$

where $F_1^{\circ n}$ is the first component of the n -th iterate of F .

Step 2. Assume by contradiction that a fiber $\{w = \tilde{w}_0\}$, with $\tilde{w}_0 \in E$, contains a disk D whose forward orbit avoids the bulging Fatou components of f_0 . Then the restriction of $F^{\circ n}$ to D is bounded and hence a normal family. Therefore, up to shrinking D there exists a subsequence $F^{\circ n_j}$ such that $F^{\circ n_j}|_D$ converges, uniformly

on compact subsets of D , to a point ζ in the Julia set of f_0 . Moreover, there exists $\varepsilon > 0$ so that $F^{on}(D) \cap (W_0 \times D(0, \varepsilon))$ is empty for all $n \in \mathbb{N}$.

Step 3. Each critical point x contained in the Julia set is eventually mapped into a repelling periodic point, and up to considering an iterate of F we may assume that it is eventually mapped into a repelling fixed point with multiplier μ , with $|\mu| > 1$. The main tool to control the orbits of the critical points of F is obtained using a linearization map of the unstable manifold of the repelling fixed point, given by a map $\Phi : \mathbb{C} \rightarrow \mathbb{C}$ satisfying $\Phi(\mu t) = f_0^k \circ \Phi(t)$ for some $k \in \mathbb{N}$. Thanks to [52, Proposition 10], there exist $\tilde{C} > 1$ and $0 < \gamma < 1$ so that

$$(4) \quad \text{Area}(F^{on}(D)) \leq \tilde{C}\gamma^n.$$

Step 4. We may assume that ζ does not lie in the post-critical set, and we may choose U and $r > 0$ such that $D(\zeta, r) \cap (U \cup W_0) = \emptyset$. Let $j_1 \in \mathbb{N}$ be such that $F_1^{onj}(D) \subseteq D(\zeta, \frac{r}{2})$ for all $j \geq j_1$, and consider O_j the connected component of $(F^{onj})^{-1}(D(\zeta, r) \times \{\lambda^{nj}\tilde{w}_0\})$ containing D . Then $D \subseteq O_j \subseteq D(0, R) \times \{\tilde{w}_0\}$, and we can study the proper holomorphic function $F_1^{onj} : O_j \rightarrow D(\zeta, r)$. Thanks to (3), such a map has at most $d_j = C\sqrt{n_j}$ critical points.

Step 5. It is possible (see [52, Proposition 28]) to find a uniform constant $C_1 > 0$ so that if $f : \mathbb{D} \rightarrow \mathbb{D}$ is a proper holomorphic function of degree d , the set $R \subset \mathbb{D}$ has Poincaré area equal to A , and $d \cdot A^{1/2d} < 8$, then the Poincaré area of $f^{-1}(R)$ is at most $C_1 d^3 A^{1/d}$. Then, setting $R_j = F_1^{onj}(D)$ and denoting by A_j its Poincaré area $\text{Area}_{D(\zeta, r)}(R_j)$ with respect to $D(\zeta, r)$, for $j \geq j_1$, we have $R_j \subseteq D(\zeta, r)$, and we can estimate A_j applying (4). Therefore there exists $j_2 \geq j_1$ such that $d_j A_j^{1/2d_j} < 1/8$ for all $j \geq j_2$. This implies

$$\text{Area}_{D(0, R)}(D) \leq \text{Area}_{O_j}(D) \leq C_2 d_j^3 A_j^{1/d_j} \leq M n_j^{3/2} \gamma^{n_j^{3/2}}$$

where $M > 0$. The contradiction follows from the fact that the last expression will converge to zero as j increases towards infinity. \square

Thanks to the fact that in particular E is dense, Peters and Smit are able to give a negative answer to Question 2 when the action on the invariant fiber is subhyperbolic. They also obtain as a corollary that the only Fatou components of F are the bulging ones, since the topological degree of F equals the one of f_0 , implying that the only Fatou components that can be mapped onto the bulging Fatou components of f_0 are exactly those bulging Fatou components.

Theorem 8.7 (Peters-Smit [52]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1). Assume that the origin is an attracting fixed point for g with corresponding basin \mathcal{B}_g , and the polynomial $f_0(z) := f(z, 0)$ is subhyperbolic. Then F has no wandering Fatou component over \mathcal{B}_g .*

Very recently, Ji was able in [38] to generalize Lilov's Theorem 8.2 to polynomial skew-products with an invariant geometrically attracting fiber under the hypothesis that the multiplier of the invariant fiber is sufficiently small. More precisely he proved the following result.

Theorem 8.8 (Ji [38]). *Let $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be a polynomial skew-product of the form (1) of degree $d \geq 2$. Let $\{w = w_0\}$ be an attracting invariant fiber for F and let \mathcal{B}_{w_0} be the basin of the attracting fixed point c . Then there exists $\lambda_0(w_0, f) > 0$*

depending only on f and c such that if $|g'(w_0)| < \lambda_0$, then there are no wandering Fatou components in $\mathcal{B}_{w_0} \times \mathbb{C}$.

The proof of this result follows Lilov's strategy. The main difficulty is due to the breaking down of Lilov's argument in the geometrically attracting case as we pointed out at the beginning of this section. Ji is able to overcome such difficulty by adapting a one-dimensional result due to Denker, Przytycki and Urbanski in [24] in this case. Such result is used to obtain estimates of the size of bulging Fatou components and of the size of forward images of wandering Fatou disks.

Ji also proved in [39] that if $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is a polynomial skew-product with an attracting invariant fiber L such that the restriction of F to L is non-uniformly hyperbolic, then the Fatou set in the basin of L coincides with the union of the basins of attracting cycles, and the Julia set in the basin of L has Lebesgue measure zero. Therefore there are no wandering Fatou components in the basin of the invariant attracting fiber.

8.3. Parabolic invariant fiber and wandering domains. A first contribution to the investigation of this case is due to Vivas, who proved a parametrization result [62, Theorem 3.1] for the unstable manifolds for *special* parabolic skew-product of \mathbb{C}^2 . Vivas used this parametrization as the main tool to prove the analogue of Theorem 8.5 for special parabolic skew-product. However, this construction does not allow to construct a wandering Fatou component in a neighbourhood of the parabolic invariant fiber.

In [4], together with Astorg, Dujardin and Peters, we proved the existence of polynomial skew-products of \mathbb{C}^2 , extending to holomorphic endomorphisms of $\mathbb{P}^2(\mathbb{C})$, having a wandering Fatou component. The key tool consists in using parabolic implosion techniques on polynomial skew-products, and this idea was initially suggested by Lyubich. The main strategy is to combine slow convergence to an invariant parabolic fiber and parabolic transition in the fiber direction, to produce orbits shadowing those of the so-called Lavaurs map, that we defined in the previous section.

Theorem 8.9 ([4]). *There exists a holomorphic endomorphism $F : \mathbb{P}^2(\mathbb{C}) \rightarrow \mathbb{P}^2(\mathbb{C})$, induced by a polynomial skew-product mapping $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$, having a wandering Fatou component. More precisely, let $f : \mathbb{C} \rightarrow \mathbb{C}$ and $g : \mathbb{C} \rightarrow \mathbb{C}$ be polynomials of the form*

$$(5) \quad f(z) = z + z^2 + O(z^3) \text{ and } g(w) = w - w^2 + O(w^3).$$

If the Lavaurs map $\mathcal{L}_f : \mathcal{B}_f \rightarrow \mathbb{C}$ has an attracting fixed point, then the skew-product $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ defined by

$$(6) \quad F(z, w) := \left(f(z) + \frac{\pi^2}{4}w, g(w) \right)$$

has a wandering Fatou component.

The orbits in these wandering Fatou components are bounded and the approach used in the proof is essentially local. Notice that if f and g have the same degree, F extends to a holomorphic endomorphism of $\mathbb{P}^2(\mathbb{C})$. Moreover we can obtain examples in arbitrary dimension $k \geq 2$ by simply considering products mappings of the form (F, Q) , where Q has a fixed Fatou component.

A first step in the proof of Theorem 8.9 is to find a parabolic polynomial f whose Lavaurs map \mathcal{L}_f admits an attracting fixed point.

Proposition 8.10 ([4, Proposition B]). *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be the cubic polynomial defined by*

$$f(z) = z + z^2 + az^3 \quad \text{with } a \in \mathbb{C}.$$

If $r > 0$ is sufficiently close to 0 and a belongs to the disk $D(1 - r, r)$, then the Lavaurs map $\mathcal{L}_f : \mathcal{B}_f \rightarrow \mathbb{C}$ admits an attracting fixed point.

Idea of the proof. We consider

$$\mathcal{U}_f := \psi_f^{-1}(\mathcal{B}_f) \quad \text{and} \quad \mathcal{E}_f := \varphi_f \circ \psi_f : \mathcal{U}_f \rightarrow \mathbb{C}.$$

The open set \mathcal{U}_f contains an upper half-plane and a lower half-plane, and it is invariant under T_1 . Like the Lavaurs map, the map \mathcal{E}_f commutes with T_1 , therefore $\mathcal{E}_f - \text{Id}$ is periodic of period 1 and admits a Fourier expansion in a upper half-plane:

$$\mathcal{E}_f(Z) = Z + \sum_{k \geq 0} c_k e^{2\pi i k Z}.$$

Using the expansion of φ_f and ψ_f near infinity, we obtain with an elementary computation:

$$\begin{aligned} \mathcal{E}_f(Z) &= \varphi_f \circ \psi_f(Z) = Z + (1 - a) \log(-Z) + o(1) \\ &\quad - (1 - a) \log(Z + (1 - a) \log(-Z) + o(1)) + o(1) \\ &= Z + (1 - a) \log(Z) - \pi i(1 - a) - (1 - a) \log(Z) + o(1) \\ &= Z - \pi i(1 - a) + o(1), \end{aligned}$$

and so $c_0 = -\pi i(1 - a)$.

Thanks to a more elaborate argument, based on the notion of finite type analytic map introduced by Adam Epstein, it is possible to prove that:

$$\mathcal{E}_f(Z) = Z - \pi i(1 - a) + c_1 e^{2\pi i Z} + o(e^{2\pi i Z}) \quad \text{with } c_1 \neq 0.$$

It then follows that for $a \neq 1$ close to 1, \mathcal{E}_f has a fixed point Z_f with multiplier ρ_f satisfying

$$c_1 e^{2\pi i Z_f} \sim \pi i(1 - a) \quad \text{and} \quad \rho_f - 1 \sim 2\pi i c_1 e^{2\pi i Z_f} \sim -2\pi^2(1 - a) \quad \text{as } a \rightarrow 1.$$

Therefore, for $r > 0$ sufficiently close to 0 and $a \in D(1 - r, 1)$, the multiplier ρ_f belongs to the unit disk and Z_f is an attracting fixed point of \mathcal{E}_f . This concludes the proof since $\psi_f : \mathcal{U}_f \rightarrow \mathcal{B}_f$ semi-conjugates \mathcal{E}_f to \mathcal{L}_f , and so, the fact that Z_f is an attracting fixed point of \mathcal{E}_f implies that the point $\psi_f(Z_f)$ is an attracting fixed point of \mathcal{L}_f . \square

The key result in the proof of Theorem 8.9 relies on a non-autonomous analogue of Lavaurs estimates in the setting of skew-products.

Let \mathcal{B}_f and \mathcal{B}_g be the parabolic basins of 0 under iteration of respectively f and g . One of the key points is to choose $(\tilde{z}_0, \tilde{w}_0) \in \mathcal{B}_f \times \mathcal{B}_g$ so that the first coordinate of $F^{\circ m}(\tilde{z}_0, \tilde{w}_0)$ returns infinitely many times close to the attracting fixed point of \mathcal{L}_f . The proof is designed so that the return times are the integers n^2 for $n \geq n_0$. Therefore, we need to analyze the orbit segment between n^2 and $(n + 1)^2$, which is of length $2n + 1$.

Proposition 8.11 ([4]). *As $n \rightarrow +\infty$, the sequence of maps*

$$\mathbb{C}^2 \ni (z, w) \mapsto F^{2n+1}(z, g^{n^2}(w)) \in \mathbb{C}^2$$

converges locally uniformly in $\mathcal{B}_f \times \mathcal{B}_g$ to the map

$$\mathcal{B}_f \times \mathcal{B}_g \ni (z, w) \mapsto (\mathcal{L}_f(z), 0) \in \mathbb{C} \times \{0\}.$$

Idea of the proof. Let \mathcal{B}_g the parabolic basin of 0 under iteration of g . For all $w \in \mathcal{B}_g$, the orbit $g^{om}(w)$ converges to 0 like $1/m$. We want to analyze the behaviour of F starting at $(z, g^{n^2}(w))$ during $2n+1$ iterates. For large n , the first coordinate of F along this orbit segment is approximately

$$f(z) + \varepsilon^2 \text{ with } \frac{\pi}{\varepsilon} \simeq 2n.$$

As we saw in the previous section, a rough statement of Lavaurs Theorem from parabolic implosion gives us that if $\frac{\pi}{\varepsilon} = 2n$, then for large n , the $(2n)^{\text{th}}$ iterate of $f(z) + \varepsilon^2$ is approximately equal to $\mathcal{L}_f(z)$ on \mathcal{B}_f .

Our setting is different since in our case ε keeps decreasing along the orbit. Indeed on the first coordinate we are taking the composition of $2n+1$ transformations of the form

$$f(z) + \varepsilon_k^2 \text{ with } \frac{\pi}{\varepsilon_k} \simeq 2n + \frac{k}{n} \text{ and } 1 \leq k \leq 2n+1.$$

The key step in the proof of the statement consists in a detailed analysis of this non-autonomous situation, proving that the decay of ε_k is counterbalanced by taking *exactly* one additional iterate of F . \square

With this proposition in hand, the proof of Theorem 8.9 is easily completed.

Proof of Theorem 8.9. Let ξ be an attracting fixed point of \mathcal{L}_f and let $V \subset \mathcal{B}_f$ be a disk centered at ξ and such that $\mathcal{L}_f(V)$ is compactly contained in V . Therefore $\mathcal{L}_f^{\circ k}(V)$ converges to ξ as $k \rightarrow +\infty$. Let $W \Subset \mathcal{B}_g$ be an arbitrary disk.

Thanks to Proposition 8.11, there exists $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$,

$$\pi_1 \circ F^{\circ(2n+1)}(V \times g^{n^2}(W)) \Subset V,$$

where $\pi_1 : \mathbb{C}^2 \rightarrow \mathbb{C}$ denotes the projection on the first coordinate, $\pi_1(z, w) := z$.

Let U be a connected component of the open set $F^{-n_0^2}(V \times g^{n_0^2}(W))$. Then for every integer $n \geq n_0$, we have

$$(7) \quad F^{\circ n^2}(U) \subseteq V \times g^{n^2}(W).$$

In fact, this holds by assumption for $n = n_0$. Now if the inclusion is true for some $n \geq n_0$, then

$$\begin{aligned} \pi_1 \circ F^{\circ(n+1)^2}(U) &= \pi_1 \circ F^{\circ(2n+1)} \left(F^{\circ n^2}(U) \right) \\ &\subset \pi_1 \circ F^{\circ(2n+1)} \left(V \times g^{n^2}(W) \right) \subset V, \end{aligned}$$

from which (7) follows. This yields that the sequence $(F^{\circ n^2})_{n \geq 0}$ is uniformly bounded, and hence normal, on U . Moreover, any cluster value of this sequence of maps is constant and of the form $(z, 0)$ for some $z \in V$, and $(z, 0)$ is a limit value (associated to a subsequence $(n_k)_{k \in \mathbb{N}}$) if and only if $(\mathcal{L}_f(z), 0)$ is a limit value (associated to the subsequence $(1 + n_k)_{k \in \mathbb{N}}$). Therefore the set of cluster limits is

totally invariant under $\mathcal{L}_f : V \rightarrow V$, and so it must coincide with the attracting fixed point ξ of \mathcal{L}_f . Therefore the sequence $(F^{\circ n^2})_{n \geq 0}$ converges locally uniformly to $(\xi, 0)$ on U .

The sequence $(F^{\circ m})_{m \in \mathbb{N}}$ is locally bounded on U if and only if there exists a subsequence $(m_k)_{k \in \mathbb{N}}$ such that $(F^{\circ m_k}|_U)_{k \in \mathbb{N}}$ has the same property. In fact, since \overline{W} is compact, there exists $R > 0$ such that if $|z| > R$, then for every $w \in W$, (z, w) escapes locally uniformly to infinity under iteration. The domain U is therefore contained in the Fatou set of F .

Let Ω be the component of the Fatou set \mathcal{F}_F containing U . For any integer $j \geq 0$, the sequence of maps $(F^{\circ n^2+j})_{n \in \mathbb{N}}$ converges locally uniformly to $F^{\circ j}(\xi, 0) = (f^{\circ j}(\xi), 0)$ on U and hence on Ω . Therefore the sequence $(F^{\circ n^2})_{n \in \mathbb{N}}$ converges locally uniformly to $(f^{\circ j}(\xi), 0)$ on $F^{\circ j}(\Omega)$. If i, j are nonnegative integers such that $F^{\circ i}(\Omega) = F^{\circ j}(\Omega)$, then $f^{\circ i}(\xi) = f^{\circ j}(\xi)$, and so $i = j$ because ξ cannot be pre-periodic under iteration of f , since it belongs to the parabolic basin \mathcal{B}_f . This proves that Ω is not (pre-)periodic under iteration of F , and so it is a wandering Fatou component for F . \square

We end this subsection with some explicit examples satisfying the assumption of Theorem 8.9.

Example 8.12. Example As a consequence of Proposition 8.10 we obtain that if $f : \mathbb{C} \rightarrow \mathbb{C}$ is the cubic polynomial $f(z) = z + z^2 + az^3$, and g is as in (5), then the polynomial skew-product F defined in (6) admits a wandering Fatou component for $r > 0$ sufficiently small and $a \in D(1 - r, r)$.

It is also interesting to search for real polynomial mappings with wandering Fatou domains intersecting \mathbb{R}^2 . We also have such examples.

Example 8.13. Example [4, Proposition C] Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be the degree 4 polynomial defined by

$$f(z) := z + z^2 + bz^4 \text{ with } b \in \mathbb{R}.$$

There exist parameters $b \in (-8/27, 0)$ such that for g as in (5), the polynomial skew-product F defined in (6) has a wandering Fatou component intersecting \mathbb{R}^2 .

Astorg, Boc-Thaler and Peters constructed in [3] a second example of polynomial skew-products, arising from similar techniques, but with distinctly different dynamical behaviour. Instead of wandering domains arising from a Lavaurs map with an attracting fixed point, they constructed a domain arising from a Lavaurs map with a fixed point of Siegel type.

In [36], Hahn and Peters constructed polynomial automorphisms of \mathbb{C}^4 with wandering Fatou components. Their four-dimensional automorphisms lie in a one-parameter family, depending on a parameter $\delta \in \mathbb{C} \setminus \{0\}$, and degenerating to the two-dimensional polynomial map in equation (5) of Theorem 8.9 as δ converges to 0.

Berger and Biebler recently proved in [10] the existence of a locally dense set of polynomial automorphisms of \mathbb{C}^2 with real coefficients, having a wandering Fatou component. These Fatou components have nonempty real trace and their statistical behaviour is historic with high emergence. The proof is based on a geometric model for parameter families of surface real mappings.

8.4. Elliptic invariant fiber. With Peters we investigated in [51] the case of invariant fibers at the center of a Siegel disk. More precisely, we considered a polynomial skew-product of the form (1) having an *elliptic invariant fiber*. As before, we can assume without loss of generality that the invariant fiber is $\{w = 0\}$ and so we have $g(0) = 0$ and $g'(0) = e^{2\pi i\theta}$ with $\theta \in \mathbb{R} \setminus \mathbb{Q}$. We assume that the origin belongs to a Siegel disk for g and hence g is locally holomorphically linearizable near $w = 0$. Therefore, up to a local change of coordinates we may assume that F is of the form

$$F(z, w) = (f(z, w), \lambda \cdot w),$$

where $f_w(z) := f(z, w)$ is a polynomial in z with coefficients depending holomorphically on w . We assume that the degree of the polynomial f_w is constant near $\{w = 0\}$, and at least 2.

In this case we only have a partial answer to Question 1. In fact, while we already know that the attracting Fatou components of f_0 always bulge, the general situation appears to be more complicated as there might be resonance phenomena. In [51] we proved the following local result, implying that all parabolic Fatou components of a polynomial skew-product with an elliptic invariant fiber bulge if the rotation number satisfies the Brjuno condition.

Proposition 8.14 ([51, Proposition 2]). *Let F be a holomorphic skew-product of the form*

$$(8) \quad F(z, w) = (f_w(z), g(w))$$

with $g(w) = \lambda w + O(w^2)$, $f_0(0) = 0$, and $f'_0(0) = 1$. Assume λ is a Brjuno number. If $f_0(w) \equiv w$, then F is holomorphically linearizable at the origin. If $f_0(z) = z + f_{0,k+1}z^{k+1} + O(z^{k+2})$ with $f_{0,k+1} \neq 0$ for some $k \geq 1$, then for any $h \geq 0$ there exists a local holomorphic change of coordinates near the origin conjugating F to a map of the form $\tilde{F}(z, w) = (\tilde{f}(z, w), \lambda w)$ satisfying

$$(9) \quad \tilde{f}(z, w) = z + f_{0,k+1}z^{k+1} + \dots + f_{0,k+h+1}z^{k+h+1} + \sum_{j \geq h} z^{k+j+2} \alpha_{k+j+2}(w),$$

where, for $j \geq h$, $\alpha_{k+j+2}(w)$ is a holomorphic function in w such that $\alpha_{k+j+2}(0) = f_{0,k+j+2}$.

We recall that a complex number is called *Brjuno* (see [15] for more details) if

$$(10) \quad \sum_{k=0}^{+\infty} \frac{1}{2^k} \log \frac{1}{\omega(2^{k+1})} < +\infty,$$

where $\omega(m) = \min_{2 \leq k \leq m} |\lambda^k - \lambda|$ for any $m \geq 2$, and a quadratic polynomial $\lambda w + w^2$ is linearizable near 0 if and only if λ is Brjuno (see [15, 16, 64]). For higher degree polynomials the Brjuno condition (10) is sufficient, but necessity is still an open question.

The proof of the previous proposition makes use of a procedure inspired by the Poincaré-Dulac normalization process [2, Chapter 4] aiming to conjugate the given polynomial skew-product to a skew-product in a simpler form. At each step of the usual Poincaré-Dulac normalization procedure we can use a polynomial change of coordinates to eliminate all non-resonant monomials of a given degree. A similar idea is used here, and thanks to the skew-product structure of the germ and the Brjuno assumption we are able to eliminate all non-resonant terms of a given degree

in the powers of z by means of changes of coordinates that are polynomial in z with holomorphic coefficients in w . It turns out that there are some differences in between the first degrees as it is pointed out in [51, Section 2].

The arithmetic Brjuno condition is strongly needed in the proof of Proposition 8.14 and it is natural to ask whether parabolic Fatou components of f_0 always bulge, or it is possible to characterize when they bulge. We do not have a complete answer to such question. However we can prove the following result.

Proposition 8.15 ([51, Proposition 10]). *Let $F(z, w) = (f_z(w), g(w))$ be a holomorphic skew-product with an elliptic linearizable invariant fiber. If F does not admit a holomorphic invariant curve on the invariant fiber, then the parabolic Fatou components of f_0 do not bulge.*

Moreover, we can construct examples showing that the Brjuno condition cannot be completely omitted. In fact by taking Cremer numbers, that is $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ and such that

$$\limsup_{m \rightarrow \infty} \frac{1}{m} \log \frac{1}{\omega(m)} = +\infty,$$

we can construct examples of polynomial skew-product not admitting a holomorphic invariant curve on the invariant fiber. An elementary example is the following polynomial skew-product

$$F(z, w) = (z + w + zw, \lambda w)$$

with λ a Cremer number (see [51, Example 8] for details). More generally, arguing as in Cremer's example [23] we can show (see [51, Proposition 9]) the existence of holomorphic skew-products without invariant holomorphic curves of the form $\{w = \varphi(z)\}$, having an elliptic invariant fiber that is not point-wise fixed.

Describing the general situation can also be complicated by resonance phenomena. For example, an invariant fiber at the center of a Siegel disk was used in [17] to construct a non-recurrent Fatou component with limit set isomorphic to a punctured disk, and in that construction the invariant fiber also contains a Siegel disk, but with opposite rotation number. Moreover, it might happen that Fatou components on the invariant fiber do not bulge. For example, we can consider the skew-product

$$F(z, w) = (\lambda z(1 + azw), \lambda^{-1}w),$$

with $a \in \mathbb{C}^*$, $\lambda = e^{2\pi i\theta}$ and $\theta \in \mathbb{R} \setminus \mathbb{Q}$. We have $F(z, 0) = (\lambda z, 0)$, but the Siegel disk around the origin in $\{w = 0\}$ is not bulging, and in fact it follows from [13] and [14] that there exists a Fatou component of parabolic type having on its boundary the origin of \mathbb{C}^2 , which is fixed by F .

We also have an answer to Question 2, under the assumption that the multiplier at the elliptic invariant fiber is Brjuno and all critical points of the polynomial acting on the invariant fiber lie in basins of attracting or parabolic cycles.

Theorem 8.16 ([51, Theorem 1]). *Let F be a polynomial skew-product of the form*

$$(11) \quad F(z, w) = (f_w(z), g(w)),$$

and let $\{w = w_0\}$ be an elliptic invariant fiber with multiplier λ . If λ is Brjuno and all critical points of the polynomial f_{w_0} lie in basins of attracting or parabolic cycles, then all Fatou components of f_{w_0} bulge, and there is a neighbourhood of the invariant fiber $\{w = w_0\}$ in which the only Fatou components of F are the bulging

Fatou components of f_{w_0} . In particular there are no wandering Fatou components in this neighbourhood.

Idea of the proof. The proof relies on the fact that, in dimension 1, if every critical point of the polynomial either (1) lies in the basin of an attracting periodic cycle, (2) lies in the basin of a parabolic periodic cycle, or (3) lies in the Julia set and after finitely many iterates is mapped to a periodic point, then it is possible to construct a conformal metric μ , defined in a backward invariant neighbourhood of the Julia set minus the parabolic periodic orbits, so that f_c is expansive with respect to this metric. It then follows that there can be no wandering Fatou components.

The key step in the proof is that for fibers $\{w = \tilde{w}_0\}$ sufficiently close to the invariant fiber $\{w = w_0\}$ we can define conformal metrics $\mu_{\tilde{w}_0}$ that depend continuously on \tilde{w}_0 , and so that F acts expansively with respect to this family of metrics (see [51, Section 3] for details). That is, for a point $(\tilde{z}_0, \tilde{w}_0) \in \mathbb{C}^2$ lying in the region where the metrics are defined, and for a non-zero tangent vector $\xi \in T_{\tilde{z}_0}(\mathbb{C}_{\tilde{w}_0})$ we have that

$$\mu_{\tilde{w}_1}(\tilde{z}_1, df_{\tilde{w}_0}\xi) > \mu_{\tilde{w}_0}(\tilde{z}_0, \xi),$$

where $(\tilde{z}_1, \tilde{w}_1) = F(\tilde{z}_0, \tilde{w}_0)$. Therefore, if $(\tilde{z}_0, \tilde{w}_0)$ does not lie in one of the bulging Fatou components and its orbit $(\tilde{z}_n, \tilde{w}_n)$ remains in the neighbourhood where the expanding metrics are defined, for any tangent vector $\xi \in T_{\tilde{z}_0}(\mathbb{C}_{\tilde{w}_0})$ we have that

$$\mu_{\tilde{w}_n} df_{\tilde{z}_{n-1}} \cdots df_{\tilde{z}_0} \xi \rightarrow \infty,$$

from which it follows that the family $(F^{o n})_{n \in \mathbb{N}}$ cannot be normal on any neighbourhood of $(\tilde{z}_0, \tilde{w}_0)$. This proves that there are no Fatou components but the bulging components near the invariant fiber $\{w = w_0\}$. \square

It is unclear whether the same techniques could be used to deal with pre-periodic critical points lying on the Julia set. The difficulty is that the property that critical points are eventually mapped onto periodic cycles is not preserved in nearby fibers.

ACKNOWLEDGEMENTS

We thank John H. Hubbard for letting us use the book in preparation [20] to write this survey. The first seven sections are adapted from the first chapter of this book. We also thank Marco Abate and Fabrizio Bianchi for useful comments on the first draft of these notes.

REFERENCES

- [1] Alexander, D.S.: A history of complex dynamics. From Schröder to Fatou and Julia. Aspects of Mathematics, Friedr. Vieweg & Sohn, Braunschweig, (1994)
- [2] Arnold, V.I.: Geometrical Methods in the Theory of Ordinary Differential Equations. Springer Verlag, New York, (1988)
- [3] Astorg, M., Boc-Thaler, L., Peters, H.: Wandering domains arising from Lavaurs maps with Siegel disks, arXiv:1907.04140, to appear in Analysis&PDE
- [4] Astorg, M., Buff, X., Dujardin, R., Peters, H., Raissy, J.: A two-dimensional polynomial mapping with a wandering Fatou component, *Ann. of Math. (2)* **184** no. 1, 263–313 (2016)
- [5] Beardon, A.F.: Iteration of rational functions. Complex analytic dynamical systems. Graduate Texts in Mathematics, 132. Springer-Verlag, New York (1991)
- [6] Bedford, E., Smillie, J.: Polynomial diffeomorphisms of \mathbb{C}^2 : currents, equilibrium measure and hyperbolicity, *Invent. Math.* **103** (1), 69–99 (1991)
- [7] Bedford, E., Smillie, J.: Polynomial diffeomorphisms of \mathbb{C}^2 : II. Stable manifolds and recurrence, *J. Am. Math. Soc.* **4** (4), 657–679 (1991)
- [8] Bedford, E., Smillie, J.: External rays in the dynamics of polynomial automorphisms of \mathbb{C}^2 , In **Contemporary Mathematics**, vol. 222, 41–79. AMS, Providence, RI (1999)
- [9] Bedford, E., Smillie, J., Ueda, T.: Semi-parabolic bifurcations in complex dimension two. *Comm. Math. Phys.* 350, no. 1, 1–29 (2017)
- [10] Berger, P., Biebler, S.: Emergence of wandering stable components, *J. Amer. Math. Soc.* 36 (2023), no. 2, 397–482.
- [11] Bianchi, F.: Parabolic implosion for endomorphisms of \mathbb{C}^2 , *J. Eur. Math. Soc. (JEMS)* 21 no. 12, 3709–3737 (2019)
- [12] Blanchard, P.: Complex analytic dynamics on the Riemann sphere, *Bull. Amer. Math. Soc.*, **11**, 85–141 (1984)
- [13] Bracci, F., Zaitsev, D.: Dynamics of one-resonant biholomorphisms, *J. Eur. Math. Soc. (JEMS)* **15** no. 1, 179–200 (2013)
- [14] Bracci, F., Raissy, J., Zaitsev, D.: Dynamics of multi-resonant biholomorphisms, *Int. Math. Res. Not. IMRN*, no. 20, 4772–4797 (2013)
- [15] Brjuno, A.D.: Analytic form of differential equations. I., *Trans. Mosc. Math. Soc.* **25**, 131–288 (1971)
- [16] Brjuno, A.D.: Analytic form of differential equations. II., *Trans. Mosc. Math. Soc.* **26**, 199–239 (1972)
- [17] Boc-Thaler, L., Fornæss, J.E., Peters, H.: Fatou components with punctured limit sets, *Ergod. Theory Dyn. Syst.* **35**, 1380–1393 (2015)
- [18] Böttcher, L.E.: The principle laws of convergence of iterates and their applications to analysis (Russian), *Izv. Kazan. Fiz.-Mat. Obshch.* **14** 155–234, (1904)
- [19] Buff, X.: Wandering Fatou component for polynomials, *Annales de la faculté des sciences de Toulouse Sér. 6*, 27 no. 2: Numéro spécial à l’occasion de KAWA Complex Analysis Winter school And workshop, 2014-2016, 445–475, (2018)
- [20] Buff, X., Hubbard, J.H.: Dynamics in one complex variable, book in preparation.
- [21] Carleson, L., Gamelin, T.W.: Complex dynamics. Universitext: Tracts in Mathematics. Springer-Verlag, New York (1993)
- [22] Cremer, H.: Über die Schrödersche Funktionalgleichung und das Schwartsche Eckenabbildungsproblem, *Ber. Verh. Sächs. Akad. Wiss. Leipzig, Math.-Phys. Kl.* **84** (1932)
- [23] Cremer, H.: Über die Häufigkeit der Nichtzentren. *Math. Ann.* **115** 573–580 (1938)
- [24] Denker, M., Przytycki, F., Urbański, M.: On the transfer operator for rational functions on the Riemann sphere *Ergodic Theory and Dynamical Systems*, 16(02) 255–266 (1996)
- [25] Devaney R.: An introduction to Chaotic Dynamical Systems, Addison-Wesley (1989)
- [26] Douady, A: Disques de Siegel et anneaux de Herman. *Séminaire Bourbaki*, Vol. 1986/87. Astérisque No. 152–153 (1987), 4, 151–172 (1988)
- [27] Douady, A: Does a Julia set depend continuously on the polynomial? *Complex dynamical systems (Cincinnati, OH, 1994)*, 91–138, *Proc. Sympos. Appl. Math.*, 49, Amer. Math. Soc., Providence, RI (1994)
- [28] Dujardin, R.: A non-laminar dynamical Green current, *Math. Ann.* **365** no. 1-2, 77–91 (2016)
- [29] Dujardin, R., Lyubich, M.: Stability and bifurcations for dissipative polynomial automorphisms of \mathbb{C}^2 . *Invent. Math.* **200** no. 2, 439–511 (2015)

- [30] Fatou, P.: Sur les solutions uniformes de certaines équations fonctionnelles, C.R. Acad. Sci. Paris **143**, 546–548 (1906)
- [31] Fatou, P.: Sur les équations fonctionnelles (Deuxième mémoire), Bull. Soc. Math. France, **48**, 33–94 (1920)
- [32] Fatou, P.: Sur les équations fonctionnelles (Troisième mémoire), Bull. Soc. Math. France, **48**, 208–314 (1920)
- [33] Fornæss, J.E., Sibony, N.: Classification of recurrent domains for some holomorphic maps, Math. Ann. **301** (4), 813–820 (1995)
- [34] Fornæss, J.E., Sibony, N.: Fatou and Julia sets for entire mappings in \mathbb{C}^k . Math. Ann. **311** no. 1, 27–40 (1998)
- [35] Fornæss, J.E., Sibony, N.: Some open problems in higher dimensional complex analysis and complex dynamics. Publ. Mat. **45** no. 2, 529–547 (2001)
- [36] Hahn, D., Peters, H.: A polynomial automorphism with a wandering Fatou component. Adv. Math. **382** (2021), Paper No. 107650, 46 pp.
- [37] Heinemann, S.-M.: Julia sets of skew-products in \mathbb{C}^2 , Kyushu J. Math. **52** (2), 299–329 (1998)
- [38] Ji, Z.: Non-wandering Fatou components for strongly attracting polynomial skew products, J. Geom. Anal. **30**, no. 1, 124–152 (2020)
- [39] Ji, Z.: Non-uniform hyperbolicity in polynomial skew products, to appear in IMRN, (2022), arXiv:1909.06084
- [40] Jonsson, M., *Dynamics of polynomial skew-products on \mathbb{C}^2* , Math. Ann. **314**, 403–447 (1999)
- [41] Julia, G.: Mémoire sur l’itération des fonctions rationnelles, J. Math. Pures Appl. **8** 47–245 (1918)
- [42] Keen, L.: Julia sets, in “Chaos and Fractals, The Mathematics behind the Computer Graphics”, edit. Devaney & Keen, Proc. Symp. Appl. Math., **39**, Amer. Math. Soc., 57–74 (1989)
- [43] Koenigs, G.: Recherches sur les intégrales de certaines équations fonctionnelles. Ann. Sci. École Norm. Sup. (3) **1**, 3–41 (1884)
- [44] Lavaurs, P.: Systèmes dynamiques holomorphes, explosion de points périodiques paraboliques, Ph.D. thesis, Université Paris-Sud, Orsay (1989)
- [45] Lilov, K.: Fatou Theory in Two Dimensions, Ph.D.thesis, University of Michigan (2004)
- [46] Lyubich, M. Y.: Dynamics of rational transformations: topological picture. Uspekhi Mat. Nauk **41**, no. 4(250), 35–95, 239 (1986)
- [47] Lyubich, M. Y., Peters, H.: Classification of invariant Fatou components for dissipative Hénon maps, Geom. Funct. Anal. **24**, 887–915 (2014)
- [48] McMullen, C.T.: Complex Dynamics and Renormalization, Annals of Math Studies vol. 135, Princeton University Press (1994)
- [49] McMullen, C.T.: Renormalization and 3-manifolds which fibers over the circle, Annals of Math Studies vol. 142, Princeton University Press (1996)
- [50] Milnor, J.: Dynamics in one complex variable. Third edition. Annals of Mathematics Studies, **160**. Princeton University Press, Princeton (2006)
- [51] Peters, H., Raissy, J.: Fatou components of polynomial elliptic skew products, Ergodic Theory and Dynamical Systems, **39**, no. 8, 2235–2247 (2019)
- [52] Peters, H., Smit, I.M.: Fatou components of attracting skew products, J. Geom. Anal. **28**, no. 1, 84–110 (2018)
- [53] Peters, H., Vivas, L.: Polynomial skew-products with wandering Fatou-disks, Math. Z. **283** no. 1-2, 349–366 (2016)
- [54] Roeder, R.: A dichotomy for Fatou components of polynomial skew-products, Conform. Geom. Dyn. **15**, 7–19 (2011)
- [55] Rosay, J.P., Rudin, W.: Holomorphic maps from \mathbb{C}^n to \mathbb{C}^n , Trans. Am. Math. Soc. **310**(1), 47–86 (1988)
- [56] Shishikura, M.: Bifurcation of parabolic fixed points. The Mandelbrot set, theme and variations, 325–363, London Math. Soc. Lecture Note Ser., 274, Cambridge Univ. Press, Cambridge, (2000)
- [57] Schröder, E.: Ueber iterirte Functionen. Math. Ann. **3** no. 2, 296–322 (1870)
- [58] Siegel, C.L.: Iteration of analytic functions. Ann. of Math. (2) **43**, 607–612 (1942)
- [59] Steinmetz, N.: Rational iteration. Complex analytic dynamical systems. De Gruyter Studies in Mathematics, **16**. Walter de Gruyter & Co., Berlin (1993)
- [60] Sullivan, D.: Quasiconformal Homeomorphisms and Dynamics I. Solution of the Fatou-Julia Problem on Wandering Domains, Ann. of Math. (2) **122** no. 3, 401–418 (1985)

- [61] Ueda, T.: Fatou sets in complex dynamics on projective spaces, *J. Math. Soc. Jpn.* **46**, 545–555 (1994)
- [62] Vivas, L.: Parametrization of unstable manifolds and Fatou disks for parabolic skew-products. *Complex Anal. Synerg.* 4 no. 1, Paper No. 1 (2018)
- [63] Vivas, L.: Non-autonomous parabolic bifurcation. *Proc. Amer. Math. Soc.* 148, no. 6, 2525–2537 (2020)
- [64] Yoccoz, J.-C.: Théorème de Siegel, nombres de Brjuno et polynômes quadratiques, *Astérisque* **231**, 3–88 (1995)

E-mail address: `xavier.buff@math.univ-toulouse.fr`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR5219, UNIVERSITÉ DE TOULOUSE, CNRS, UPS, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `jasmin.raissy@math.u-bordeaux.fr`

UNIV. BORDEAUX, CNRS, BORDEAUX INP, IMB, UMR 5251, F-33400 TALENCE, FRANCE & INSTITUT UNIVERSITAIRE DE FRANCE (IUF)