

# TOEPLITZ OPERATORS AND SKEW CARLESON MEASURES FOR WEIGHTED BERGMAN SPACES ON STRONGLY PSEUDOCONVEX DOMAINS

MARCO ABATE<sup>†</sup>, SAMUELE MONGODI, AND JASMIN RAISSY<sup>\*</sup>

ABSTRACT. In this paper we study mapping properties of Toeplitz-like operators on weighted Bergman spaces of bounded strongly pseudoconvex domains in  $\mathbb{C}^n$ . In particular we prove that a Toeplitz operator built using as kernel a weighted Bergman kernel of weight  $\beta$  and integrating against a measure  $\mu$  maps continuously (when  $\beta$  is large enough) a weighted Bergman space  $A_{\alpha_1}^{p_1}(D)$  into a weighted Bergman space  $A_{\alpha_2}^{p_2}(D)$  if and only if  $\mu$  is a  $(\lambda, \gamma)$ -skew Carleson measure, where  $\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}$  and  $\gamma = \frac{1}{\lambda} \left( \beta + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right)$ . This theorem generalizes results obtained by Pau and Zhao on the unit ball, and extends and makes more precise results obtained by Abate, Raissy and Saracco on a smaller class of Toeplitz operators on bounded strongly pseudoconvex domains.

## 1. INTRODUCTION

Carleson measures are a powerful tool and an interesting object to study, introduced by Carleson [6] in his celebrated solution of the corona problem. Let  $A$  be a (usually) Banach space of holomorphic functions on a domain  $D \subset \mathbb{C}^n$ ; given  $p > 0$ , a finite positive Borel measure  $\mu$  on  $D$  is a *Carleson measure* for  $A$  and  $p$  if there is a continuous inclusion  $A \hookrightarrow L^p(\mu)$ , that is, if there exists a constant  $C > 0$  such that

$$\forall f \in A \quad \int_D |f|^p d\mu \leq C \|f\|_A^p.$$

We shall also say that  $\mu$  is a *vanishing Carleson measure* for  $A$  and  $p$  if the inclusion  $A \hookrightarrow L^p(\mu)$  is compact.

In this paper we are interested in Carleson measures for weighted Bergman spaces  $A_\beta^p(D)$ , that is spaces of holomorphic functions on a domain  $D \Subset \mathbb{C}^n$  which are  $p$ -integrable with respect to the measure  $\delta^\beta \nu$ , where  $\nu$  is the Lebesgue measure,  $\delta$  is the Euclidean distance from the boundary of  $D$  and  $\beta \in \mathbb{R}$ ; we shall denote by  $A^p(D)$  the (unweighted) Bergman space  $A_0^p(D)$ .

Carleson measures for (possibly weighted) Bergman spaces have been studied by several authors, including Hastings [13], Oleinik and Pavlov [26], Oleinik [25] and Luecking [24] for the unit disk  $\Delta \subset \mathbb{C}$ ; Cima and Wogen [8], Duren and Weir [11], Zhu [31] and Kaptanoğlu [18] for the unit ball  $B^n \subset \mathbb{C}^n$ ; Zhu [30] for bounded symmetric domains; Cima and Mercer [7], Abate and Saracco [3], Abate, Raissy and Saracco [4], Hu, Lv and Zhu [16] and Abate and Raissy [5] for strongly pseudoconvex domains.

One of the reasons of the interest for Carleson measures is that they can be characterized in several different ways, even without any reference to function spaces. A particularly important characterization relies on the intrinsic Kobayashi geometry of the domain  $D \Subset \mathbb{C}^n$ . Given  $z_0 \in D$  and  $r \in (0, 1)$ ,

---

2010 Mathematics Subject Classification: 32A36 (primary), 32A25, 32Q45, 32T15, 46E22, 46E15, 47B35 (secondary).

*Keywords:* Carleson measure; Toeplitz operator; strongly pseudoconvex domain; weighted Bergman space.

<sup>†</sup> Partially supported by PRA grant “Sistemi dinamici in analisi, geometria, logica e meccanica celeste”, University of Pisa.

<sup>\*</sup> Partially supported by the FIRB2012 grant “Differential Geometry and Geometric Function Theory”, RBFR12W1AQ 002.

let  $B_D(z, r)$  denote the Kobayashi ball of  $D$  with center  $z_0$  and radius  $\frac{1}{2} \log \frac{1+r}{1-r}$ . If  $\mu$  is a finite positive Borel measure on  $D$ , for any  $r \in (0, 1)$  and  $\theta \in \mathbb{R}$  we can compare the  $\mu$ -measure and the Lebesgue measure of the Kobayashi balls by using the functions

$$\hat{\mu}_{r,\theta}(z) = \frac{\mu(B_D(z, r))}{\nu(B_D(z, r))^\theta}.$$

It turns out that the behavior of  $\hat{\mu}_{r,\theta}$  can be used to decide whether  $\mu$  is Carleson for a given weighted Bergman space. Indeed we have the following statement:

**Theorem 1.1** (Abate-Raissy-Saracco [4], Hu-Lv-Zhu [16]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex smooth domain and  $\mu$  a finite positive Borel measure on  $D$ . Choose  $0 < p, q < +\infty$  and  $\alpha > -1$ , and denote by  $\delta: D \rightarrow \mathbb{R}^+$  the Euclidean distance from the boundary of  $D$ . Then:*

- (i) *if  $p \leq q$ , then  $\mu$  is a Carleson measure for  $A_\alpha^p(D)$  and  $q$  if and only if  $\hat{\mu}_{r,q/p} \delta^{-\alpha q/p} \in L^\infty(D)$  for some (and hence any)  $r \in (0, 1)$ ;*
- (ii) *if  $p > q$ , then  $\mu$  is a Carleson measure for  $A_\alpha^p(D)$  and  $q$  if and only if  $\hat{\mu}_{r,1} \delta^{-\alpha q/p} \in L^{\frac{p}{p-q}}(D)$  for some (and hence any)  $r \in (0, 1)$ .*

In view of this theorem it is natural to say that a measure  $\mu$  is a  $(\lambda, \alpha)$ -skew Carleson measure if  $\lambda \geq 1$  and  $\hat{\mu}_{r,\lambda} \delta^{-\alpha \lambda} \in L^\infty(D)$ , or if  $\lambda < 1$  and  $\hat{\mu}_{r,1} \delta^{-\alpha \lambda} \in L^{\frac{1}{1-\lambda}}(D)$ . When  $\lambda = 1$  (i.e.,  $p = q$ ) we shall say that  $\mu$  is a  $\alpha$ -Carleson measure.

Other characterizations can be given in terms of  $r$ -lattices and of the Berezin transform of the measure  $\mu$  (see Section 2 of this paper for details); but here we are interested in a different kind of characterization, an application of Carleson measures to mapping properties of Toeplitz operators.

Roughly speaking, a Toeplitz operator is the composition of a projection and a multiplication. More precisely, if  $X$  is a Banach algebra,  $Y \subset X$  a Banach subspace,  $P: X \rightarrow Y$  a linear projection and  $f \in X$ , then the Toeplitz operator  $T_f$  of symbol  $f$  is given by  $T_f(g) = P(fg)$ .

In complex analysis, the most important projection is the Bergman projection  $B$ , which is the orthogonal projection of the space  $L^2(D)$  onto the (unweighted) Bergman space  $A^2(D)$ , where  $D \Subset \mathbb{C}^n$  is a bounded domain. The Bergman projection is an integral operator of the form

$$Bf(z) = \int_D K(z, w) f(w) d\nu(w),$$

where  $K: D \times D \rightarrow \mathbb{C}$  is the Bergman kernel of  $D$ . It turns out that the Bergman projection can be extended to  $L^p(D)$  for all  $p > 0$  and maps  $L^p(D)$  into  $A^p(D)$ . Čučković and McNeal [9] suggested to study the mapping properties of Toeplitz operators, associated to the Bergman projection, of the form

$$T_{\delta^\beta} f(z) = \int_D K(z, w) f(w) \delta(w)^\beta d\nu(w);$$

in particular they were interested in determining for which values of  $\beta \in \mathbb{R}$  the operator  $T_{\delta^\beta}$  would map a Bergman space  $A^p(D)$  into a Bergman space  $A^q(D)$ . In the paper [4] we realized that to properly address Čučković and McNeal's questions it is useful to consider the larger class of Toeplitz operators associated to measures. If  $\mu$  is a finite positive Borel measure on  $D$  then the Toeplitz operator of symbol  $\mu$  is given by

$$T_\mu f(z) = \int_D K(z, w) f(w) d\mu(w);$$

clearly, the Toeplitz operator  $T_{\delta^\beta}$  considered by Čučković and McNeal is the Toeplitz operator of symbol the measure  $\delta^\beta \nu$ . Toeplitz operators with a measure as symbol have been studied, for instance, by Kaptanoğlu [18] on the unit ball of  $\mathbb{C}^n$ , by Li [21] and Li and Lueckling [22] in strongly pseudoconvex

domains, and by Schuster and Varolin [29] in the setting of weighted Bargmann-Fock spaces on  $\mathbb{C}^n$ ; they already noticed relationships between their mapping properties and Carleson properties of  $\mu$ .

In [4] we performed a detailed study of how Carleson properties of  $\mu$  were related to mapping properties of  $T_\mu$ , proving results like the following:

**Theorem 1.2** (Abate-Raissy-Saracco [4]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex smooth domain,  $\mu$  a finite positive Borel measure on  $D$  and take  $1 < p < q < +\infty$ . Then the following assertions are equivalent:*

- (i)  $T_\mu: A^p(D) \rightarrow A^q(D)$  continuously;
- (ii)  $\mu$  is a  $\left(1 + \frac{1}{p} - \frac{1}{q}, 0\right)$ -skew Carleson measure.

In proving this theorem we realized that the natural setting to study the mapping properties of Toeplitz operators of this kind is given by weighted Bergman spaces, and we obtained several results showing that if  $T_\mu$  maps a weighted Bergman space into another weighted Bergman space then  $\mu$  is  $(\lambda, \alpha)$ -skew Carleson for suitable  $\lambda$  and  $\alpha$ , and conversely that if  $\mu$  is  $(\lambda, \alpha)$ -skew Carleson then  $T_\mu$  maps a suitable weighted Bergman space into another suitable weighted Bergman space. Unfortunately, we got only a few clean “if and only if” statements; moreover, we were mainly interested in mapping spaces  $A_\alpha^p(D)$  in spaces  $A_\beta^q(D)$  with  $q \geq p$ , and we did not discuss the case  $p > q$ .

This paper is devoted to prove instead a neat and general “if and only if” statement, following ideas introduced by Pau and Zhao [27] in the unit ball. To do so we proceed by further enlarging the class of Toeplitz operators we are considering. Given  $\beta > -1$ , the orthogonal projection  $P_\beta: L^2(\delta^\beta \nu) \rightarrow A_\beta^2(D)$  is still represented by an integral operator of the form

$$P_\beta f(z) = \int_D K_\beta(z, w) f(w) \delta(w)^\beta d\nu(w),$$

where the *weighted Bergman kernel*  $K_\beta: D \times D \rightarrow \mathbb{C}$  has properties similar to those of the usual Bergman kernel (see Section 2). The *Toeplitz operator*  $T_\mu^\beta$  of *symbol*  $\mu$  and *exponent*  $\beta$  is given by

$$T_\mu^\beta f(z) = \int_D K_\beta(z, w) f(w) d\mu(w).$$

Then the main result of this paper is the following:

**Theorem 1.3.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex smooth domain. Let  $0 < p_1, p_2 < +\infty$  and  $-1 < \alpha_1, \alpha_2 < +\infty$ . Suppose that  $\beta \in \mathbb{R}$  satisfies*

$$n + 1 + \beta > n \max \left\{ 1, \frac{1}{p_j} \right\} + \frac{1 + \alpha_j}{p_j}$$

for  $j = 1, 2$ . Put

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}$$

and, if  $\lambda \neq 0$ , put

$$\gamma = \frac{1}{\lambda} \left( \beta + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then, for any finite positive Borel measure  $\mu$  on  $D$ , the following statements are equivalent:

- (i)  $T_\mu^\beta: A_{\alpha_1}^{p_1}(D) \rightarrow A_{\alpha_2}^{p_2}(D)$  continuously;
- (ii) the measure  $\mu$  is a  $(\lambda, \gamma)$ -skew Carleson measure.

In particular, Theorem 1.2 is now obtained as a consequence of Theorem 1.3 by taking  $\alpha_1 = \alpha_2 = \beta = 0$  and  $1 < p_1 < p_2 < +\infty$ .

The paper is structured as follows. In Section 2 we collect a number of preliminary results, on the Kobayashi geometry of strongly pseudoconvex domains, on the weighted Bergman kernels, and on the known characterizations of skew Carleson measures. Section 3 is devoted to the proof of Theorem 1.3, while in Section 4 we prove a version of Theorem 1.3 for vanishing skew Carleson measures, showing that (under the same hypotheses on the parameters)  $T_\mu^\beta: A_{\alpha_1}^{p_1}(D) \rightarrow A_{\alpha_2}^{p_2}(D)$  is compact if and only if the measure  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure.

## 2. PRELIMINARY RESULTS

In this section we collect definitions and preliminary results that we shall use in the rest of the paper.

From now on,  $D \Subset \mathbb{C}^n$  will be a bounded strongly pseudoconvex domain in  $\mathbb{C}^n$  with smooth  $C^\infty$  boundary. Furthermore, we shall use the following notations:

- $\delta: D \rightarrow \mathbb{R}^+$  will denote the Euclidean distance from the boundary of  $D$ , that is  $\delta(z) = d(z, \partial D)$ ;
- given two non-negative functions  $f, g: D \rightarrow \mathbb{R}^+$  we shall write  $f \preceq g$  to say that there is  $C > 0$  such that  $f(z) \leq Cg(z)$  for all  $z \in D$  (the constant  $C$  is independent of  $z \in D$ , but it might depend on other parameters, such as  $r, \theta$ , etc.);
- given two strictly positive functions  $f, g: D \rightarrow \mathbb{R}^+$  we shall write  $f \approx g$  if  $f \preceq g$  and  $g \preceq f$ , that is if there is  $C > 0$  such that  $C^{-1}g(z) \leq f(z) \leq Cg(z)$  for all  $z \in D$ ;
- $\nu$  will be the Lebesgue measure;
- $\mathcal{O}(D)$  will denote the space of holomorphic functions on  $D$ , endowed with the topology of uniform convergence on compact subsets;
- given  $0 < p < +\infty$ , the *Bergman space*  $A^p(D)$  is the (Banach if  $p \geq 1$ ) space  $L^p(D) \cap \mathcal{O}(D)$ , endowed with the  $L^p$ -norm;
- more generally, if  $\mu$  is a positive finite Borel measure on  $D$  and  $0 < p < +\infty$  we shall denote by  $L^p(\mu)$  the set of complex-valued  $\mu$ -measurable functions  $f: D \rightarrow \mathbb{C}$  such that

$$\|f\|_{p,\mu} := \left[ \int_D |f(z)|^p d\mu(z) \right]^{1/p} < +\infty;$$

- if  $\alpha > -1$  we shall write  $\nu_\alpha = \delta^\alpha \nu$ , we shall denote by  $A_\alpha^p(D)$  the *weighted Bergman space*

$$A_\alpha^p(D) = L^p(\delta^\alpha \nu) \cap \mathcal{O}(D),$$

and we shall write  $\|\cdot\|_{p,\alpha}$  instead of  $\|\cdot\|_{p,\delta^\alpha \nu}$ ;

- $K: D \times D \rightarrow \mathbb{C}$  will be the Bergman kernel of  $D$ , and for each  $z_0 \in D$  we shall denote by  $k_{z_0}: D \rightarrow \mathbb{C}$  the *normalized Bergman kernel* defined by

$$k_{z_0}(z) = \frac{K(z, z_0)}{\sqrt{K(z_0, z_0)}} = \frac{K(z, z_0)}{\|K(\cdot, z_0)\|_2};$$

- given  $r \in (0, 1)$  and  $z_0 \in D$ , we shall denote by  $B_D(z_0, r)$  the Kobayashi ball of center  $z_0$  and radius  $\frac{1}{2} \log \frac{1+r}{1-r}$ .

We refer to, e.g., [1, 2, 17, 19], for definitions, basic properties and applications to geometric function theory of the Kobayashi distance; and to [14, 15, 20, 28] for definitions and basic properties of the Bergman kernel.

Let us now recall a few results we shall need on the Kobayashi geometry of strongly pseudoconvex domains.

**Lemma 2.1** ([3, Lemma 2.2]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Then there is  $C > 0$  such that*

$$\frac{1-r}{C} \delta(z_0) \leq \delta(z) \leq \frac{C}{1-r} \delta(z_0)$$

for all  $r \in (0, 1)$ ,  $z_0 \in D$  and  $z \in B_D(z_0, r)$ .

**Lemma 2.2.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain,  $\beta \in \mathbb{R}$  and  $r \in (0, 1)$ . Then*

$$\nu_\beta(B_D(\cdot, r)) \approx \delta^{n+1+\beta},$$

where the constant depends on  $r$ .

*Proof.* For  $\beta = 0$  the result can be found in [21, Corollary 7] and [3, Lemma 2.1]. If  $\beta \neq 0$  Lemma 2.1 yields

$$\nu_\beta(B_D(z_0, r)) = \int_{B_D(z_0, r)} \delta(z)^\beta d\nu(z) \approx \delta(z_0)^\beta \nu(B_D(z_0, r))$$

and we are done.  $\square$

We shall also need the existence of suitable coverings by Kobayashi balls. Recall that for a bounded domain  $D \Subset \mathbb{C}^n$ , given  $r > 0$ , a  $r$ -lattice in  $D$  is a sequence  $\{a_k\} \subset D$  such that  $D = \bigcup_k B_D(a_k, r)$  and there exists  $m > 0$  such that any point in  $D$  belongs to at most  $m$  balls of the form  $B_D(a_k, R)$ , where  $R = \frac{1}{2}(1+r)$ .

The existence of  $r$ -lattices in bounded strongly pseudoconvex domains is ensured by the following result:

**Lemma 2.3** ([3, Lemma 2.5]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Then for every  $r \in (0, 1)$  there exists an  $r$ -lattice in  $D$ .*

We shall use a submean estimate for nonnegative plurisubharmonic functions on Kobayashi balls:

**Lemma 2.4** ([3, Corollaries 2.7 and 2.8]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Given  $r \in (0, 1)$ , set  $R = \frac{1}{2}(1+r) \in (0, 1)$ . Then there exists a constant  $K_r > 0$  depending on  $r$  such that*

$$\forall z_0 \in D \quad \chi(z_0) \leq \frac{K_r}{\nu(B_D(z_0, r))} \int_{B_D(z_0, r)} \chi d\nu$$

and

$$\forall z_0 \in D \quad \forall z \in B_D(z_0, r) \quad \chi(z) \leq \frac{K_r}{\nu(B_D(z_0, R))} \int_{B_D(z_0, R)} \chi d\nu$$

for every nonnegative plurisubharmonic function  $\chi: D \rightarrow \mathbb{R}^+$ .

Now we collect a few results on the weighted Bergman kernels. Given  $\beta > -1$ , the *weighted Bergman projection* is the orthogonal projection  $P_\beta: L^2(\nu_\beta) \rightarrow A_\beta^2(D)$ , where  $\nu_\beta = \delta^\beta \nu$ . It is known (see, e.g., [12]), that there exists a function  $K_\beta: D \times D \rightarrow \mathbb{C}$  such that

$$P_\beta f(z) = \int_D K_\beta(z, w) f(w) \delta(w)^\beta d\nu(w)$$

for all  $f \in L^2(\nu_\beta)$ . Moreover,  $K_\beta(z, w)$  is holomorphic in  $z$ , we have  $K_\beta(w, z) = \overline{K_\beta(z, w)}$  and

$$f(z) = \int_D K_\beta(z, w) f(w) \delta(w)^\beta d\nu(w)$$

for all  $f \in A_\beta^2(D)$ . The function  $K_\beta$  is called the *weighted Bergman kernel* of  $D$ . For  $a \in D$ , the *normalized weighted Bergman kernel* of  $D$  is

$$k_{\beta,a}(z) = \frac{K_\beta(z,a)}{\sqrt{K_\beta(a,a)}}.$$

When  $\beta = 0$  we recover the usual Bergman kernel, and we shall write  $K$ , respectively  $k_a$ , instead of  $K_0$ , respectively  $k_{0,a}$ .

We shall need a few estimates on the behaviour of the weighted Bergman kernel. They are analogous to the classical estimates for the Bergman kernel and follow from the results obtained by Engliš [12] on the asymptotic behaviour of the weighted Bergman kernel. The first one is the following.

**Lemma 2.5.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain and let  $\beta > -1$ . Then*

$$\|K_\beta(\cdot, z_0)\|_{2,\beta} = \sqrt{K_\beta(z_0, z_0)} \approx \delta(z_0)^{-(n+1+\beta)/2} \quad \text{and} \quad \|k_{\beta,z_0}\|_{2,\beta} \equiv 1$$

for all  $z_0 \in D$ .

*Proof.* The first equality, and hence the result for  $k_{\beta,z_0}$ , is well-known, as well as the whole statement for  $\beta = 0$  (see, e.g., [14]).

If  $\beta \neq 0$ , then thanks to the results in [12], the weighted Bergman kernel is smooth outside the boundary diagonal; so, in particular, if  $z_0 \in D' \Subset D$  the norm  $\|K_\beta(\cdot, z_0)\|_{p,\beta}$  is bounded by a constant depending only on  $D'$ ,  $p$  and  $\beta$ .

Therefore, we only have to estimate the boundary behaviour. Let  $q \in \partial D$  and let  $U$  be a neighbourhood of  $q$  with coordinates  $(z', z_n) = (z_1, \dots, z_n)$  centered in  $q$  such that

$$D \cap U = \{(z', z_n) \in U : \operatorname{Re}(z_n) > \psi(z')\}$$

where  $-\psi$  is strongly plurisubharmonic with  $\nabla\psi \neq 0$ . Set  $r(z) = \operatorname{Re}(z_n) - \psi(z')$ . We consider an almost-sesquianalytic extension of  $r(z)$  on  $U \times U$ , i.e., a function, which we denote again by  $r$ , such that:

- $r(z, w) = \overline{r(w, z)}$ ,
- the first derivatives of  $r$  with respect to  $\bar{z}$  and  $w$  vanish at infinite order along  $z = w$ ,
- $r(z, z) = r(z)$ .

It easily follows from these properties that

$$\frac{\partial}{\partial z_j} r(O) = \frac{\partial}{\partial z_j} r(O, O),$$

and similarly for the other derivatives. Therefore we have

$$|2r(z, w) - r(z) - r(w)| = c_1 |z_n - w_n| + \sum_{j=1}^{n-1} c_2^j |z_j - w_j|^2 + O(\|z - w\|^3).$$

Moreover  $|2r(z, w) - r(z) - r(w)|$  is positive outside  $z = w$ , and so  $c_1 > 0$ . Therefore in a neighbourhood of  $(O, O)$  we have that

$$|r(z, w)| \approx \left( r(z) + r(w) + |z_n - w_n| + \sum_{j=1}^{n-1} |z_j - w_j|^2 \right).$$

The results in [12] imply that  $K_\beta(z, w)$  is asymptotic to  $c(z, w)r(z, w)^{-n-1-\beta}$  for a suitable function  $c \in \mathcal{C}^\infty(\overline{D} \times \overline{D})$ . Therefore on  $U$  we have

$$|K_\beta(z, w)| \approx \left( r(z) + r(w) + |z_n - w_n| + \sum_{j=1}^{n-1} |z_j - w_j|^2 \right)^{-n-1-\beta}.$$

Thus, following the same proof as in the classical case, we obtain the assertion.  $\square$

A similar estimate, but with uniform constants on Kobayashi balls, is the following.

**Lemma 2.6.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain and let  $\beta > -1$ . Then for every  $r \in (0, 1)$  there exist  $c_r > 0$  and  $\delta_r > 0$  such that if  $z_0 \in D$  satisfies  $\delta(z_0) < \delta_r$  then*

$$\frac{c_r}{\delta(z_0)^{n+1+\beta}} \leq |K_\beta(z, z_0)| \leq \frac{1}{c_r \delta(z_0)^{n+1+\beta}}$$

and

$$\frac{c_r}{\delta(z_0)^{n+1+\beta}} \leq |k_{\beta, z_0}(z)|^2 \leq \frac{1}{c_r \delta(z_0)^{n+1+\beta}}$$

for all  $z \in B_D(z_0, r)$ .

*Proof.* If  $\beta = 0$  then this is proven in [21, Theorem 12] and [3, Lemma 3.2 and Corollary 3.3]. If  $\beta \neq 0$ , then thanks to the results in [12], we have that

$$(1) \quad K_\beta(z, z_0) \approx c(z, z_0) \left( r(z) + r(z_0) + |z_n - z_{0,n}| + \sum_{j=1}^{n-1} |z_j - z_{0,j}|^2 \right)^{-(n+1+\beta)}$$

in suitable local coordinates around a point of the boundary diagonal, i.e., if  $d(z_0, \partial D)$ ,  $d(z, \partial D)$  and  $\|z - z_0\|$  are small enough. By the completeness of the Kobayashi metric, there exists  $\delta_r > 0$  such that every  $z \in B_D(z_0, r)$  satisfies such condition if  $\delta(z_0) < \delta_r$ . The assertion then follows by arguing as in [21, Theorem 12] or as in [3, Lemma 3.2 and Corollary 3.3].  $\square$

**Remark 2.1.** Note that in the previous lemma the estimates from above hold even when  $\delta(z_0) \geq \delta_r$ , possibly with a different constant  $c_r$ . Indeed, when  $\delta(z_0) \geq \delta_r$  and  $z \in B_D(z_0, r)$  by Lemma 2.1 there is  $\tilde{\delta}_r > 0$  such that  $\delta(z) \geq \tilde{\delta}_r$ ; as a consequence we can find  $M_r > 0$  such that  $|K_\beta(z, z_0)| \leq M_r$  as soon as  $\delta(z_0) \geq \delta_r$  and  $z \in B_D(z_0, r)$ , and the assertion follows from the fact that  $D$  is a bounded domain.

A very useful integral estimate generalizing the analogous ones for the unweighted Bergman kernel (see [21, Corollary 11, Theorem 13] and [4, Theorem 2.7]) is the following:

**Theorem 2.7.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain,  $z_0 \in D$  and  $\alpha, \beta > -1$ . Then for  $0 < p < +\infty$  and  $\alpha - \beta < (n + \beta + 1)(p - 1)$  we have*

$$\int_D |K_\beta(\zeta, z_0)|^p \delta(\zeta)^\alpha d\nu(\zeta) \preceq \delta(z_0)^{\alpha - \beta - (n + \beta + 1)(p - 1)}.$$

In particular,

$$\|K(\cdot, z_0)\|_{p, \alpha} \preceq \delta(z_0)^{\frac{n+1+\alpha}{p} - (n+1+\beta)}.$$

*Proof.* If  $\beta = 0$  then this is proven in [21, Corollary 11, Theorem 13] and [4, Theorem 2.7]. If  $\beta \neq 0$ , then it suffices to use (1) and follow the same proof as in the unweighted case.  $\square$

Finally, the normalized Bergman kernel can be used to build functions belonging to suitable weighted Bergman spaces:

**Lemma 2.8.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain, and  $\beta > -1$ . Given  $0 < p < +\infty$  and  $-1 < \alpha < \min\{(n + \beta + 1)p - np - 1, (n + \beta + 1)p - n - 1\}$ , set*

$$\tau = \frac{n + 1 + \beta}{2} - \frac{n + 1 + \alpha}{p}.$$

For each  $a \in D$  set  $f_a = \delta(a)^\tau k_{\beta,a}$ . Let  $\{a_k\}$  be an  $r$ -lattice and  $\mathbf{c} = \{c_k\} \in \ell^p$ , and put

$$f = \sum_{k=0}^{\infty} c_k f_{a_k}.$$

Then  $f \in A_\alpha^p(D)$  with  $\|f\|_{p,\alpha} \preceq \|\mathbf{c}\|_p$ .

*Proof.* If  $\beta = 0$  then this is a consequence of [16, Lemma 2.6]. If  $\beta \neq 0$ , then it suffices to use the estimates given by Theorem 2.7 and follow the same proof as in the unweighted case.  $\square$

We also need to recall a few definitions and results about Carleson measures.

**Definition 2.9.** Let  $0 < p, q < +\infty$  and  $\alpha > -1$ . A  $(p, q; \alpha)$ -skew Carleson measure is a finite positive Borel measure  $\mu$  such that

$$\int_D |f(z)|^q d\mu(z) \preceq \|f\|_{p,\alpha}^q$$

for all  $f \in A_\alpha^p(D)$ . In other words,  $\mu$  is  $(p, q; \alpha)$ -skew Carleson if  $A_\alpha^p(D) \hookrightarrow L^q(\mu)$  continuously. In this case we shall denote by  $\|\mu\|_{p,q;\alpha}$  the operator norm of the inclusion  $A_\alpha^p(D) \hookrightarrow L^q(\mu)$ . Furthermore, a  $(p, q; \alpha)$ -skew Carleson measure is *vanishing* if

$$\lim_{j \rightarrow +\infty} \int_D |f_j(z)|^q d\mu(z) = 0$$

for any bounded sequence  $\{f_j\}_{j \in \mathbb{N}} \subset A_\alpha^p(D)$  converging to 0 uniformly on any compact subset of  $D$ . For  $p \geq 1$ ,  $\mu$  is a vanishing  $(p, q; \alpha)$ -skew Carleson if and only if  $A_\alpha^p(D) \hookrightarrow L^q(\mu)$  compactly (see, e.g., [4, Lemma 4.5]).

**Remark 2.2.** When  $p = q$  we recover the usual (non-skew) notion of Carleson measure for  $A_\alpha^p(D)$ .

**Definition 2.10.** Let  $\theta \in \mathbb{R}$ , and let  $\mu$  be a finite positive Borel measure on  $D$ . Given  $r \in (0, 1)$ , let  $\hat{\mu}_{r,\theta}: D \rightarrow \mathbb{R}$  be defined by

$$\hat{\mu}_{r,\theta}(z) = \frac{\mu(B_D(z, r))}{\nu(B_D(z, r))^\theta};$$

we shall write  $\hat{\mu}_r$  for  $\hat{\mu}_{r,1}$ .

We say that  $\mu$  is a *geometric  $\theta$ -Carleson measure* if  $\hat{\mu}_{r,\theta} \in L^\infty(D)$  for all  $r \in (0, 1)$ , that is if for every  $r > 0$  we have

$$\mu(B_D(z, r)) \preceq \nu(B_D(z, r))^\theta$$

for all  $z \in D$ , where the constant depends only on  $r$ .

Furthermore, we shall say that  $\mu$  is a *geometric vanishing  $\theta$ -Carleson measure* if

$$\lim_{z \rightarrow \partial D} \hat{\mu}_{r,\theta}(z) = 0$$

for all  $r \in (0, 1)$ .



Notice that Lemma 2.2 yields

$$(2) \quad \hat{\mu}_{r,\theta} \approx \delta^{-(n+1)(\theta-1)} \hat{\mu}_r .$$

In [4] we proved (among other things) that, if  $p \geq 1$ , a measure  $\mu$  is  $(p, p; \alpha)$ -skew Carleson if and only if it is geometric  $\theta$ -Carleson, where  $\theta = 1 + \frac{\alpha}{n+1}$ . Hu, Lv and Zhu in [16] have given a similar geometric characterization of  $(p, q; \alpha)$ -skew Carleson measures for all values of  $p$  and  $q$ ; to recall their results we need another definition.

**Definition 2.11.** Let  $\mu$  be a finite positive Borel measure on  $D$ , and  $s > 0$ . The *Berezin transform* of level  $s$  of  $\mu$  is the function  $B^s \mu: D \rightarrow \mathbb{R}^+ \cup \{+\infty\}$  given by

$$B^s \mu(z) = \int_D |k_z(w)|^s d\mu(w) .$$

The geometric characterization of  $(p, q; \alpha)$ -skew Carleson measures is different according to whether  $p \leq q$  or  $p > q$ . We first recall the characterization for the case  $p \leq q$ .

**Theorem 2.12** ([16, Theorem 3.1], [5, Theorem 2.15]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $0 < p \leq q < +\infty$  and  $\alpha > -1$ ; set  $\theta = 1 + \frac{\alpha}{n+1}$ . Then the following assertions are equivalent:*

- (i)  $\mu$  is a  $(p, q; \alpha)$ -skew Carleson measure;
- (ii)  $\mu$  is a geometric  $\frac{q}{p}\theta$ -Carleson measure;
- (iii) there exists  $r_0 \in (0, 1)$  such that  $\hat{\mu}_{r_0, \frac{q}{p}\theta} \in L^\infty(D)$ ;
- (iv) for some (and hence any)  $r \in (0, 1)$  we have  $\hat{\mu}_{r, \frac{q}{p}} \delta^{-\alpha \frac{q}{p}} \in L^\infty(D)$ ;
- (v) for some (and hence any)  $r \in (0, 1)$  and some (and hence any)  $r$ -lattice  $\{a_k\}$  in  $D$  we have

$$\forall k \in \mathbb{N} \quad \mu(B_D(a_k, r)) \preceq \nu(B_D(a_k, r))^{\frac{q}{p}\theta} ;$$

- (vi) for some (and hence all)  $s > \theta \frac{q}{p}$  we have

$$B^s \mu \preceq \delta^{(n+1)(\theta \frac{q}{p} - \frac{s}{2})} ;$$

Moreover we have

$$(3) \quad \|\mu\|_{p,q;\alpha} \approx \|\hat{\mu}_{r, \frac{q}{p}\theta}\|_\infty \approx \|\hat{\mu}_{r, \frac{q}{p}} \delta^{-\alpha \frac{q}{p}}\|_\infty \approx \|\delta^{(n+1)(\frac{s}{2} - \theta \frac{q}{p})} B^s \mu\|_\infty .$$

The geometric characterization of  $(p, q; \alpha)$ -skew Carleson measures when  $p > q$  has a slightly different flavor.

**Theorem 2.13** ([16, Theorem 3.3], [5, Theorem 2.16]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $0 < q < p < +\infty$  and  $\alpha > -1$ ; put  $\theta = 1 + \frac{\alpha}{n+1}$ . Then the following assertions are equivalent:*

- (i)  $\mu$  is a  $(p, q; \alpha)$ -skew Carleson measure;
- (ii)  $\mu$  is a vanishing  $(p, q, \alpha)$ -skew Carleson measure;
- (iii)  $\hat{\mu}_r \delta^{-\alpha \frac{q}{p}} \in L^{\frac{p}{p-q}}(D)$  for some (and hence any)  $r \in (0, 1)$ ;
- (iv) for some (and hence any)  $r \in (0, 1)$  and for some (and hence any)  $r$ -lattice  $\{a_k\}$  in  $D$  we have  $\{\hat{\mu}_{r, \theta \frac{q}{p}}(a_k)\} \in \ell^{\frac{p}{p-q}}$ ;
- (v) for some (and hence all)  $s > \theta \frac{q}{p} + \frac{n}{n+1} \left(1 - \frac{q}{p}\right)$  we have

$$\delta^{-(n+1)(\theta \frac{q}{p} - \frac{s}{2} + \frac{p-q}{p})} B^s \mu \in L^{\frac{p}{p-q}}(D) ;$$

Moreover we have

$$(4) \quad \|\mu\|_{p,q;\alpha} \approx \|\hat{\mu}_r \delta^{-\alpha \frac{q}{p}}\|_{\frac{p}{p-q}} \approx \|\hat{\mu}_{r,\theta \frac{q}{p}}(a_k)\|_{\frac{p}{p-q}} \approx \|\delta^{-(n+1)(\theta \frac{q}{p} - \frac{s}{2} + \frac{p-q}{p})} B^s \mu\|_{\frac{p}{p-q}}.$$

We also have a geometric characterization of vanishing  $(p, q; \alpha)$ -skew Carleson measures when  $p \leq q$ :

**Theorem 2.14** ([16, Theorem 3.1], [4, Theorem 4.10]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $0 < p \leq q < +\infty$  and  $\alpha > -1$ ; set  $\theta = 1 + \frac{\alpha}{n+1}$ . Then the following assertions are equivalent:*

- (i)  $\mu$  is a vanishing  $(p, q; \alpha)$ -skew Carleson measure;
- (ii)  $\mu$  is a geometric vanishing  $\frac{q}{p}\theta$ -Carleson measure;
- (iii) there exists  $r_0 \in (0, 1)$  such that  $\lim_{z \rightarrow \partial D} \hat{\mu}_{r_0, \frac{q}{p}}(z) = 0$ ;
- (iv) for some (and hence any)  $r \in (0, 1)$  we have  $\lim_{z \rightarrow \partial D} \hat{\mu}_{r, \frac{q}{p}}(z) \delta(z)^{-\alpha \frac{q}{p}} = 0$ ;
- (v) for some (and hence any)  $r \in (0, 1)$  and some (and hence any)  $r$ -lattice  $\{a_k\}$  in  $D$  we have

$$\lim_{k \rightarrow +\infty} \hat{\mu}_{r_0, \frac{q}{p}}(a_k) = \lim_{k \rightarrow +\infty} \hat{\mu}_{r, \frac{q}{p}}(a_k) \delta(a_k)^{-\alpha \frac{q}{p}} = 0;$$

- (vi) for some (and hence all)  $s > \theta \frac{q}{p}$  we have

$$\lim_{z \rightarrow \partial D} \delta(z)^{(n+1)(\frac{s}{2} - \theta \frac{q}{p})} B^s \mu(z) = 0.$$

A consequence of these theorems is that the property of being  $(p, q; \alpha)$ -skew Carleson actually depends only on the quotient  $q/p$  and on  $\alpha$ . We shall then introduce the following definition:

**Definition 2.15.** Take  $\lambda, \alpha \in \mathbb{R}$ . A finite positive Borel measure  $\mu$  on  $D$  is a  $(\lambda, \alpha)$ -skew Carleson measure if

- $\lambda \geq 1$  and  $\hat{\mu}_{r,\lambda} \delta^{-\alpha \lambda} \in L^\infty(D)$  for some (and hence any)  $r \in (0, 1)$ , and we shall put  $\|\mu\|_{\lambda,\alpha} = \|\hat{\mu}_{r,\lambda} \delta^{-\alpha \lambda}\|_\infty$ ; or,
- $\lambda < 1$  and  $\hat{\mu}_r \delta^{-\alpha \lambda} \in L^{\frac{1}{1-\lambda}}(D)$  for some (and hence any)  $r \in (0, 1)$ , and we shall put  $\|\mu\|_{\lambda,\alpha} = \|\hat{\mu}_r \delta^{-\alpha \lambda}\|_{\frac{1}{1-\lambda}}$ .

Notice that by [16, Lemma 2.3] different  $r$ 's yield equivalent norms.

Furthermore we say that  $\mu$  is *vanishing  $(\lambda, \alpha)$ -skew Carleson measure* if

- $\lambda \geq 1$  and  $\lim_{z \rightarrow \partial D} \hat{\mu}_{r,\lambda}(z) \delta(z)^{-\alpha \lambda} = 0$  for some (and hence any)  $r \in (0, 1)$ ; or,
- $\lambda < 1$  and  $\mu$  is a  $(\lambda, \alpha)$ -skew Carleson measure.

So a measure is (vanishing)  $(p, q; \alpha)$ -skew Carleson if and only if it is (vanishing)  $(q/p, \alpha)$ -skew Carleson. Notice that the definition of  $(0, \alpha)$ -skew Carleson does not depend on  $\alpha$ .

This definition has the following easy (but useful) consequence.

**Lemma 2.16** ([5, Lemma 2.18]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain,  $\lambda > 0$  and  $\alpha > -1$ . Let  $\mu$  be a  $(\lambda, \alpha)$ -skew Carleson measure, and  $\beta > -\lambda(\alpha + 1)$ . Then  $\mu_\beta = \delta^\beta \mu$  is a  $(\lambda, \alpha + \frac{\beta}{\lambda})$ -skew Carleson measure with  $\|\mu_\beta\|_{\lambda, \alpha + \frac{\beta}{\lambda}} \approx \|\mu\|_{\lambda, \alpha}$ .*

We end this section by recalling the main result in [5], which gives a characterisation of  $(\lambda, \gamma)$ -skew Carleson measures on bounded strongly pseudoconvex domain through products of functions in weighted Bergman spaces.

**Theorem 2.17** ([5, Theorem 1.1]). *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain, and let  $\mu$  be a positive finite Borel measure on  $D$ . Fix an integer  $k \geq 1$ , and let  $0 < p_j, q_j < +\infty$  and  $-1 < \alpha_j < +\infty$  be given for  $j = 1, \dots, k$ . Set*

$$\lambda = \sum_{j=1}^k \frac{q_j}{p_j} \quad \text{and} \quad \gamma = \frac{1}{\lambda} \sum_{j=1}^k \frac{\alpha_j q_j}{p_j}.$$

*Then  $\mu$  is a  $(\lambda, \gamma)$ -skew Carleson measure if and only if there exists  $C > 0$  such that*

$$(5) \quad \int_D \prod_{j=1}^k |f_j(z)|^{q_j} d\mu(z) \leq C \prod_{j=1}^k \|f_j\|_{p_j, \alpha_j}^{q_j}$$

*for any  $f_j \in A_{\alpha_j}^{p_j}(D)$ .*

### 3. TOEPLITZ OPERATORS AND SKEW CARLESON MEASURES ON WEIGHTED BERGMAN SPACES

This section is devoted to the proof of our main Theorem 1.3. We shall need the following preliminary result:

**Lemma 3.1.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $1 < p < +\infty$ ,  $-1 < \alpha, \alpha' < +\infty$  and put*

$$\beta = \frac{\alpha}{p} + \frac{\alpha'}{p'},$$

*where  $p'$  the conjugate exponent of  $p$ . Then the functional*

$$(f, g)_\beta = \int_D f(z) \overline{g(z)} d\nu_\beta(z)$$

*is a duality pairing between  $A_\alpha^p(D)$  and  $A_{\alpha'}^{p'}(D)$ , where  $\nu_\beta = \delta^\beta \nu$ .*

*Proof.* The continuous dual of  $A_\alpha^p(D)$  is  $A_{\alpha'}^{p'}(D)$ , with the usual pairing

$$\langle f, h \rangle = \int_D f(z) \overline{h(z)} d\nu_\alpha(z).$$

Therefore

$$(f, g)_\beta = \int_D f(z) \overline{g(z)} d\nu_\beta(z) = \int_D f(z) \overline{g(z)} \delta(z)^{\beta-\alpha} d\nu_\alpha(z) = \langle f, g \delta^{\beta-\alpha} \rangle$$

is a duality pairing between  $A_\alpha^p(D)$  and  $A_{\alpha'}^{p'}(D)$  as soon as  $h = g \delta^{\beta-\alpha} \in A_{\alpha'}^{p'}(D)$ , i.e., as soon as

$$\int_D |g(z)|^{p'} \delta(z)^{(\beta-\alpha)p'} d\nu_\alpha(z) < +\infty$$

which is true because the choice of  $\beta$  yields  $(\beta - \alpha)p' + \alpha = \alpha'$ . □

Now we can prove Theorem 1.3:

**Theorem 3.2.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $0 < p_1, p_2 < +\infty$  and  $-1 < \alpha_1, \alpha_2 < +\infty$ . Suppose that  $\beta \in \mathbb{R}$  satisfies*

$$(6) \quad n + 1 + \beta > n \max \left\{ 1, \frac{1}{p_j} \right\} + \frac{1 + \alpha_j}{p_j}$$

*for  $j = 1, 2$ . Put*

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}$$

and, if  $\lambda \neq 0$ , put

$$\gamma = \frac{1}{\lambda} \left( \beta + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then for any positive Borel measure  $\mu$  on  $D$  the following statements are equivalent:

- (i)  $T_\mu^\beta : A_{\alpha_1}^{p_1}(D) \rightarrow A_{\alpha_2}^{p_2}(D)$  continuously;
- (ii)  $\mu$  is a  $(\lambda, \gamma)$ -skew Carleson measure.

Moreover, one has

$$\|T_\mu^\beta\|_{A_{\alpha_1}^{p_1}(D) \rightarrow A_{\alpha_2}^{p_2}(D)} \approx \|\mu\|_{\lambda, \gamma}.$$

*Proof.* The proof is divided into several cases.

(i) $\Rightarrow$ (ii) We consider two cases:  $\lambda \geq 1$  and  $\lambda < 1$ .

*Case 1.* Assume  $\lambda \geq 1$ . Let  $a \in D$  and consider  $f_a = K_\beta(\cdot, a)$ . By (6) with  $j = 1$ , we get  $(n+1+\beta)p_1 > n+1+\alpha_1$ , which is equivalent to  $\alpha_1 - \beta < (n+\beta+1)(p_1-1)$ , so, by Theorem 2.7, for  $a \in D$  we have that

$$(7) \quad \|K_\beta(\cdot, a)\|_{p_1, \alpha_1}^{p_1} \leq \delta(a)^{n+1+\alpha_1-(n+1+\beta)p_1};$$

in particular,  $f_a \in A_{\alpha_1}^{p_1}(D)$ . We can then apply the Toeplitz operator to  $f_a$  and consider the value of the resulting function for  $z = a$ :

$$(8) \quad \begin{aligned} T_\mu^\beta f_a(a) &= \int_D K_\beta(a, w) f_a(w) d\mu(w) = \int_D |K_\beta(a, w)|^2 d\mu(w) \\ &\geq \int_{B_D(a, r)} |K_\beta(a, w)|^2 d\mu(w) \geq \frac{\mu(B_D(a, r))}{\delta(a)^{2(n+\beta+1)}} \end{aligned}$$

as soon as  $a$  is close enough to  $\partial D$ , where, in the last inequality, we used Lemma 2.6.

Moreover, by Lemma 2.4

$$(9) \quad \begin{aligned} T_\mu^\beta f_a(a) &= [|T_\mu^\beta f_a(a)|^{p_2}]^{1/p_2} \leq \frac{1}{\nu(B_D(a, r))^{1/p_2}} \left[ \int_{B_D(a, r)} |T_\mu^\beta f_a(\zeta)|^{p_2} d\nu(\zeta) \right]^{1/p_2} \\ &\leq \frac{\delta(a)^{-\alpha_2/p_2}}{\nu(B_D(a, r))^{1/p_2}} \left[ \int_{B_D(a, r)} |T_\mu^\beta f_a(\zeta)|^{p_2} \delta(\zeta)^{\alpha_2} d\nu(\zeta) \right]^{1/p_2} \\ &\leq \delta(a)^{-(n+1+\alpha_2)/p_2} \|T_\mu^\beta f_a\|_{p_2, \alpha_2} \leq \|T_\mu^\beta\| \delta(a)^{-(n+1+\alpha_2)/p_2} \|f_a\|_{p_1, \alpha_1}, \end{aligned}$$

where we used Lemma 2.2 and Lemma 2.1.

Combining (7), (8) and (9) we conclude that

$$(10) \quad \begin{aligned} \mu(B_D(a, r)) &\leq \|T_\mu^\beta\| \delta(a)^{(n+1+\beta)+(n+1+\alpha_1)/p_1-(n+1+\alpha_2)/p_2} = \|T_\mu^\beta\| \delta(a)^{(n+1+\gamma)\lambda} \\ &\approx \|T_\mu^\beta\| \nu(B_D(a, r))^{(n+1+\gamma)\lambda/(n+1)}. \end{aligned}$$

This means that  $\mu$  is a geometric  $\lambda \left(1 + \frac{\gamma}{n+1}\right)$ -Carleson measure, which, by Theorem 2.12, is equivalent to  $\mu$  being a  $(\lambda, \gamma)$ -skew Carleson measure. Moreover,

$$\|\mu\|_{\lambda, \gamma} \leq \|T_\mu^\beta\|.$$

*Case 2.* Assume  $\lambda < 1$ , that is  $p_2 < p_1$ . In this case, we can adapt the proof of [5, Proposition 3.4] and we report here the complete proof for the sake of completeness.

Let  $\{a_k\}$  be an  $r$ -lattice in  $D$ , and  $\{r_k\}$  a sequence of Rademacher functions (see [10, Appendix A]). Set

$$\tau = \frac{n+1+\beta}{2} - \frac{n+1+\alpha_1}{p_1},$$

and, for every  $a \in D$ , put  $f_a = \delta(a)^\tau k_{\beta,a}$ . Then Lemma 2.8 implies that

$$f_t = \sum_{k=0}^{\infty} c_k r_k(t) f_{a_k}$$

belongs to  $A_{\alpha_1}^{p_1}(D)$  for all  $\mathbf{c} = \{c_k\} \in \ell^{p_1}$ , and  $\|f_t\|_{p_1, \alpha_1} \leq \|\mathbf{c}\|_{p_1}$ .

Since, by assumption,  $T_\mu^\beta$  is bounded from  $A_{\alpha_1}^{p_1}$  to  $A_{\alpha_2}^{p_2}$  we have

$$\begin{aligned} \|T_\mu^\beta f_t\|_{p_2, \alpha_2}^{p_2} &= \int_D \left| \sum_{k=0}^{\infty} c_k r_k(t) T_\mu^\beta f_{a_k}(z) \right|^{p_2} d\nu_{\alpha_2}(z) \\ &\leq \|T_\mu^\beta\|^{p_2} \|f_t\|_{p_1, \alpha_1}^{p_2} \leq \|T_\mu^\beta\|^{p_2} \|\mathbf{c}\|_{p_1}^{p_2}. \end{aligned}$$

Integrating both sides on  $[0, 1]$  with respect to  $t$  and using Khinchine's inequality (see, e.g., [24]) we obtain

$$\int_D \left( \sum_{k=0}^{\infty} |c_k|^2 |T_\mu^\beta f_{a_k}(z)|^2 \right)^{p_2/2} d\nu_{\alpha_2}(z) \leq \|T_\mu^\beta\|^{p_2} \|\mathbf{c}\|_{p_1}^{p_2}.$$

Set  $B_k = B_D(a_k, r)$ . We consider two cases:  $p_2 \geq 2$  and  $0 < p_2 < 2$ .

If  $p_2 \geq 2$ , using the fact that  $\|\mathbf{a}\|_{p_2/2} \leq \|\mathbf{a}\|_1$  for every  $\mathbf{a} \in \ell^1$  we get

$$\begin{aligned} \sum_{k=0}^{\infty} |c_k|^{p_2} \int_{B_k} |T_\mu^\beta f_{a_k}(z)|^{p_2} d\nu_{\alpha_2}(z) \\ \leq \int_D \left( \sum_{k=0}^{\infty} |c_k|^2 |T_\mu^\beta f_{a_k}(z)|^2 \chi_{B_k}(z) \right)^{p_2/2} d\nu_{\alpha_2}(z) \\ \leq \int_D \left( \sum_{k=0}^{\infty} |c_k|^2 |T_\mu^\beta f_{a_k}(z)|^2 \right)^{p_2/2} d\nu_{\alpha_2}(z). \end{aligned}$$

If instead  $0 < p_2 < 2$ , using Hölder's inequality, we obtain

$$\begin{aligned} \sum_{k=0}^{\infty} |c_k|^{p_2} \int_{B_k} |T_\mu^\beta f_{a_k}(z)|^{p_2} d\nu_{\alpha_2}(z) \\ \leq \int_D \left( \sum_{k=0}^{\infty} |c_k|^2 |T_\mu^\beta f_{a_k}(z)|^2 \right)^{\frac{p_2}{2}} \left( \sum_{k=0}^{\infty} \chi_{B_k}(z) \right)^{1-\frac{p_2}{2}} d\nu_{\alpha_2}(z) \\ \leq \int_D \left( \sum_{k=0}^{\infty} |c_k|^2 |T_\mu^\beta f_{a_k}(z)|^2 \right)^{p_2/2} d\nu_{\alpha_2}(z), \end{aligned}$$

where we used the fact that each  $z \in D$  belongs to no more than  $m$  of the  $B_k$ .

Summing up, for any  $p_2 > 0$  we have

$$\sum_{k=0}^{\infty} |c_k|^{p_2} \int_{B_k} |T_\mu^\beta f_{a_k}(z)|^{p_2} d\nu_{\alpha_2}(z) \leq \|T_\mu^\beta\|^{p_2} \|\mathbf{c}\|_{p_1}^{p_2}.$$

Now Lemmas 2.2, 2.1 and 2.4 yield

$$|T_\mu^\beta f_{a_k}(a_k)|^{p_2} \preceq \delta(a_k)^{-(n+1+\alpha_2)} \int_{B_k} |T_\mu^\beta f_{a_k}(z)|^{p_2} d\nu_{\alpha_2}(z),$$

and so we have

$$\sum_{k=0}^{\infty} |c_k|^{p_2} \delta(a_k)^{n+1+\alpha_2} |T_\mu^\beta f_{a_k}(a_k)|^{p_2} \preceq \|T_\mu^\beta\|^{p_2} \|\mathbf{c}\|_{p_1}^{p_2}.$$

On the other hand, using Lemmas 2.5 and 2.6, we obtain

$$\begin{aligned} T_\mu^\beta f_{a_k}(a_k) &= \delta(a_k)^\tau \int_D K_\beta(a_k, w) k_{\beta, a_k}(w) d\mu(w) \\ &\succeq \delta(a_k)^{\tau + \frac{n+1+\beta}{2}} \int_D |K_\beta(a_k, w)|^2 d\mu(w) \\ &\geq \delta(a_k)^{n+1+\beta - \frac{n+1+\alpha_1}{p_1}} \int_{B_D(a_k, r)} |K_\beta(a_k, w)|^2 d\mu(w) \\ &\succeq \delta(a_k)^{n+1+\beta - \frac{n+1+\alpha_1}{p_1}} \frac{\mu(B_D(a_k, r))}{\delta(a_k)^{2(n+1+\beta)}} = \frac{\mu(B_D(a_k, r))}{\delta(a_k)^{n+1+\beta + \frac{n+1+\alpha_1}{p_1}}}. \end{aligned}$$

Putting all together we get

$$\sum_{k=0}^{\infty} |c_k|^{p_2} \left( \frac{\mu(B_D(a_k, r))}{\delta(a_k)^{(n+1+\gamma)\lambda}} \right)^{p_2} \preceq \|T_\mu^\beta\|^{p_2} \|\mathbf{c}\|_{p_1}^{p_2},$$

since

$$n+1+\beta + \frac{n+1+\alpha_1}{p_1} - \frac{n+1+\alpha_2}{p_2} = (n+1+\gamma)\lambda.$$

Now, set  $\mathbf{d} = \{d_k\}$ , where

$$d_k = \frac{\mu(B_D(a_k, r))}{\delta(a_k)^{(n+1+\gamma)\lambda}}.$$

Then by duality we get  $\{d_k^{p_2}\} \in \ell^{p_1/(p_1-p_2)}$  with  $\|\{d_k^{p_2}\}\|_{p_1/(p_1-p_2)} \preceq \|T_\mu^\beta\|^{p_2}$ , because  $p_1/(p_1-p_2)$  is the conjugate exponent of  $p_1/p_2 > 1$ . This means that  $\mathbf{d} \in \ell^{p_1 p_2/(p_1-p_2)} = \ell^{1/(1-\lambda)}$  with

$$\|\mathbf{d}\|_{\frac{1}{1-\lambda}} \preceq \|T_\mu^\beta\|,$$

and the assertion then follows from Theorem 2.13 (notice that the proof in [16] that  $\{\hat{\mu}_{r, \lambda \theta}(a_k)\} \in \ell^{\frac{1}{1-\lambda}}$  implies  $\hat{\mu}_r \delta^{-\lambda \gamma} \in L^{\frac{1}{1-\lambda}}(D)$ , where  $\theta = 1 + \frac{\gamma}{n+1}$ , holds also for  $\lambda \leq 0$ ).

(ii)  $\Rightarrow$  (i) We consider three cases:  $p_2 > 1$ ,  $p_2 = 1$  and  $0 < p_2 < 1$ .

Case 1. If  $p_2 > 1$ , let  $p'_2 > 1$  be the conjugate exponent of  $p_2$ , and choose  $\alpha'_2 \in \mathbb{R}$  so that

$$(11) \quad \beta = \frac{\alpha_2}{p_2} + \frac{\alpha'_2}{p'_2}.$$

An easy computation shows that  $\alpha'_2 = \alpha_2 + (\beta - \alpha_2)p'_2$ , and then  $\alpha'_2 > -1$  follows from (6) for  $j = 2$ .

Take  $f \in A_{\alpha_1}^{p_1}(D)$  and  $h \in A_{\alpha'_2}^{p'_2}(D)$ . Then

$$(12) \quad \begin{aligned} (T_\mu^\beta f, h)_\beta &= \int_D \overline{h(z)} \int_D K_\beta(z, w) f(w) d\mu(w) d\nu_\beta(z) \\ &= \int_D \int_D \overline{K_\beta(w, z) h(z)} f(w) d\nu_\beta(z) d\mu(w) = \int_D \overline{h(w)} f(w) d\mu(w). \end{aligned}$$

Therefore, as  $\mu$  is  $(\lambda, \gamma)$ -skew Carleson, by Theorem 2.17, we have

$$|(T_\mu^\beta f, h)_\beta| \preceq \|\mu\|_{\lambda, \gamma} \|f\|_{p_1, \alpha_1} \|h\|_{p_2', \alpha_2'}$$

because, by our hypotheses,

$$\lambda = \frac{1}{p_1} + \frac{1}{p_2'} \quad \text{and} \quad \gamma = \frac{1}{\lambda} \left( \frac{\alpha_1}{p_1} + \frac{\alpha_2'}{p_2'} \right).$$

As this holds for every  $h \in A_{\alpha_2'}^{p_2'}(D)$ , that is, by Lemma 3.1, for every continuous functional on  $A_{\alpha_2}^{p_2}(D)$ , we conclude that

$$\|T_\mu^\beta f\|_{p_2, \alpha_2} \preceq \|\mu\|_{\lambda, \gamma} \|f\|_{p_1, \alpha_1},$$

that is  $T_\mu^\beta$  is bounded from  $A_{\alpha_1}^{p_1}(D)$  to  $A_{\alpha_2}^{p_2}(D)$  and  $\|T_\mu^\beta\| \preceq \|\mu\|_{\lambda, \gamma}$ .

*Case 2.* If  $p_2 = 1$ , that is  $\lambda = \frac{1}{p_1}$ , condition (6) for  $j = 2$  implies  $\beta - \alpha_2 > 0$ . Take  $f \in A_{\alpha_1}^{p_1}(D)$ . Then

$$\begin{aligned} \|T_\mu^\beta f\|_{1, \alpha_2} &\leq \int_D \int_D |K_\beta(z, w)| |f(w)| d\mu(w) d\nu_{\alpha_2}(z) \\ &= \int_D |f(w)| \int_D |K_\beta(z, w)| \delta(z)^{\alpha_2 - \beta} d\nu_\beta(z) d\mu(w) \preceq \int_D |f(w)| \delta(w)^{\alpha_2 - \beta} d\mu(w) \end{aligned}$$

by Theorem 2.7.

Now, as  $\mu$  is  $(\lambda, \gamma)$ -skew Carleson, Lemma 2.16 implies that  $\delta^{\alpha_2 - \beta} \mu$  is  $\left(\frac{1}{p_1}, \alpha_1\right)$ -skew Carleson, with  $\|\delta^{\alpha_2 - \beta} \mu\|_{1/p_1, \alpha_1} \approx \|\mu\|_{\lambda, \gamma}$ . Theorems 2.12 and 2.13 then implies that  $\delta^{\alpha_2 - \beta} \mu$  is  $(p_1, 1; \alpha_1)$ -skew Carleson, and so we obtain

$$\|T_\mu^\beta f\|_{1, \alpha_2} \preceq \|\mu\|_{\lambda, \gamma} \|f\|_{p_1, \alpha_1},$$

as desired.

*Case 3.* If  $0 < p_2 < 1$ , thanks to Lemma 2.3 we can find a  $r$ -lattice  $\{a_k\}$  and  $m \in \mathbb{N}$  such that for every  $z \in D$  there exist at most  $m$  values of  $k$  such that  $z \in B_D(a_k, R)$ , where  $R = \frac{1}{2}(1+r)$ . Put  $B_k = B_D(a_k, r)$  and  $\tilde{B}_k = B_D(a_k, R)$ .

By Lemmas 2.2, 2.1 and 2.4 for  $w \in B_k$  we have

$$|f(w)|^{p_1} \preceq \frac{1}{\nu_{\alpha_1}(B_k)} \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta)$$

and

$$|K_\beta(z, w)|^{p_2} \preceq \frac{1}{\nu_{\alpha_2}(B_k)} \int_{\tilde{B}_k} |K_\beta(z, \zeta)|^{p_2} d\nu_{\alpha_2}(\zeta).$$

Therefore, integrating on  $B_k$  we get

$$\begin{aligned} \int_{B_k} |K_\beta(z, w)| |f(w)| d\mu(w) \\ \preceq \frac{\mu(B_k)}{\nu_{\alpha_1}(B_k)^{1/p_1} \nu_{\alpha_2}(B_k)^{1/p_2}} \left( \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{1/p_1} \left( \int_{\tilde{B}_k} |K_\beta(z, \zeta)|^{p_2} d\nu_{\alpha_2}(\zeta) \right)^{1/p_2}. \end{aligned}$$

Since  $p_2 < 1$ , summing over  $k$  we get

$$|T_\mu^\beta f(z)|^{p_2} \preceq \sum_{k=1}^{\infty} \frac{\mu(B_k)^{p_2}}{\nu_{\alpha_1}(B_k)^{p_2/p_1} \nu_{\alpha_2}(B_k)} \left( \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} \int_{\tilde{B}_k} |K_\beta(z, \zeta)|^{p_2} d\nu_{\alpha_2}(\zeta).$$

Integrating in  $z$  over  $D$  with respect to  $\nu_{\alpha_2}$  we obtain

$$(13) \quad \|T_\mu^\beta f\|_{p_2, \alpha_2}^{p_2} \preceq \sum_{k=1}^{\infty} \frac{\mu(B_k)^{p_2}}{\delta(a_k)^{(n+1+\gamma)\lambda p_2}} \left( \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}},$$

thanks to Lemma 2.2, Lemma 2.1 and Theorem 2.7, that we can apply because of (6) for  $j = 2$ .

Now, if  $\lambda \geq 1$  we have that

$$\mu(B_k) \preceq \|\mu\|_{\lambda, \gamma} \delta(a_k)^{(n+1+\gamma)\lambda},$$

and so (13) yields

$$\|T_\mu^\beta f\|_{p_2, \alpha_2}^{p_2} \preceq \|\mu\|_{\lambda, \gamma}^{p_2} \sum_{k=1}^{\infty} \left( \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} \preceq \|\mu\|_{\lambda, \gamma}^{p_2} \|f\|_{p_1, \alpha_1}^{p_2}.$$

On the other hand, if  $\lambda < 1$  (that is  $p_1/p_2 > 1$ ), by Hölder inequality we have

$$\begin{aligned} & \sum_{k=1}^{\infty} \frac{\mu(B_k)^{p_2}}{\delta(a_k)^{(n+1+\gamma)\lambda p_2}} \left( \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} \\ & \leq \left( \sum_{k=1}^{\infty} \left( \frac{\mu(B_k)}{\delta(a_k)^{(n+1+\gamma)\lambda}} \right)^{\frac{p_1 p_2}{p_1 - p_2}} \right)^{\frac{p_1 - p_2}{p_1}} \left( \sum_{k=1}^{\infty} \int_{\tilde{B}_k} |f(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}}. \end{aligned}$$

Now, the proof of the implication (b) $\Rightarrow$ (c) in [16, Lemma 2.5] applied with  $s = -\gamma\lambda$  and  $p = p_1 p_2 / (p_1 - p_2)$  yields

$$\left\{ \frac{\mu(B_k)}{\delta(a_k)^{(n+1+\gamma)\lambda}} \right\}_{k \geq 1} \in \ell^{\frac{p_1 p_2}{p_1 - p_2}}$$

and

$$\left\| \left\{ \frac{\mu(B_k)}{\delta(a_k)^{(n+1+\gamma)\lambda}} \right\}_{k \geq 1} \right\|_{\ell^{\frac{p_1 p_2}{p_1 - p_2}}} \approx \|\mu\|_{\lambda, \gamma}.$$

So

$$\|T_\mu^\beta f\|_{p_2, \alpha_2}^{p_2} \preceq \|\mu\|_{\lambda, \gamma}^{p_2} \|f\|_{p_1, \alpha_1}^{p_2},$$

and we are done in this case too.  $\square$

#### 4. COMPACT TOEPLITZ OPERATORS AND VANISHING SKEW CARLESON MEASURES

In this section we shall prove a version of Theorem 3.2 concerning compact Toeplitz operators and vanishing skew-Carleson measures. The only interesting case is  $\lambda \geq 1$ , because for  $\lambda < 1$  (that is  $p_2 < p_1$ ) all  $(\lambda, \gamma)$ -skew Carleson measures are vanishing (Theorem 2.13) and all continuous operators from  $A_{\alpha_1}^{p_1}(D)$  to  $A_{\alpha_2}^{p_2}(D)$  are compact (see, e.g., [23, Proposition 2.c.3]).

To deal with the case  $\lambda \geq 1$  we shall need the following version of Theorem 2.17, whose proof is analogous to the proof of [27, Theorem 4.1]:

**Theorem 4.1.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain, and let  $\mu$  be a positive finite Borel measure on  $D$ . Fix an integer  $k \geq 1$ , and let  $0 < p_j, q_j < +\infty$  and  $-1 < \alpha_j < +\infty$  be given for  $j = 1, \dots, k$ . Set*

$$\lambda = \sum_{j=1}^k \frac{q_j}{p_j} \quad \text{and} \quad \gamma = \frac{1}{\lambda} \sum_{j=1}^k \frac{\alpha_j q_j}{p_j}.$$

*Assume that  $\lambda \geq 1$ . Then the following statements are equivalent:*

- (i)  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure.



- (ii) For any sequence  $\{f_{1,\ell}\}_\ell$  in the unit ball of  $A_{\alpha_1}^{p_1}(D)$  converging to 0 uniformly on compact sets in  $D$  we have

$$\lim_{\ell \rightarrow \infty} F(\ell) = 0 ,$$

where

$$F(\ell) = \sup \left\{ \int_D |f_{1,\ell}(z)|^{q_1} \prod_{j=2}^k |f_j(z)|^{q_j} d\mu(z) \mid \|f_j\|_{p_j, \alpha_j} \leq 1, j = 2, \dots, k \right\} .$$

- (iii) For any  $k$  sequences  $\{f_{1,\ell}\}, \dots, \{f_{k,\ell}\}$  in the unit balls of  $A_{\alpha_1}^{p_1}(D), \dots, A_{\alpha_k}^{p_k}(D)$ , respectively, which are all convergent to 0 uniformly on compact sets in  $D$ , we have

$$\lim_{\ell \rightarrow \infty} \int_D |f_{1,\ell}(z)|^{q_1} \cdots |f_{k,\ell}(z)|^{q_k} d\mu(z) = 0 .$$

*Proof.* Assume (i) is satisfied, that is  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure. Let  $\{f_{1,\ell}\}_{\ell \in \mathbb{N}}$  be a sequence in the unit ball of  $A_{\alpha_1}^{p_1}(D)$  which converges to 0 uniformly on compact subsets of  $D$ , and for  $j = 2, \dots, k$  let  $f_j$  be an arbitrary function in the unit ball of  $A_{\alpha_j}^{p_j}(D)$ . Given  $r > 0$ , let us set  $D_r = \{z \in D \mid \delta(z) < r\}$ . Then  $\mu_r = \mu|_{D_r}$  is a  $(\lambda, \gamma)$ -skew Carleson measure, and

$$\lim_{r \rightarrow 0} \|\mu_r\|_{\lambda, \gamma} = 0$$

because  $\mu$  is vanishing. Fix  $\varepsilon > 0$ . Then if  $r > 0$  is small enough Theorem 2.17 yields

$$(14) \quad \int_{D_r} |f_{1,\ell}(z)|^{q_1} |f_2(z)|^{q_2} \cdots |f_k(z)|^{q_k} d\mu(z) = \int_D |f_{1,\ell}(z)|^{q_1} |f_2(z)|^{q_2} \cdots |f_k(z)|^{q_k} d\mu_r(z) \leq \varepsilon .$$

On the other hand, thanks to the uniform convergence of  $f_{1,\ell}$  to 0 on compact subsets of  $D$ , we can find  $M \in \mathbb{N}$  such that for any  $\ell > M$  we have  $|f_{1,\ell}(z)| < \varepsilon$  for all  $z \in D \setminus D_r$ . Therefore applying again Theorem 2.17 we have

$$(15) \quad \begin{aligned} \int_{D \setminus D_r} |f_{1,\ell}(z)|^{q_1} |f_2(z)|^{q_2} \cdots |f_k(z)|^{q_k} d\mu(z) &\leq \varepsilon \int_D |f_2(z)|^{q_2} \cdots |f_k(z)|^{q_k} d\mu(z) \\ &= \varepsilon \int_D |1|^{q_1} |f_2(z)|^{q_2} \cdots |f_k(z)|^{q_k} d\mu(z) \leq \varepsilon . \end{aligned}$$

These last two estimates together imply (ii).

It is evident that (ii) implies (iii). To prove that (iii) implies (i) we follow the same construction as in the proof of Theorem 2.17. Choose  $\sigma_1, \dots, \sigma_k \in \mathbb{N}^*$  such that

$$p_j \sigma_j > \max \left\{ 1, 1 + \frac{\alpha_j}{n+1} \right\}$$

for all  $j = 1, \dots, k$ , and

$$\sum_{j=1}^k q_j \sigma_j > \lambda \gamma ,$$

and set

$$r_j = \frac{(n+1)\sigma_j}{2} - \frac{n+1+\alpha_j}{p_j} .$$

For any  $a \in D$  and  $j = 1, \dots, k$ , consider

$$f_{j,a}(z) = \delta(a)^{r_j} k_a(z)^{\sigma_j} .$$

Then, since  $\alpha_j < (n+1)(p_j\sigma_j - 1)$  by the choice of  $\sigma_j$  we know (Theorem 2.7) that  $\|f_{j,a}\|_{p_j,\alpha_j} \leq 1$  for all  $j = 1, \dots, k$ ; moreover it is easy to see that

$$\lim_{a \rightarrow \partial D} |f_{j,a}(z)| = 0$$

uniformly on any compact subset of  $D$ . Therefore (iii) yields

$$(16) \quad \lim_{a \rightarrow \partial D} \int_D \prod_{j=1}^k |f_{j,a}(z)|^{q_j} d\mu(z) = 0.$$

Now, we have

$$\int_D \prod_{j=1}^k |f_{j,a}(z)|^{q_j} d\mu(z) = \int_D |k_a(z)|^{\sum_j q_j \sigma_j} \delta(a)^{\sum_j q_j r_j} d\mu(z),$$

and

$$\sum_{j=1}^k q_j r_j = (n+1) \sum_{j=1}^k \left[ \frac{q_j \sigma_j}{2} - \theta_j \frac{q_j}{p_j} \right] = \frac{n+1}{2} \sum_{j=1}^k q_j \sigma_j - (n+1) \lambda \gamma.$$

Therefore, setting  $s = \sum_j \sigma_j q_j > \lambda \gamma$ , (16) becomes

$$\lim_{a \rightarrow \partial D} \delta(a)^{(n+1)(\frac{s}{2} - \lambda \gamma)} \int_D |k_a(z)|^s d\mu(z) = \lim_{a \rightarrow \partial D} \delta(a)^{(n+1)(\frac{s}{2} - \lambda \gamma)} B^s \mu(a) = 0,$$

where  $B^s \mu$  is the Berezin transform of level  $s$  of  $\mu$ , and so  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure thanks to Theorem 2.14.  $\square$

We can now prove the following result:

**Theorem 4.2.** *Let  $D \Subset \mathbb{C}^n$  be a bounded strongly pseudoconvex domain. Let  $0 < p_1 \leq p_2 < +\infty$  and  $-1 < \alpha_1, \alpha_2 < +\infty$ . Suppose that  $\beta \in \mathbb{R}$  satisfies*

$$(17) \quad n + 1 + \beta > n \max \left\{ 1, \frac{1}{p_j} \right\} + \frac{1 + \alpha_j}{p_j}$$

for  $j = 1, 2$ . Put

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}$$

and

$$\gamma = \frac{1}{\lambda} \left( \beta + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then for any positive Borel measure  $\mu$  on  $D$  the following statements are equivalent:

- (i)  $T_\mu^\beta : A_{\alpha_1}^{p_1}(D) \rightarrow A_{\alpha_2}^{p_2}(D)$  compactly;
- (ii)  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure.

*Proof.* Assume that (i) holds. Since  $T_\mu^\beta$  is compact, it maps every bounded sequence in  $A_{\alpha_1}^{p_1}(D)$  converging uniformly to 0 on compact subsets of  $D$  to a sequence strongly converging to 0 in  $A_{\alpha_2}^{p_2}(D)$ .

We consider a sequence  $\{a_k\} \in D$  such that  $\lim_{k \rightarrow +\infty} \delta(a_k) = 0$  and we set

$$f_k(z) = \delta(a_k)^{(n+1+\beta) - (n+1+\alpha_1)/p_1} K_\beta(z, a_k).$$

Thanks to Theorem 2.7, we have that

$$\|f_k\|_{p_1, \alpha_1}^{p_1} \leq 1.$$

Moreover, for any  $L \Subset D$  there exists a constant  $C_1 > 0$  such that  $|K_\beta|$  is bounded from above by  $C_1$  on  $L \times D$ . Therefore for every  $z \in L$  we have that

$$|f_k(z)| \leq C_1 \delta(a_k)^{n+1+\beta-(n+1+\alpha_1)/p_1}$$

and so, since since our hypotheses give us that  $(n+1+\beta) - (n+1+\alpha_1)/p_1 > 0$ , we get

$$\lim_{k \rightarrow +\infty} \sup_{z \in L} |f_k(z)| \leq \lim_{k \rightarrow +\infty} C_1 \delta(a_k)^{n+1+\beta-(n+1+\alpha_1)/p_1} = 0.$$

Hence the compactness of  $T_\mu^\beta$  implies

$$(18) \quad \lim_{k \rightarrow +\infty} \|T_\mu^\beta f_k\|_{p_2, \alpha_2} = 0.$$

Now, the same computations as in the proof of the implication (i)  $\implies$  (ii) of Theorem 3.2 yield

$$\frac{\mu(B_D(a_k, r))}{\delta(a_k)^{(n+1+\beta)+(n+1+\alpha_1)/p_1}} \preceq T_\mu^\beta f_k(a_k)$$

and

$$T_\mu^\beta f_k(a_k) \preceq \delta(a_k)^{-(n+1+\alpha_2)/p_2} \|T_\mu^\beta f_k\|_{p_2, \alpha_2}.$$

Therefore

$$\frac{\mu(B_D(a_k, r))}{\delta(a_k)^{(n+1+\gamma)\lambda}} \preceq \|T_\mu^\beta f_k\|_{p_2, \alpha_2},$$

which, together with (18) and Theorem 2.14, implies that  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure.

Conversely, assume that  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure with  $\lambda \geq 1$ , and let  $\{g_k\}_{k \in \mathbb{N}}$  be a bounded sequence in  $A_{\alpha_1}^{p_1}(D)$  converging uniformly to 0 on compact subsets of  $D$ . We want to prove that the bounded sequence  $\{T_\mu^\beta g_k\}_{k \in \mathbb{N}} \subset A_{\alpha_2}^{p_2}(D)$  converges strongly to 0 in  $A_{\alpha_2}^{p_2}(D)$ . We consider two cases:  $p_2 > 1$  and  $0 < p_2 \leq 1$ .

If  $p_2 > 1$  then, as in the proof of Theorem 3.2, thanks to Lemma 3.1, denoting by  $p_2'$  the conjugate exponent of  $p_2$  and  $\alpha_2'$  the number defined in (11), using (12) we have

$$\|T_\mu^\beta g_k\|_{p_2, \alpha_2} \approx \sup_{\|h\|_{p_2', \alpha_2'} \leq 1} |\langle h, T_\mu^\beta g_k \rangle_\beta| \leq \sup_{\|h\|_{p_2', \alpha_2'} \leq 1} \int_D |h(z)| |g_k(z)| d\mu(z),$$

and Theorem 4.1 yields that the last integral converges to 0 as  $k$  tends to  $+\infty$ .

If  $0 < p_2 \leq 1$ , for any  $r$ -lattice  $\{a_j\}$  we consider the associated balls  $\{B_j = B_D(a_j, r)\}$  and  $\{\tilde{B}_j = B_D(a_j, R)\}$ , where  $R = (1+r)/2$ , as usual. Using (13) we obtain that

$$(19) \quad \|T_\mu^\beta g_k\|_{p_2, \alpha_2}^{p_2} \preceq \sum_{j=1}^{\infty} \frac{\mu(B_j)^{p_2}}{\delta(a_j)^{(n+1+\gamma)\lambda p_2}} \left( \int_{\tilde{B}_j} |g_k(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{p_2/p_1}.$$

Let  $\varepsilon > 0$ . Since  $\mu$  is a vanishing  $(\lambda, \gamma)$ -skew Carleson measure by Theorem 2.14 there exists  $j_0 > 0$  such that

$$\frac{\mu(B_j)}{\delta(a_j)^{(n+1+\gamma)\lambda}} < \varepsilon$$

for all  $j > j_0$ . Choose  $\delta_0 > 0$  such that  $\tilde{B}_j \subset L = \{z \in D \mid \delta(z) \geq \delta_0\} \Subset D$  for all  $j \leq j_0$ . We can then split the sum in the right-hand-side of (19) into two parts. For the first part we have

$$\sum_{j=1}^{j_0} \frac{\mu(B_j)^{p_2}}{\delta(a_j)^{(n+1+\gamma)\lambda p_2}} \left( \int_{\tilde{B}_j} |g_k(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} \preceq \left( \sup_L |g_k|^{p_1} \right)^{\frac{p_2}{p_1}} \sum_{j=1}^{j_0} \frac{\mu(B_j)^{p_2} \nu_{\alpha_1}(B_j)^{\frac{p_2}{p_1}}}{\delta(a_j)^{(n+1+\gamma)\lambda p_2}}$$

and clearly the right-hand-side converges to 0 as  $k$  tends to  $+\infty$ .

On the other hand we have

$$\begin{aligned} \sum_{j=j_0+1}^{\infty} \frac{\mu(B_j)^{p_2}}{\delta(a_j)^{(n+1+\gamma)\lambda p_2}} \left( \int_{\tilde{B}_j} |g_k(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} &< \varepsilon^{p_2} \sum_{j=j_0+1}^{\infty} \left( \int_{\tilde{B}_j} |g_k(\zeta)|^{p_1} d\nu_{\alpha_1}(\zeta) \right)^{\frac{p_2}{p_1}} \\ &\preceq \varepsilon^{p_2} \|g_k\|_{p_1, \alpha_1}^{p_2} \preceq \varepsilon^{p_2} \end{aligned}$$

because the sequence  $\{g_k\}$  is norm-bounded. Therefore  $\lim_{k \rightarrow +\infty} \|T_{\mu}^{\beta} g_k\|_{p_2, \alpha_2} = 0$ , and this concludes the proof.  $\square$

## REFERENCES

- [1] M. Abate, *Iteration theory of holomorphic maps on taut manifolds*, Research and Lecture Notes in Mathematics. Complex Analysis and Geometry, Mediterranean Press, Rende, 1989.
- [2] ———, *Angular derivatives in several complex variables*, Real methods in complex and CR geometry, 2004, pp. 1–47.
- [3] M. Abate and A. Saracco, *Carleson measures and uniformly discrete sequences in strongly pseudoconvex domains*, J. Lond. Math. Soc. (2) **83** (2011), no. 3, 587–605, DOI 10.1112/jlms/jdq092.
- [4] M. Abate, J. Raissy, and A. Saracco, *Toeplitz operators and Carleson measures in strongly pseudoconvex domains*, J. Funct. Anal. **263** (2012), no. 11, 3449–3491, DOI 10.1016/j.jfa.2012.08.027.
- [5] M. Abate and J. Raissy, *Skew Carleson measures in strongly pseudoconvex domains*, Complex Anal. Oper. Theory **13** (2019), no. 2, 405–429, DOI 10.1007/s11785-018-0823-4.
- [6] L. Carleson, *Interpolations by bounded analytic functions and the corona problem*, Ann. of Math. **76** (1962), 547–559.
- [7] J.A. Cima and P.R. Mercer, *Composition operators between Bergman spaces on convex domains in  $\mathbb{C}^n$* , J. Operator Theory **33** (1995), 363–369.
- [8] J.A. Cima and W.R. Wogen, *A Carleson measure theorem for the Bergman space on the ball*, J. Operator Theory **7** (1982), 157–165.
- [9] Ž. Čučković and J.D. McNeal, *Special Toeplitz operators on strongly pseudoconvex domains*, Rev. Mat. Iberoam. **22** (2006), 851–866.
- [10] P.L. Duren, *Extension of a theorem of Carleson*, Bull. Amer. Math. Soc. **75** (1969), 143–146, DOI 10.1090/S0002-9904-1969-12181-6.
- [11] P.L. Duren and R. Weir, *The pseudohyperbolic metric and Bergman spaces in the ball*, Trans. Amer. Math. Soc. **359** (2007), 63–76.
- [12] M. Engliš, *Toeplitz operators and weighted Bergman kernels*, J. Funct. Anal. **255** (2008), no. 6, 1419–1457, DOI 10.1016/j.jfa.2008.06.026.
- [13] W.W. Hastings, *A Carleson measure theorem for Bergman spaces*, Proc. Amer. Math. Soc. **52** (1975), 237–241.
- [14] L. Hörmander,  *$L^2$  estimates and existence theorems for the  $\bar{\partial}$  operator*, Acta Math. **113** (1965), 89–152, DOI 10.1007/BF02391775.
- [15] ———, *An introduction to complex analysis in several variables*, Third, North-Holland Mathematical Library, vol. 7, North-Holland Publishing Co., Amsterdam, 1990.
- [16] Z. Hu, X. Lv, and K. Zhu, *Carleson measures and balayage for Bergman spaces of strongly pseudoconvex domains*, Math. Nachr. **289** (2016), no. 10, 1237–1254. MR3520714
- [17] M. Jarnicki and P. Pflug, *Invariant distances and metrics in complex analysis*, extended, De Gruyter Expositions in Mathematics, vol. 9, Walter de Gruyter GmbH & Co. KG, Berlin, 2013.
- [18] H.T. Kaptanoğlu, *Carleson measures for Besov spaces on the ball with applications*, J. Funct. Anal. **250** (2007), 483–520.
- [19] S. Kobayashi, *Hyperbolic complex spaces*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 318, Springer-Verlag, Berlin, 1998.
- [20] S.G. Krantz, *Function theory of several complex variables*, Second, The Wadsworth & Brooks/Cole Mathematics Series, Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 1992.
- [21] H. Li, *BMO, VMO and Hankel operators on the Bergman space of strictly pseudoconvex domains*, J. Funct. Anal. **106** (1992), 375–408.
- [22] H. Li and D.H. Luecking, *Schatten class of Hankel and Toeplitz operators on the Bergman space of strongly pseudoconvex domains*, Multivariable operator theory (Seattle, WA, 1993), 1995, pp. 237–257.

- [23] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. I and II*, Springer-Verlag, Berlin, 1996.
- [24] D. Luecking, *A technique for characterizing Carleson measures on Bergman spaces*, Proc. Amer. Math. Soc. **87** (1983), no. 4, 656–660, DOI 10.2307/2043353.
- [25] V.L. Oleinik, *Embeddings theorems for weighted classes of harmonic and analytic functions*, J. Soviet Math. **9** (1978), 228–243.
- [26] V.L. Oleinik and B.S. Pavlov, *Embedding theorems for weighted classes of harmonic and analytic functions*, J. Soviet Math. **2** (1974), 135–142.
- [27] J. Pau and R. Zhao, *Carleson measures and Toeplitz operators for weighted Bergman spaces on the unit ball*, Michigan Math. J. **64** (2015), 759–796.
- [28] R.M. Range, *Holomorphic functions and integral representations in several complex variables*, Graduate Texts in Mathematics, vol. 108, Springer-Verlag, New York, 1986.
- [29] A.P. Schuster and D. Varolin, *Toeplitz operators and Carleson measures on generalized Bargmann-Fock spaces*, Integral Equations Operator Theory **72** (2012), 363–392.
- [30] K. Zhu, *Positive Toeplitz operators on weighted Bergman spaces of bounded symmetric domains*, J. Operator Theory **20** (1988), 329–357.
- [31] ———, *Spaces of holomorphic functions in the unit ball*, Graduate Texts in Mathematics, vol. 226, Springer-Verlag, New York, 2005.

MARCO ABATE, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO PONTECORVO 5, I-56127 PISA, ITALY.  
*E-mail address:* `marco.abate@unipi.it`

SAMUELE MONGODI, POLITECNICO DI MILANO, DIPARTIMENTO DI MATEMATICA, VIA BONARDI, 9 I-20133 MILANO, ITALY  
*E-mail address:* `samuele.mongodi@polimi.it`

JASMIN RAISSY, INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR5219, UNIVERSITÉ DE TOULOUSE, CNRS, UPS, F-31062 TOULOUSE CEDEX 9, FRANCE  
*E-mail address:* `jraissy@math.univ-toulouse.fr`