

# **A reexamination of the QBO-period modulation by the solar cycle**

P. Fischer

Université Bordeaux 1, IMB, INRIA Team MC2, Talence, France.

K.K. Tung

Department of Applied Mathematics, University of Washington, Seattle,

Washington, USA.

---

P. Fischer, Université Bordeaux 1, IMB, INRIA Team MC2, 351 Cours de la Liberation, 33405  
Talence, France. (Patrick.Fischer@math.u-bordeaux1.fr)

K.K. Tung, Department of Applied Mathematics, Box 352420, University of Washington, Seat-  
tle WA 98105. (tung@amath.washington.edu)

Using the updated Singapore wind from 1953-2007 for the lower stratosphere 70hPa-10hPa, courtesy of Barbara Naujokat of Free University of Berlin, we examine the variation of the period of the Quasi-Biennial Oscillation (QBO) as a function of height and its modulation in time by the 11-year solar cycle. The analysis is supplemented by the ERA-40 reanalysis up to 1 hPa. Previously it was reported that the descent of the easterly shear zone tends to stall near 30 hPa during solar minimum, leading to a lengthened QBO westerly duration near 44-50 hPa, and the reported anti-correlation of the westerly duration and the solar cycle. Using an objective method, Continuous Wavelet Transform (CWT), for the determination of local QBO period, we find that the whole QBO period is almost invariant with respect to height, so that the stalling mechanism affects only the partition of the whole period between easterly and westerly durations. Using this longest dataset available for equatorial stratospheric wind, which spans five and half solar cycles (six solar minima), we find that in three solar minima, the QBO period is lengthened, while in the remaining almost three solar cycles, the QBO period is lengthened instead at solar maxima. We suggest that the decadal variation of the QBO period originates in the upper stratosphere, where the solar-ozone radiative influence is strong. The solar modulation of the QBO period is found to be non-stationary; the averaged effect cannot be determined unless the data record is much longer. In shorter records, the correlation can change

<sup>27</sup> sign, as we have found in segments of the longest record available, with or  
<sup>28</sup> without lag.

## 1. Introduction

The Quasi-Biennial Oscillation (QBO) is a dominant oscillation of the equatorial stratospheric zonal wind, whose period is irregular but averages to about 28 months. The classical mechanism of QBO [Holton and Lindzen, 1972; Lindzen and Holton, 1968] attributes the period of the QBO to internal interactions between the waves and the mean flow in the equatorial stratosphere. Later modifications to the theory take into account the QBO's secondary circulation [Plumb and Bell, 1982] and the upwelling branch at the equator of the Brewer-Dobson circulation in affecting the descent rate and hence the period [Baldwin *et al.*, 2001; Kinnersley and Pawson, 1996].

Whether the QBO's period is affected by external forcing, such as the 11-year variation in the solar radiation (especially its variation in the UV component), is an intriguing open question. Quiroz [1981], using 12-month running mean of the Balboa data (9.0N, 79.6W), was the first to point out that there is a decadal variation in the QBO's period. Salby and Callaghan [2000] found, using radiosonde data near the equator at 45 hPa from 1956-1996 (from Free University of Berlin), that the duration of the westerly phase varies on a decadal cycle from 12 months to 23 months, being longer during solar minimum and shorter during solar maximum. The easterly phase at that level, on the other hand, seems to be always about 12 months long in duration. The authors suggested that easterlies near 30 hPa tend to stall during solar min instead of descending and replacing the westerlies below, thus prolonging the westerly phase near 45 hPa.

The descent of the easterlies tends to stall more easily than that of the westerlies because the QBO's self-induced secondary circulation is upward for easterly momentum

acceleration [*Plumb and Bell*, 1982]. Slowing down the descent of both the easterly and westerly phases near the equator is the upward branch of the Brewer-Dobson circulation, which however is stronger during the easterly phase of the QBO. The Brewer-Dobson circulation is remotely forced by planetary-wave breaking and dissipation in the polar stratosphere [*Holton et al*, 1995; *Randel et al.*, 2002; *Hood and Soukharev*, 2003]. This is consistent with the finding of *Dunkerton* [1990] that this stalling almost always occurs during Northern Hemisphere winter, when the planetary wave dissipation is the strongest. It has been long suggested [*Labitzke*, 1982] that more Stratospheric Sudden Warmings (SSWs) occur in late winter in Northern Hemisphere during solar max than during solar min. This result has recently been established statistically by *Camp and Tung* [2007]. Since SSWs produce downwelling at the pole and upwelling at the equator, it would seem more plausible that the stalling of the descent of the QBO in the equatorial region should occur during solar max instead of the solar min., at least in late winter, unless the proposed stalling mechanism is not the dominant factor in controlling the QBO period. *Kodera and Kuroda* [2002], on the other hand, presented data which showed that the upwelling branch in the subtropics of the Brewer-Dobson circulation is weakened during solar max. No data is shown equatorward of 20 degrees of latitude and below 10 hPa, a region of interest for the equatorial QBO. Furthermore, their composite uses only two solar cycles, from 1979-1998, during which two volcano eruptions occurred (El Chichon and Pinatubo).

*Soukharev and Hood* [2001] confirmed the conclusion of *Salby and Callaghan* [2000] using a composite analysis of the band-pass filtered 10 to 70hPa equatorial zonal wind for the

period from January 1957 to December 1999 (from Free University of Berlin). For each solar maximum (and for each solar minimum), two westerly and two easterly phases of the equatorial zonal wind were composited after alignment. The eight westerly or easterly phases of the zonal wind were aligned in such a way that the zero-wind lines all start at 10hPa in month zero. They found that the westerly phase in the lower stratosphere tend to last longer at solar minimum than at solar maximum, with the largest difference observed at the 40-50hPa levels.

*Gabis and Troshichev* [2006] pointed out that their data analysis is not consistent with the implicit assumption of *Soukharev and Hood* [2001] and *Salby and Callaghan* [2000] that there are more stalling of the easterlies and the prolongation of the westerlies in years of solar minimum since almost half of all short QBO durations occur near solar minima (1962, 1974, 1996 and 1998).

There are other questions that remain unanswered. For example, if stalling of the easterlies at 30 hPa is the relevant mechanism for the prolonged duration of the westerly phase below, should one expect the period of the QBO to be different above and below 30 hPa? When the duration of the westerly phase is prolonged, does the duration of the easterly phase become shortened so that the period of the QBO itself is unchanged? In the GCM model experiment of *Palmer and Gray* [2005] both the easterly duration and the westerly duration are shortened during solar max, in contrast to the 2D model of *McCormack* [2003], where as westerly is shortened, easterly is lengthened, during solar max.

93 *Hamilton* [2002] examined a longer equatorial record (than *Salby and Callaghan* [2000])  
 94 from 1953 to 2001, also from Free University of Berlin. He found that while the correlation  
 95 with the solar flux and the westerly duration is -0.46 over the 17 westerly phases during  
 96 the 1956-1996 period studied by *Salby and Callaghan* [2000], the correlation falls to an  
 97 insignificant -0.10 when computed over the 22 westerly phases in the longer record. In  
 98 particular Hamilton pointed out that towards the end of his record in the 21th century,  
 99 the relationship discovered by Salby and Callaghan appears to fail. It would be interesting  
 100 to examine a longer data record further into the 21th century, when the stratosphere is  
 101 not known to be contaminated by a major volcanic eruption. The eruption of Pinatubo  
 102 in 1991 was cited as a possible reason for some of the problems with the correlation of  
 103 *Salby and Callaghan* [2000] in the early 1990s.

104 Using 44 years of ERA-40 data from January 1958 to December 2001, spanning 18.5  
 105 QBO cycles, *Pascoe et al.* [2005] arrived at a conclusion consistent with that of *Salby*  
 106 *and Callaghan* [2000] and *Soukharev and Hood* [2001]. They found that the mean time  
 107 for the easterly shear zone to descend from 20 to 44 hPa is 2 months less under solar-  
 108 max conditions than under solar min conditions. This rapid descent of the easterly shear  
 109 zone cuts short the westerly phase of QBO in the lower stratosphere during solar max  
 110 periods. In particular the authors found that a Spearson's rank correlation with the solar  
 111 radio flux of the easterly descent rate for the period from 1958 to 1990 is a rather high  
 112 0.84 at 14 month lag. However, they also pointed out that the correlation breaks down  
 113 during the 1990s, but they attributed the anomalous climate of the tropical atmosphere  
 114 after the eruption of Pinatubo in June 1991 for this breakdown. This explanation can

possibly be ruled out if we extend the data to 2007, since volcanic aerosols do not stay in the stratosphere most likely not longer than 3 years. Most of the radiative effects by volcanic aerosols are from sulphate aerosols formed as a result of oxidation of the sulphate gas emitted by explosive volcanic eruptions into the stratosphere. The sedimentation e-folding time for sulphate aerosols is typically about one year [*Lambert et al.*, 1993], while most of the other ash particulates sediment out of the stratosphere within three months. *Ramaswamy et al.* [2001] in the IPCC 3rd Assessment Report and *Forster et al.* [2007] in the 4th Assessment Report commented that the stratosphere is now the cleanest since the satellite era.

In this work, we shall reexamine the possibility of a decadal solar-cycle modulation of the period of the QBO using an objective method and the longest record available. It is the same in situ dataset of near equatorial wind at 50 hPa that *Hamilton* [2002] used, distributed by the Stratospheric Research Group at the Free University of Berlin (FUB), here updated by B. Naujokat to span from 1953 to 2007. Figure 1 shows the time-height cross-section of the FUB wind. It shows the descent of the easterly shear zone to have more inter-annual variability compared with that of the westerly shear zone, as pointed out by previous authors. In this extended record, the stalling of the easterlies and the subsequent prolongation of the duration of the westerlies occurred irregularly: under solar min conditions of 1964, 1977 and 1987, but also under solar max conditions of 1967 and 2000. The corresponding time-height cross-section of the ERA-40 data can be found in *Pascoe et al.* [2005], and no update is available. Figure 2 shows this 40-year record up to 1 hPa in a time-height cross-section. The extended height coverage shows the transition



137 of the Semi-Annual Oscillation (SAO) near the stratopause into the QBO below, and  
 138 the tendency of synchronization of certain westerly phases of the two phenomena, as  
 139 the QBO always starts in a westerly SAO and ends when another SAO (several SAO  
 140 periods later) descends into the QBO altitude. As the QBO descends into the mid and  
 141 lower stratosphere, there are large (up to 12 month) changes in its easterly and westerly  
 142 durations, but the whole QBO period appears to be within one to two months of its value  
 143 in the upper stratosphere. These features of the QBO will be demonstrated more clearly  
 144 later in this paper. They impose a constraint on the possible mechanism(s) responsible  
 145 for the decadal variation of the QBO period. For example, since the radiative heating by  
 146 the volcanic aerosols is not expected to be vertically uniform throughout the stratosphere,  
 147 it cannot be responsible for the vertically uniform part of decadal variation of the QBO  
 148 period. Its effect on the QBO period is then seen to be at most 2 months, the maximum  
 149 extent of the vertical variation in period.

## 2. Methods

150 We use both an objective method and a subjective method for determining the QBO  
 151 period, each has its own advantages and disadvantages. Calculating the period of the  
 152 QBO has been a subjective procedure. It has usually involved visually determining when  
 153 a descending westerly (or easterly) first crosses zero at a particular level and when it later  
 154 goes back above zero. Such a procedure is sensitive to calibration errors and monthly  
 155 averaging. At the lower levels, such as 70 hPa, the presence of higher frequency zero  
 156 crossings renders this subjective method less useful, as there are many zero crossings  
 157 in sub QBO periods. At the upper levels, at 7 hPa and above, the method also fails

because of the simultaneous presence of SAO and QBO. The use of Continuous Wavelet Transform, which can determine the local intrinsic period of an oscillation, gives a more objective method that is not sensitive to the location of the zero-wind line. With this method the QBO period can be determined from 70 hPa to 1 hPa. However, this method does not separately determine the westerly and the easterly parts of the QBO period; for that we still need to rely on the traditional subjective methods. Most of our results do not depend critically on the use of the CWT method, and can be reproduced, but with some ambiguity, by some subjective methods, except in the upper stratosphere. The mathematical details of the CWT method are discussed in the Appendix.

Note that because of the continuous nature of the CWT method, the period thus determined varies smoothly, and does not show, for example, the discrete jump from 24 months to 30 months in the QBO period sometimes seen in the raw data. What is seen in the CWT result should be interpreted as some running average of neighboring periods.

### 3. Results and Discussion

#### 3.1. QBO-period variation

Figure 3 shows the local period of the QBO oscillation as determined by applying the CWT to the FUB data at 50 hPa for the period 1953-2007. This radiosonde dataset of near equatorial zonal wind consists of observations at Canton Island (January 1953 - August 1967) , Gan, Maldives (September 1967 - December 1975) and Singapore (since January 1976). It is the extended version of that used by *Salby and Callaghan* [2000] and *Hamilton* [2002]. It shows decadal variations around the mean period of 28 months. Because of this variation of periods, a Fourier analysis would give a broad spectrum of

QBO period but is unable to locate the times with long or short periods. The color scheme in this figure shows the amplitude of the wavelet coefficients, with darker color indicating higher amplitudes. There is a decadal *amplitude* modulation of the QBO, which will be the subject of a separate paper. Here we focus on the frequency modulation. The maximum amplitude of these wavelet coefficients is marked in a dashed line in Figure 3. This marks the dominant period of the equatorial zonal wind, i.e. the period associated with the most amplitude (or kinetic energy). This is the period we will be focusing on. Below it we also plot the sunspot number as a function of years, which is used as a proxy for the 11-year solar-cycle flux.

Figure 3 shows that, consistent with *Salby and Callaghan* [2000], the period of the QBO reaches its maximum during the solar min of 1965, when the dominant period is 33 months, the solar min of 1976, when the dominant period is 30 months, and the solar min of 1986, when the dominant period is 31 months. Other than these three solar minima mentioned by *Salby and Callaghan* [2000], however, the anti-correlation with the solar cycle breaks down. In the solar min of 1997, the dominant QBO period reaches a low of close to 25 months, consistent instead with the finding of *Hamilton* [2002]. Going forward in time, the correlation is the reverse of that of *Salby and Callaghan* [2000]. That is, during solar max, the QBO period is longer, while during solar min the QBO period is shorter. The in-phase relationship appears to commence around 1991, and probably as early as the late 1970 if one considers relative variations (see Figure 4 later). Prior to 1960 (from 1955-1960), the period variation is also approximately in-phase with the solar cycle, as Hamilton already pointed out. Over the almost 6 cycles spanned by the FUB

data, three cycles show anti-correlation of QBO period with solar flux, while the other two and a half cycles show in-phase correlation. As a consequence, the correlation coefficient between the two is close to zero (-0.03) for the long record of 1953-2007. It is intriguing to note the alternate correlation and anti-correlation of the QBO period with the solar cycle, which is a different behavior than two curves not related to each other at all that could also give a zero correlation coefficient. The calculation is repeated with the ERA-40 data. The result is very close to what we have shown here using the FUB data for the period of overlap.

This CWT procedure is repeated for all levels of the FUB data and the result is plotted in Figure 4. It shows the same prominent decadal variation at all levels and that the differences are small with respect to height. Minor differences of about a month in period exist at the higher levels (20 and 15 hPa) and at 70hPa prior to 1958. *Gabis and Troshichev* [2006] previously demonstrated the constancy of the QBO period with height, but only in an average for the period of 1953-2003; see their Figure 2c. Here we showed that this is true locally at each time in the record.

Since this plot uses an enlarged scale for the QBO period, it shows the variation of the period during 1980s and 1990s better than in Figure 3. The almost in-phase relationship between the relative QBO-period variation and the solar index appears as early as late 1970s and lasts till the end of the record in 2007, although in the 1980s and 1990s, the QBO variation appears to lead the solar index variation.

While the period variation of the QBO is almost the same at all heights in the lower stratosphere, above 30 hPa it is the easterly period variation that is responsible for most

of the variation of the whole QBO period, but below 30 hPa it is the westerly period that controls the whole QBO period variation. Figure 5 shows that, interestingly, the easterly period at 15 hPa and the westerly period at 50 hPa vary synchronously. The amplitude of the variation is also about the same, from 12 to 23 months. This observational result can be understood as follows. When the mean equatorial upwelling is strong it slows down the descent of the easterlies. The effect being more noticeable on the easterlies than on the westerlies as explained in the Introduction. Because the descent of the QBO east phase is preferentially stalled near the 30 hPa level in NH winter-spring, the duration of the easterly QBO phase is typically longer than the westerly phase above this level while the reverse is true below this level. This description explains the differing behavior of easterlies and westerlies above and below the stalling level, while the whole period of the QBO remains the same at these levels. Mechanistically, the upward propagating gravity waves and equatorial waves that are responsible for inducing the descent of the easterlies and westerlies need to pass through the region at 50 hPa of a prolonged duration of westerlies. The filtering effect by the lower stratospheric westerlies means that there is no deposition of westerly wave momentum above this level. As a result, the easterly duration is prolonged at 15 hPa. In the upper stratosphere, the easterly phase ends at around 3 hPa when the westerly wanes in the lower stratosphere, and westerly-phased gravity waves again propagate up to the upper stratosphere and initiate the next westerly descent. It is in this way that the upper level easterlies and lower level westerlies are related in their duration.

### 3.2. Comparison with the subjective methods

Figure 6 shows in the upper panel a comparison of the period as determined objectively using Continuous Wavelet Transform and the subjectively determined period by measuring the period between successive zero crossing of the zonal wind. It turns out that the result for the QBO period obtained by the subjective method is very different depending on whether one defines the full QBO period as easterly plus westerly, or as westerly plus easterly. The easterly plus westerly period appears to have a few more oscillations in 1980-1990 than the period as determined by westerly plus easterly durations. Our method turns out to be consistent with the average of these two definitions, provided that monthly averages are used in the subjective method.

On the lower panel of Figure 6, the “period” of the westerly phase and that of the easterly phase are separately determined by the subjective method. These are consistent with the results of *Salby and Callaghan* [2000] and *Hamilton* [2002], but for the longer data record. It shows a decadal variation of the westerly period that tracks quite closely that of the full QBO period as determined by the Continuous Wavelet Transform. Therefore our conclusion that there is no correlation of the QBO period with the solar cycle also applies to the westerly phase duration separately. The correlation coefficient of the westerly-phase duration variation with the sunspot time series for the full period of 1953-2007 is less than -0.10, also negligible.

*Hamilton and Hsieh* [2002] proposed using the circular nonlinear principal component analysis to objectively analyze and characterize the quasi-periodic QBO oscillation. They found that a single time series of the QBO phase can be found for data at all levels.

Although their method is not specifically aimed at studying the frequency variation of the QBO, a period variation similar to our CWT result was obtained in their Figure 10, but with much high-frequency irregular oscillations. *Hamilton and Hsieh* [2002] also concluded, based on their shorter record, that there is “no clear connection with the 11-year solar cycle”.

### 3.3. Behavior at different pressure levels in the ERA-40 data

These CWT calculations were repeated with the ERA-40 data up to 1 hPa. (Above that level the QBO is no longer the dominant oscillation because of the presence of SAO, and the CWT method does not yield an unambiguous QBO period.) The results are similar to those obtained using the FUB data for the period, except that the ERA-40 data extend to higher levels. This is shown in Figure 7. It is seen that the QBO period is almost constant in height, with the possible exception during the decade of 1980-1990, when changes of about one to two months in QBO period are seen around 20 hPa. During this period, El Chichon erupted in 1982 and Pinatubo in 1991. This implies the influence of the volcanic aerosols on QBO period is at most 2 months. In the recent decade, when the stratosphere is clean, the QBO period is again almost independent of height, to within one month. This again implies that perturbations to the stratosphere, such as ENSO and SSWs, which should be nonuniform with height, are not the dominant mechanism affecting the QBO period.

We show the CWT frequency-time plot at 2 hPa in Figure 8. Even at this high altitude, the QBO period and its variations are well separated (in frequency domain) from that of the SAO. The amplitude connections between SAO, the annual cycle and the QBO are

seen in vertical strings of darker color, which show the synchronization of the QBO with the SAO and the annual cycle.

### 3.4. Lagged correlations

It has been suggested that the correlation of the westerly-phase duration variation and the sunspot time series can be improved by considering lags between the two signals, and by looking at 40 hPa (L.L. Hood, personal communication, 2007). We have mentioned earlier that the QBO period is in phase with the solar cycle index in the decades of 1990s and 2000s, and also over half a cycle in the 1950s. This in-phase relationship is opposite to the behavior in the intervening three decades. This fact does not change with lagged correlation. Nevertheless, during the middle three decades considered by *Salby and Callaghan* [2000], the anti-correlation coefficient can be improved with 22-month lag. The improvement is large enough that even with the opposite phased correlation in the later decades the overall anti-correlation is improved in the extended record considered here.

The time series for period durations obtained using the subjective method has only one data point for each QBO period, and needs to be interpolated for the consideration of lagged correlation. Using a piecewise cubic Hermite interpolation method, the whole QBO period time series and the westerly phase duration time series have been produced with monthly increments at 40 hPa. At this altitude, the period considered by *Salby and Callaghan* [2000], 1956-1996, yields a greatly improved anti-correlation with the solar index of -0.79 for the westerly duration at 22 month lag, while for the period 1957-1999 studied by *Soukharev and Hood* [2001] the anti-correlation of the westerly duration with solar cycle is improved to -0.74 with 22 months lag. These correlation coefficients remain



significant, though smaller, at -0.64, over the extended period studied by us, while the zero-lag correlation is much smaller at -0.27.

For the whole QBO period, the correlation coefficient is also improved by considering a lag, but the improvement is not large enough to be statistically significant. (The correlation coefficient at zero lag is roughly zero for the full QBO signal, and reaches a maximum magnitude of -0.32 with 27 month lag.)

Despite this improvement in anti-correlation of the westerly duration at 40 hPa with the solar index, the fact remains that the recent decade shows opposite phased behavior than the middle three decades. This may be another sign that the QBO period response to solar cycle forcing may be non-stationary, and that a much longer time series than the one currently available is needed to see if the two are correlated, and to determine the sign of the correlation.

#### 4. Conclusion

Previous authors have pointed to the anti-correlation of the duration of the westerly phase of the QBO at the equator with the solar cycle in the three decades, 1960s, 1970s and 1980s, in the lower stratosphere. In particular, during the three solar minima in this period the duration of the westerly phase reaches a local maximum. In this paper an expanded study is undertaken for the longest period of record available from the Free University of Berlin, and for the levels between 70 and 15 hPa. This study is supplemented by the ERA-40 data, up to 1 hPa.

We have three main results: (1) The period of the QBO is almost the same in the vertical in the stratosphere. (2) There is a decadal variation of the QBO period of up to 12 months;

below 30 hPa such a variation of the whole QBO period is reflected in the variation of the westerly duration, while above 30 hPa it is the easterly duration that follows the decadal variation of the whole QBO period. (3) The decadal variation of the whole QBO period, and that of the westerly duration below 30 hPa, shows anti-correlation with the solar cycle during the middle three decades but positive correlation in the remaining decades in the 55-year record examined. This may be a sign of non-stationary behavior. These results are not entirely new, and have been mentioned by previous authors. It is now confirmed by us using the longest stratospheric record available. The synthesis of these results is discussed below.

Previous attention has focused on the decadal variation of the duration of the westerly QBO at the equator in the lower stratosphere, especially at around 40-50 hPa, and pointed to the mechanism of the stalling of the descent of the easterly shear zone near 30 hPa in prolonging the westerly duration below. Our new result shows that there is the same decadal variation of the whole QBO period, and so this proposed mechanism on the descent rate of easterly shear zones, if it works, only explains the partition of the whole QBO period into the two parts: Below 30 hPa, the westerly duration is longer than the easterly duration and it is the former which contains the decadal variation in the whole QBO period. Since the same decadal variation of the whole QBO period exists throughout the stratosphere, the origin of its variability may arguably be above the level we examined, although this conclusion is only tentative until checked by a model. This is because the wave-mean flow interaction in the QBO phenomenon is known to be able to transmit lower stratospheric influence to the upper levels through the filtering effect (see the discussion

earlier on the effect of a prolonged westerly region in the lower stratosphere delaying the onset of the next westerly phase in the upper stratosphere) . A simple possibility is the radiative perturbation by the solar max in the upper stratosphere that affects the QBO period in that photochemical region. This same decadal period variation is then preserved for all heights in the stratosphere with small variations of a month or two, much smaller than the decadal variation itself, which can be as large as 12 months. These small variations of a month or two can be caused by a number of mechanisms, such as volcanic aerosol heating, ENSO or polar stratospheric sudden warming.

With respect to what the data reveals concerning the solar-cycle modulation of the QBO period, we have found that in the longer equatorial zonal wind dataset from Free University of Berlin (1953-2007), which spans almost six solar cycles, the conclusion is mixed. During three of the cycles the period of the QBO is anti-correlated with the solar cycle, while in the remaining almost three cycles, there is correlation with the solar-cycle flux. The period previously considered by *Salby and Callaghan* [2000] contains three anti-correlated periods with one “straddling” period in 1992, which could have been discounted as due to Pinatubo. With our longer record extending into 2007, when there has not been a major volcanic eruption since 1991, it becomes more difficult to attribute the “anomalous” behavior to volcanic aerosols. We are not suggesting that there is no solar modulation of the QBO period, but the possibility exists that there is considerable unforced variability in the QBO period, so that the time series of the QBO is non-stationary even if the solar-cycle forcing is held fixed, as in the model of Mayr et al (2003). This is also consistent with the suggestion of *McCormack et al.* [2007] that perhaps 150 years are needed to identify

a statistically significant modulation of the QBO period in a 2D model simulation. This possibility will be examined more in a forthcoming modeling paper.

**Acknowledgments.** The research was supported by the National Science Foundation, Climate Dynamics Program, under grant ATM-0332364, and the DGA (French Defense Department) under contract 06.60.018.00.470.75.01. KK Tung would like to thank Professor Yuk Yung, Dr Run-Lie Shin and graduate student Le Kuai of Caltech, whose modeling work raises the question of whether the QBO period should become longer or shorter if the solar-cycle variation is magnified, and prompted the authors to reexamine the existing data for our current climate. P. Fischer would like to thank Dr. Dianne Casper for many fruitful conversations. The authors wish to thank Dr. Lon Hood for his helpful suggestions on the lagged correlation. We are also grateful to two anonymous reviewers for their detailed comments and suggestions on an earlier version of the manuscript.

## 5. Appendix: Continuous Wavelet Transform

We recall here only the main ideas underlying the wavelet theory and we refer to *Mallat* [1998] for a complete description. Like the complex exponential functions of the Fourier decomposition, wavelets can be used as basis functions for the representation of a signal. But, unlike the complex exponential functions, they are able to restore the temporal information as well as the frequency information. Functions depending on two real variables  $a$  and  $b$ , linked to frequency and time, respectively, are used to define the mathematical transformation:

$$WT_b^a = \int dt s(t) \psi_{a,b}(t), \quad (1)$$

where  $\psi_{a,b}(t)$  plays the same part as the exponential functions in the Fourier transform. Starting with a function  $\psi$  well localized in time and frequency spaces, a family of analyzing functions can be constructed:

$$\psi_{a,b}(t) = |a|^{-1/2} \psi \left( \frac{t-b}{a} \right). \quad (2)$$

where  $b$  is a time parameter and  $a$  is analogous to a period. The initial function  $\psi$  is called the mother wavelet and has to verify the following condition:

$$\int d\xi \frac{|\hat{\psi}|^2}{|\xi|} = K < \infty. \quad (3)$$

This condition means that any oscillating function localized in both spaces and whose integral over the whole space  $\mathbb{R}$  is null can be used as a mother wavelet. Similar to the definition of the inverse Fourier transform, it is also possible to define a reconstruction formula that allows one to rewrite  $s(t)$  as an expansion on the corresponding basis. The coefficients  $WT_b^a$  defined in (1) give a time-scale representation of the initial signal allowing the detection of transient components or singularities.

The results presented in this paper have been obtained using the tenth derivative of the Gaussian as a wavelet mother. Higher derivatives imply a longer support (i.e. range over a longer time window), and therefore yield a smoother result. When the support is too long, CWT begins to resemble Fourier transform and does not yield local time

information anymore. The chosen wavelet mother is appropriate for the examination of  
 decadal variation of QBO period. A link between the Fourier period and the wavelet  
 scale can be derived analytically for each wavelet mother. The results are given here in  
 a time-period representation rather than the usual time-frequency representation of the  
 wavelet theory.

## References

- Angel J.K. (1986), On the variation in period and amplitude of the quasi-biennial oscillation in the equatorial stratosphere 1951-1985, *Monthly Weather Rev.*, *114*, 2272–2278.
- Baldwin, M., L. Gray, T. Dunkerton, K. Hamilton, P. Haynes, W. Randel, J. Holton, M. Alexander, I. Hirota, T. Horinouchi, D. Jones, J. Kinnnersley, C. Marquardt, K. Sato, and M. Takahashi (2001), The quasi-biennial oscillation, *Reviews of Geophysics*, *39*, 179–229.
- Camp, C. and K-K. Tung (2007), The influence of the solar cycle and QBO on the late winter stratospheric polar vortex, *J. Atmos. Sci.*, *64*, 1267–1283.
- Dunkerton, T. (1990), Annual variation of deseasonalized mean flow acceleration in the equatorial lower stratosphere, *J. Met. Soc. Japan*, *68*, 499–508.
- Forster, P., et al. (2007), Changes in Atmospheric Constituents and Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis*.
- Gabis, I. and O. Troshichev (2005), QBO cycle identified by changes in height profile of the zonal winds: new regularities. *J. Atmos. and Solar-Terr. Phys.*, *67*, 33-44.
- Gabis, I. and O. Troshichev (2006), Influence of solar UV irradiance on the quasi-biennial oscillation of zonal winds in the equatorial stratosphere, *J. Atmos. and Solar-Terr.*

426 *Phys.*, 68, 1987-1999.

427 Hamilton, K. (2002), On the quasi-decadal modulation of the stratospheric QBO period,  
428 *J. Climate*, 15, 2562–2565.

429 Hamilton, K. and W. Hsieh (2002), Representation of the quasi-biennial oscillation in  
430 the tropical stratospheric wind by nonlinear principal component analysis, *J. Geophys.*  
431 *Res.*, 10 D15, 4232, 10.1029/2001JD001250.

432 Holton, J. and R. Lindzen (1972), Updated theory for quasi-biennial cycle of tropical  
433 stratosphere, *J. Atmos. Sci.*, 29, 1076–1080.

434 Holton, J., P.H. Haynes, M.E. McIntyre, A.R. Douglass, R. B. Hood and L. Pfister (1995),  
435 Stratosphere-troposphere exchange, *Rev. Geophys.*, 33, 403-439.

436 Hood, L.L. and B.E. Soukharev (2003), Quasi-decadal variability of the tropical lower  
437 stratosphere: the role of extratropical wave forcing, *J. Atmos. Sci.*, 60, 2389–2403.

438 Kinnersley, J. and S. Pawson (1996), The descent rates of the shear zones of the equatorial  
439 QBO, *J. Atmos. Sci.*, 53, 1937–1949.

440 Kodera, K., and Y. Kuroda (2002), Dynamical response to the solar cycle, *J. Geophys.*  
441 *Res.*, 107, 4749, doi:10.1029/2002JD002224.

442 Labitzke, K. (1982), On the interannual variability of the middle stratosphere during the  
443 northern winters, *J. Met. Soc. Japan*, 60, 124–139.

444 A. Lambert, et al., (1993), Measurements of the evolution of the Mt. Pinatubo aerosol  
445 cloud by ISAMS, *Geophys. Res. Lett.*, 20, 1287-1290.

446 Lindzen, R. and J. Holton (1968), A theory of quasi-biennial oscillation, *J. Atmos. Sci.*,  
447 25, 1095–1107.

Mallat, S. (1998), *A wavelet tour of signal processing*. Academic Press.

McCormack, J.P. (2003), the influence of the 11-year solar cycle on the quasi-biennial oscillation, *Geophys. Rev. Lett*, *30*, 2162, doi:10.1029/2003GL018314.

McCormack, J.P., D.E. Siskind, and L.L. Hood (2007), Solar-QBO interaction and its impact on stratospheric ozone in a zonally averaged photochemical transport model of the middle atmosphere, *J. Geophys. Res.*, *112*, D16109, doi:10.1029/2006JD008369.

Palmer, M.A., and L.J. Gray (2005), Modeling the atmospheric response to solar irradiance changes using a GCM with a realistic QBO, *Geophys. Rev. Lett*, *33*, L24701, doi:10.1029/2005GL023809.

Pascoe, C., L. Gray, S. Crooks, M. Jukes, and M. Baldwin (2005), The quasi-biennial oscillation: Analysis using era-40 data, *J. Geophys. Res.*, *110*, D08105, doi:10.1029/2004JD004941.

Plumb, R. and R. Bell (1982), A model of the quasi-biennial oscillation on an equatorial beta-plane, *Q.J.R. Meteorol. Soc.*, *108*, 335–352.

Quiroz, R. (1981), Period modulation of the stratospheric quasi-biennial oscillation. *Monthly Weather Review*, *109*, 665–674.

Ramaswamy, V et al. (2001), Radiative forcing of climate change. In: *Climate Change 2001, the Scientific Basis*. pp349-416.

Randel, W.J., R. Garcia, F. Wu (2002), Time-dependent upwelling in the tropical lower stratosphere estimated from the zonal-mean momentum budget. *J. Atmos. Sci.*, *59*, 2141–2152.



- 469 Salby, M. and P. Callaghan (2000), Connection between the solar cycle and the QBO:  
470 The missing link, *J. Climate*, *13*, 2652–2662.
- 471 Soukharev, B.E. and L.L. Hood (2001), Possible solar modulation of the equatorial quasi-  
472 biennial oscillation: Additional statistical evidence, *J. Geophys. Res.*, *106*, D14, 14855–  
473 14860.

**Figure captions:**

**Figure 1:** Time-height section of the FUB zonal wind. Easterlies are shown shaded.

Contour increment is  $5 \text{ m.s}^{-1}$

**Figure 2:** Time-height section of the monthly mean ERA-40 zonal wind, maintained at European Center for Medium Range Weather Forecasting. Easterlies are shown shaded.

Contour increment is  $5 \text{ m.s}^{-1}$ .

**Figure 3:** Local period in months of the 50 hPa FUB wind as determined by the CWT.

The darker region denotes location of higher amplitude of the zonal wind. The dashed line traces the location of the maximum amplitude. Superimposed, in solid line, is the sunspot number (monthly averaged), which is a proxy for the solar cycle radiative flux variability.

**Figure 4:** The QBO period as determined by the CWT method using the FUB data for various pressure levels in the lower stratosphere.


**Figure 5:** The period of the westerly phase at 50 hPa (in solid line) and that of the easterly phase at 15 hPa as a function of year.

**Figure 6:** Upper panel: A comparison of the QBO period as determined by the CWT method (in solid line) and that determined subjectively by adding the westerly phase period followed by the easterly phase period (in dashed line), and by adding the easterly phase period followed by the westerly phase period (in dotted line)

Lower panel: The westerly phase duration (in dashed line) and the easterly phase duration (in dotted line). All at 50 hPa.


496 **Figure 7:** The QBO period as determined by the CWT method using the ERA-40 data  
497 for various pressure levels in the stratosphere. Local maxima are shown shaded. Contour  
498 increment is one month.

499 **Figure 8:** Same as Figure 3, except for the zonal wind at 2 hPa (ERA-40).




**Figure 1.** Time-height section of the FUB zonal wind. Easterlies are shown shaded.


Contour increment is  $5 \text{ m.s}^{-1}$




**Figure 2.** Time-height section of the monthly mean ERA-40 zonal wind, maintained at European Center for Medium Range Weather Forecasting. Easterlies are shown shaded. Contour increment is  $5 \text{ m.s}^{-1}$ .



**Figure 3.** Local period in months of the 50 hPa FUB wind as determined by the CWT. The darker region denotes location of higher amplitude of the zonal wind. The dashed line traces the location of the maximum amplitude. Superimposed, in solid line, is the sunspot number (monthly averaged), which is a proxy for the solar cycle radiative flux variability.




**Figure 4.** The QBO period as determined by the CWT method using the FUB data for various pressure levels in the lower stratosphere.




**Figure 5.** The period of the westerly phase at 50 hPa (in solid line) and that of the easterly phase at 15 hPa as a function of year.






**Figure 6.** Upper panel: A comparison of the QBO period as determined by the CWT method (in solid line) and that determined subjectively by adding the westerly phase period followed by the easterly phase period (in dashed line), and by adding the easterly phase period followed by the westerly phase period (in dotted line)

Lower panel: The westerly phase duration (in dashed line) and the easterly phase duration (in dotted line). All at 50 hPa



**Figure 7.** The QBO period as determined by the CWT method using the ERA-40 data for various pressure levels in the stratosphere. Local maxima are shown shaded. Contour increment is one month.



**Figure 8.** Same as Figure 3, except for the zonal wind at 2 hPa.