

Dissipative boundary conditions for 2×2 hyperbolic systems of conservation laws for entropy solutions in BV*

Jean-Michel Coron[†] Sylvain Ervedoza[‡] Shyam Sundar Ghoshal[§]
Olivier Glass[¶] Vincent Perrollaz^{||}

Abstract

In this article, we investigate the BV stability of 2×2 hyperbolic systems of conservation laws with strictly positive velocities under dissipative boundary conditions. More precisely, we derive sufficient conditions guaranteeing the exponential stability of the system under consideration for entropy solutions in BV. Our proof is based on a front tracking algorithm used to construct approximate piecewise constants solutions whose BV norms are controlled through a Lyapunov functional. This Lyapunov functional is inspired by the one proposed in J. Glimm's seminal work [16], modified with some suitable weights in the spirit of the previous works [10, 9].

Contents

1	Introduction	2
1.1	Setting and main result	2
1.2	Previous results	4
1.3	Outline	8
2	Preliminaries	9
2.1	On the quantity $TV_{[0,L]}^*$	9
2.2	On the Riemann problem	10
2.2.1	Usual Riemann problem	10
2.2.2	Boundary Riemann problem	11

*This work was supported by the ERC advanced grant 266907 (CPDENL) of the 7th Research Framework Programme (FP7). The authors acknowledge the support of the UMI IFCAM (Indo-French Center for Applied Mathematics).

[†]Sorbonne Universités, UPMC Univ Paris 06, Laboratoire Jacques-Louis Lions, CNRS UMR 7598, 4 place Jussieu, 75252 Paris cedex 05, France. Email: coron@ann.jussieu.fr.

[‡]Université Paul Sabatier, Institut de Mathématiques de Toulouse, CNRS UMR 5219, 118 route de Narbonne 31062 Toulouse Cedex 9, France. Email: sylvain.ervedoza@math.univ-toulouse.fr.

[§]Gran Sasso Science Institute (GSSI), viale Francesco Crispi 7, 67100 L'Aquila, Italy. Email: shyam.ghoshal@gssi.infn.it.

[¶]Université Paris-Dauphine, Ceremade, CNRS UMR 7534, Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France. Email: glass@ceremade.dauphine.fr.

^{||}Université François Rabelais, Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 7350, Parc de Grandmont, 37200 Tours, France. Email: vincent.perrollaz@lmpt.univ-tours.fr.

3	Proof of Theorem 1.2	11
3.1	Construction of front-tracking approximations	11
3.2	A Glimm-type functional	13
3.3	Decay of Glimm's functional	14
3.4	Global definiteness of the approximations	19
3.5	Size of the rarefaction fronts	21
3.6	Lipschitz in time estimates	23
3.7	Passing to the limit: end of the proof of Theorem 1.2	25

1 Introduction

1.1 Setting and main result

The goal of this article is to study the exponential stability of 2×2 systems of conservation laws on a finite interval, by means of boundary feedbacks, in the context of weak entropy solutions. To be more precise, we consider the following setting: Let Ω be an open subset of \mathbb{R}^2 with $0 \in \Omega$, and $f : \Omega \mapsto \mathbb{R}^2$ be a smooth function (supposed to satisfy the strict hyperbolicity condition described below) and consider the following system of two conservation laws

$$\partial_t u + \partial_x(f(u)) = 0 \quad \text{for } (t, x) \in (0, \infty) \times (0, L). \quad (1.1)$$

In System (1.1), the solution $u = u(t, x) = (u_1, u_2)^T$ has 2 components and the space variable x belongs to the finite interval $(0, L)$. We assume the flux function f to satisfy the strictly hyperbolicity conditions, that is,

$$\forall u \in \Omega, \quad \text{the matrix } A(u) = Df(u) \text{ has 2 real distinct eigenvalues } \lambda_1(u) < \lambda_2(u). \quad (1.2)$$

Furthermore, we make the assumption that both velocities are positive, so that we finally get:

$$0 < \lambda_1(u) < \lambda_2(u) \quad \text{for all } u \in \Omega. \quad (1.3)$$

Note that the case of two strictly negative velocities is obviously equivalent by the change of variable $x \rightarrow L - x$.

System (1.1) is completed with boundary conditions of the form

$$u(t, 0) = Ku(t, L), \quad (1.4)$$

where K is a 2×2 (real) matrix. Here, for sake of simplicity, we assume that the boundary condition is linear but other nonlinear forms could be considered.

Clearly $u \equiv 0$ is an equilibrium of (1.1)-(1.4). We are interested in the exponential stability of this equilibrium in the BV space for entropy solutions. We recall that the space BV is natural for solutions of nonlinear hyperbolic systems of conservation laws, and is in particular the space considered in the celebrated paper by J. Glimm [16].

In order to discuss the condition that we impose on K , we further introduce the left and right eigenvectors of $A(u) = Df(u)$: for each $k = 1, 2$, we define $r_k(u)$ as a right eigenvector of $A(u)$ corresponding to the eigenvalue $\lambda_k(u)$:

$$A(u)r_k(u) = \lambda_k(u)r_k(u), \quad r_k(u) \neq 0, \quad k = 1, 2, \quad u \in \Omega. \quad (1.5)$$

We also introduce correspondingly left eigenvectors $\ell_k(u)$ of $A(u)$

$$\ell_k(u)A(u) = \lambda_k(u)\ell_k(u), \quad \text{with } \ell_k(u) \cdot r_{k'}(u) = \begin{cases} 1 & \text{if } k = k', \\ 0 & \text{if } k \neq k'. \end{cases} \quad (1.6)$$

We further impose that the hyperbolic system (1.1) is genuinely non-linear in the sense of Lax [19], i.e.

$$D\lambda_k(u) \cdot r_k(u) \neq 0 \text{ for all } u \in \Omega.$$

Changing the sign of $r_k(u)$ and $\ell_k(u)$ if necessary, we can therefore assume

$$D\lambda_k(u) \cdot r_k(u) > 0 \text{ for all } u \in \Omega. \quad (1.7)$$

As the total variation $\text{TV}_{[0,L]}$ is only a semi-norm on $BV(0, L)$ (it vanishes for constant maps), it is convenient to define the following norm on $BV(0, L)$ as

$$|u|_{BV} := \text{TV}_{[0,L]}(u) + \int_0^L |u(x)|dx, \quad u \in BV(0, L). \quad (1.8)$$

It is useful to recall that a function $u \in BV(0, L)$ has at most countably many discontinuities and have at each point left and right limits. In particular one can define $u(0^+)$ and $u(L^-)$ without ambiguity.

Now we recall that entropy solutions are weak solutions of (1.1) in the sense of distributions, which satisfy moreover *entropy conditions* for the sake of uniqueness. A way to express these entropy conditions consists in introducing entropy/entropy flux couples for (1.1) as any couple of regular functions $(\eta, q) : \Omega \rightarrow \mathbb{R}$ satisfying:

$$\forall U \in \Omega, \quad D\eta(U) \cdot Df(U) = Dq(U). \quad (1.9)$$

Of course $(\eta, q) = (\pm \text{Id}, \pm f)$ are entropy/entropy flux couples. Then we have the following definition (see [5, 12, 19]):

Definition 1.1. *A function $u \in L^\infty(0, T; BV(0, L)) \cap \mathcal{L}ip(0, T; L^1(0, L))$ is called an entropy solution of (1.1) when, for any entropy/entropy flux couple (η, q) , with η convex, one has in the sense of measures*

$$\eta(u)_t + q(u)_x \leq 0, \quad (1.10)$$

that is, for all $\varphi \in \mathcal{D}((0, T) \times (0, L))$ with $\varphi \geq 0$,

$$\int_{(0,T) \times (0,L)} (\eta(u(t, x))\varphi_t(t, x) + q(u(t, x))\varphi_x(t, x)) dx dt \geq 0. \quad (1.11)$$

These entropy conditions can be justified for instance by vanishing viscosity and are automatically satisfied by classical (by which we mean of class C^1) solutions.

Our main result is the following one:

Theorem 1.2. *Let the system (1.1) be strictly hyperbolic and genuinely nonlinear in the sense of (1.7), and assume that the velocities are positive in the sense of (1.3).*

If the matrix K satisfies

$$\inf_{\alpha \in (0, +\infty)} \left(\max \{ |\ell_1(0) \cdot Kr_1(0)| + \alpha |\ell_2(0) \cdot Kr_1(0)|, \right. \\ \left. \alpha^{-1} |\ell_1(0) \cdot Kr_2(0)| + |\ell_2(0) \cdot Kr_2(0)| \} \right) < 1, \quad (1.12)$$

then there exist positive constants $C, \nu, \epsilon_0 > 0$, such that for every $u_0 \in BV(0, L)$ satisfying

$$|u_0|_{BV} \leq \epsilon_0, \quad (1.13)$$

there exists an entropy solution u of (1.1) in $L^\infty(0, \infty; BV(0, L))$ satisfying $u(0, \cdot) = u_0(\cdot)$ and (1.4) for almost all times, such that

$$|u(t)|_{BV} \leq C \exp(-\nu t) |u_0|_{BV}, \quad t \geq 0. \quad (1.14)$$

In other words, Theorem 1.2 states that (1.12) is a sufficient condition for the (local) exponential stability of $u \equiv 0$ with respect to the BV -norm for (1.1) with the boundary condition (1.4).

Remark 1.3. *In Theorem 1.2, there is no claim of uniqueness of the solution of (1.1) with initial condition $u(0, \cdot) = u_0(\cdot)$ and boundary law (1.4). The initial-boundary value problem for hyperbolic systems of conservation laws is a very delicate matter, even in the non-characteristic case (when no characteristic speed vanishes). We refer for instance to Sablé-Tougeron [24], Amadori [1], Amadori-Colombo [2, 3], Colombo-Guerra [7], Donadello-Marson [15] and references therein. In particular, in the open loop case, it is possible to construct a Standard Riemann Semigroup as a limit of front-tracking approximations (which is also the construction method employed here). However, up to our knowledge, these results do not quite cover our feedback boundary condition (1.4), and no result of uniqueness in the same spirit as [5, Theorem 9.4] is available in our situation.*

Our approach will be based on the construction of a Lyapunov functional for solutions u of (1.1)-(1.4), in the spirit of earlier works by B. d'Andréa-Novel, G. Bastin and J.-M. Coron [10, 9]. The main difference with these works is that they consider Lyapunov functionals estimating the H^2 norm of the solutions, which are classical solutions and therefore forbids the presence of shocks. More recently in [8], a Lyapunov approach was also discussed to derive local stabilization results in C^1 . But we will rather design a Lyapunov functional that measures the BV -norm of the solution.

Our Lyapunov functional is actually inspired by the classical functional introduced by J. Glimm in [16], which measures the strength of interaction and a quadratic quantity measuring the “potential for future interactions”. In this celebrated paper, this functional was devoted to prove the existence of global solutions of hyperbolic systems in a small BV -neighborhood of a constant map. While the construction in [16] is applied to an approximating sequence generated by means of the so-called random choice method, we will follow the method of wave-front-tracking, originally introduced by C. Dafermos [13] in the context of scalar conservation laws, and extended in the context of 2×2 hyperbolic systems of conservation laws (as considered in this paper) by R. DiPerna in [14]. This method is at the core of the book [5] by A. Bressan to which we will refer several times in the following.

1.2 Previous results

Classical setting. We first recall some previous results on the exponential stability of $n \times n$ hyperbolic systems on the finite interval $(0, L)$ in the “classical case”, when no entropy conditions are needed.

For simplicity of this presentation, we assume that all the characteristic speeds are positive. Our dynamical system takes the form

$$\begin{cases} u_t + A(u)u_x = 0 & \text{for } (t, x) \in (0, \infty) \times (0, L), \\ u(t, 0) = G(u(t, L)) & \text{for } t \in (0, +\infty), \end{cases} \quad (1.15)$$

where $A : \Omega \rightarrow \mathbb{R}^{n \times n}$ and $G : \Omega \rightarrow \mathbb{R}^n$ are smooth maps such that

$$\text{for every } u \in \Omega, \text{ the eigenvalues of } A(u) \text{ are real, positive and distinct,} \quad (1.16)$$

$$G(0) = 0. \quad (1.17)$$

It follows from (1.17) that $u \equiv 0$ is an equilibrium of our dynamical system (1.15). Recall that one says that this equilibrium is exponentially stable for (1.15) with respect to the norm $|\cdot|_X$ on a linear space X of functions from $[0, L]$ into \mathbb{R}^n if there exist three positive constants C , ν and $\varepsilon_0 > 0$ such that, for every initial data $u_0 \in X$ satisfying the compatibility conditions adapted to X and such that

$$|u_0|_X \leq \varepsilon_0, \quad (1.18)$$

the solution u of (1.15) associated to the initial condition $u(0, \cdot) = u_0(\cdot)$ is such that

$$|u(t)|_X \leq C \exp(-\nu t) |u_0|_X \quad \text{for } t \geq 0. \quad (1.19)$$

Again to simplify the presentation, we moreover assume that

$$A(0) \text{ is a diagonal matrix.} \quad (1.20)$$

In fact, this can be assumed without loss of generality by performing a linear change of variables on u if necessary. Let

$$K := G'(0) \in \mathbb{R}^{n \times n}. \quad (1.21)$$

The linear case. The first studies on the exponential stability of 0 concern naturally the linear case:

$$A(u) = \Lambda = \text{diag}(\lambda_1, \dots, \lambda_n), \quad G(u) = Ku \quad \text{for } u \in \Omega. \quad (1.22)$$

Hence the dynamical system (1.15) is now

$$\begin{cases} u_t + \Lambda u_x = 0 & \text{for } (t, x) \in (0, \infty) \times (0, L), \\ u(t, 0) = Ku(t, L) & \text{for } t \in (0, +\infty). \end{cases} \quad (1.23)$$

Regarding X , we can consider various functional settings which give analogous results, due to the fact that we are in the linear regime. For instance the following classical functional spaces (with the associated usual norms) can be considered:

- (a) the Sobolev spaces $W^{m,p}$, with $m \in \mathbb{N}$ and $p \in [1, +\infty]$,
- (b) $C^m([0, L])$, with $m \in \mathbb{N}$,
- (c) $BV(0, L)$.

Then, whatever is X in the above list, 0 is exponentially stable for (1.23) for the norm $|\cdot|_X$ if and only if there exists $\delta > 0$ such that

$$\left\{ \det(\text{Id}_n - (\text{diag}(e^{-z/\lambda_1}, \dots, e^{-z/\lambda_n}))K) = 0, z \in \mathbb{C} \right\} \implies \Re(z) \leq -\delta. \quad (1.24)$$

See, e.g. (the proof of) [17, Theorem 3.5 on page 275]. For example, for

$$n = 2, A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, K = K_a := \begin{pmatrix} a & a \\ a & a \end{pmatrix}, \text{ with } a \in \mathbb{R}, \quad (1.25)$$

condition (1.24) is equivalent to

$$a \in (-1, 1/2). \quad (1.26)$$

See [17, p. 285]. Condition (1.24) is robust to small perturbations on K . However it turns out that this condition is not robust with respect to small perturbations on the λ_i . See again [17, p. 285]. This lack of robustness is of course a problem if one wants to treat the case of a nonlinear system by looking at its linearization at 0. We will say that the exponential stability of 0 is robust with respect to small perturbations on the λ_i 's if there exists $\varepsilon > 0$ such that, for every $(\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_n) \in \mathbb{R}^n$ such that

$$|\tilde{\lambda}_i - \lambda_i| \leq \varepsilon \quad \text{for } i = 1, \dots, n, \quad (1.27)$$

0 is exponentially stable for the perturbed hyperbolic system

$$\begin{cases} u_t + \tilde{\Lambda}u = 0 & \text{for } (t, x) \in (0, \infty) \times (0, L), \\ u(t, 0) = Ku(t, L) & \text{for } t \in (0, +\infty), \end{cases} \quad (1.28)$$

where

$$\tilde{\Lambda} := \text{diag}(\tilde{\lambda}_1, \dots, \tilde{\lambda}_n). \quad (1.29)$$

Then, Silkowski (see e.g. [17, Theorem 6.1 on page 286]) proved that for every X in the list (a)-(b)-(c), 0 is exponentially stable for (1.23) with an exponential stability which is robust with respect to small perturbations on the λ_i 's if and only if

$$\rho_0(K) < 1, \quad (1.30)$$

where

$$\rho_0(K) := \max \left\{ \rho(\text{diag}(e^{i\theta_1}, \dots, e^{i\theta_n})K); \theta_i \in \mathbb{R} \right\} < 1, \quad (1.31)$$

in which $\rho(M)$ denotes the spectral radius of a matrix M .

We point out that condition (1.30), in contrast with condition (1.24), does not depend on the λ_i 's. For Example (1.25), condition (1.30) is equivalent to

$$a \in (-1/2, 1/2), \quad (1.32)$$

which is more restrictive than condition (1.26).

Nonlinear systems. Let us now turn to the nonlinear system (1.15). The main previous known results on the exponential stability by means of feedback control concern, as far as we know, only the classical solutions. We introduce some notations. For $p \in [1, +\infty]$, let

$$\|x\|_p := \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \quad \text{for } x := (x_1, \dots, x_n)^T \in \mathbb{R}^n, \quad p \in [1, +\infty), \quad (1.33)$$

$$\|x\|_\infty := \max \{ |x_i|; i \in \{1, \dots, n\} \} \quad \text{for } x := (x_1, \dots, x_n)^T \in \mathbb{R}^n, \quad (1.34)$$

$$\|M\|_p := \max_{\|x\|_p=1} \|Mx\|_p \quad \text{for } M \in \mathbb{R}^{n \times n}, \quad (1.35)$$

$$\rho_p(M) := \inf \{ \|\Delta M \Delta^{-1}\|_p; \Delta \in \mathcal{D}_{n,+} \} \quad \text{for } M \in \mathbb{R}^{n \times n}, \quad (1.36)$$

where $\mathcal{D}_{n,+}$ denotes the set of all $n \times n$ real diagonal matrices whose entries on the diagonal are strictly positive. It is proved in [9] that

$$\rho_0(K) \leq \rho_p(K) \quad \text{for all } p \in [1, +\infty], \quad (1.37)$$

$$\rho_0(K) = \rho_2(K) \quad \text{for all } n \in \{1, 2, 3, 4, 5\}, \quad (1.38)$$

$$\text{for all } n > 5, \text{ there are } K \in \mathbb{R}^{n \times n} \text{ such that } \rho_0(K) < \rho_2(K). \quad (1.39)$$

(In fact, concerning (1.37), only the case $p = 2$ is treated in [9]; however the proof given in this paper can be adapted to treat the case of every $p \in [1, +\infty]$.) Let us point out that, for every $p \in [1, +\infty] \setminus \{2\}$ and every $n \geq 2$, there are examples of K such that inequality (1.37) is strict.

Now, we give known sufficient conditions for the exponential stability of 0 for (1.15) with respect to the $|\cdot|_X$ -norm.

- (i) If $X = C^m([0, L])$ with $m \in \mathbb{N} \setminus \{0\}$, a sufficient condition is:

$$\rho_\infty(K) < 1. \quad (1.40)$$

This result is proved by T. H. Qin [23], Y. C. Zhao [25], T. Li [20, Theorem 1.3 on page 173] and [8]. In fact, in [20, 23, 25], G is assumed to have a special structure; though, it is pointed out by J. de Halleux et al. in [18] that the case of a general G can be reduced to the case of this special structure. Moreover [20, 23, 25, 8] deal only with the case $m = 1$; but the proofs given there can be adapted to treat the general case $m \in \mathbb{N} \setminus \{0\}$.

- (ii) If X is the Sobolev space $W^{m,p}([0, L])$ with $m \in \mathbb{N} \setminus \{0, 1\}$ and $p \in [1, +\infty]$, a sufficient condition is:

$$\rho_p(K) < 1. \quad (1.41)$$

The case $p = 2$ is treated by J.-M. Coron, B. d'Andréa-Novel and G. Bastin in [9], and the case of general $p \in [1, +\infty]$ is treated by in J.-M. Coron and H.-M. Nguyen in [11]. In fact [9, 11] deal only with the case $m = 2$; nonetheless the proofs given there can be adapted to treat the general case $m \in \mathbb{N} \setminus \{0, 1\}$.

These conditions are only sufficient conditions for exponential stability. It is natural to ask if one can improve them. In particular, it is natural to ask if the condition $\rho_0(K) < 1$ (which seems to be the weakest possible sufficient condition: see above) is sufficient for the exponential stability in these spaces X . Of course this is true for $n = 1$. However it turns out to be false already for $n = 2$: as shown in [11], for every $n \geq 2$, there are analytic maps A such that, for every $\varepsilon > 0$, there are $K \in \mathbb{R}^{n \times n}$ satisfying

$$\rho_0(K) = \rho_2(K) < 1 < \rho_\infty(K) < 1 + \varepsilon, \quad (1.42)$$

such that, for every $m \in \mathbb{N} \setminus \{0\}$, 0 is not exponentially stable with respect to the C^m -norm for (1.15) with $G(u) := Ku$. Let us emphasize that, as already mentioned above, the first inequality of (1.42) implies that, for every $m \in \mathbb{N} \setminus \{0, 1\}$, 0 is exponentially stable with respect to the $W^{m,2}$ -norm for (1.15).

Finally, we point out that the right hand side of (1.12) is $\rho_1(K)$. Hence (1.12) is equivalent to

$$\rho_1(K) < 1. \quad (1.43)$$

Remark 1.4. *One has*

$$\rho_1(K) = \rho_\infty(K). \quad (1.44)$$

Indeed, for every matrix $M \in \mathbb{R}^{n \times n}$, one has

$$\|M\|_1 = \max \left\{ \sum_{j=1}^n |M_{ij}|; i \in \{1, \dots, n\} \right\}, \quad (1.45)$$

$$\|M\|_\infty = \max \left\{ \sum_{i=1}^n |M_{ij}|; j \in \{1, \dots, n\} \right\}. \quad (1.46)$$

In particular $\|M\|_\infty = \|M^T\|_1$, from which one easily gets

$$\rho_\infty(M) = \rho_1(M^T). \quad (1.47)$$

Using (1.45) and [20, Lemma 2.4, page 146], one has

$$\rho_1(M) = \rho_1(|M|) = \rho(|M|), \quad (1.48)$$

where, for $M \in \mathbb{R}^{n \times n}$, $\rho(M)$ is the spectral radius of M and $|M|$ is the $n \times n$ matrix whose entries are $|M|_{ij} := |M_{ij}|$. In particular

$$\rho_1(K) = \rho(|K|) = \rho(|K^T|) = \rho_1(K^T), \quad (1.49)$$

which, together with (1.47), implies (1.44).

The context of entropy solutions. As we mentioned earlier, very few results exist on the stabilization of hyperbolic systems of conservation laws in the context of entropy solutions. In fact, even in the scalar case, we are only aware of the work [22], in which a suitable stationary feedback law is shown to stabilize the solutions exponentially. Regarding the stabilization of hyperbolic systems of conservation laws, we know only two results obtaining asymptotic stabilization in open loop. The first one, obtained by A. Bressan and G.M. Coclite [6], established that for a general hyperbolic system of conservation laws with either genuinely nonlinear or linearly degenerate characteristic fields (in the sense of Lax), and characteristic speeds strictly separated from 0, one can steer asymptotically in time any initial condition on a finite interval with sufficiently small total variation to all close constant states, by suitably acting on both sides of the interval. Furthermore, the controllability result may depend on the considered class of solutions, as underlined by [6]. Indeed, in the example presented in the article [6], controllability holds in the context of classical solutions ([21]) but not in the context of weak entropy solutions, emphasizing that linearization techniques cannot be used in the BV class. The second result, due to F. Ancona and A. Marson [4] is concerned with a case of a control from a single boundary point rather than on both sides.

1.3 Outline

The article is organized as follows. In Section 2, we start with some basic remarks that will be needed in the proof of Theorem 1.2, recalling in particular some embeddings and the solvability of the Riemann problem away from the boundary and on the boundary. Section 3 gives the proof of Theorem 1.2. It is divided in several steps. First, Section 3.1 presents the construction of front-tracking approximations of solutions of (1.1)–(1.4). Section 3.2 then introduces the Lyapunov functional we will use, which is a suitably weighted Glimm functional, inspired in [16] and [10, 9]. Section 3.3 proves the exponential decay of this quantity, which is in fact the main step in our analysis. Though, in order to conclude, we shall provide further estimates, in particular guaranteeing that:

- our construction is valid for all time (Section 3.4);
- the rarefaction fronts remain small (Section 3.5);
- the solutions are uniformly bounded in $L^\infty(0, L; BV(0, T))$ for all $T > 0$ (Section 3.6).

One can then pass to the limit in our approximate solutions and prove that they converge to a suitable solution of (1.1)–(1.4) whose BV norm is exponentially decaying, see Section 3.7, thus finishing the proof of Theorem 1.2.

2 Preliminaries

Let us first point out that, using (1.12) and replacing, if necessary, $r_1(u)$ by $\alpha r_1(u)$ and $\ell_1(u)$ by $\alpha^{-1}\ell_1(u)$ for some suitable $\alpha > 0$, we may assume without loss of generality that

$$\max_{k=1,2} \{|\ell_1(0) \cdot Kr_k(0)| + |\ell_2(0) \cdot Kr_k(0)|\} < 1. \quad (2.1)$$

For the proof of Theorem 1.2, it will be convenient to define, for $u \in BV(0, L)$, the quantity $TV_{[0,L]}^*(u)$ as

$$TV_{[0,L]}^*(u) = TV_{[0,L]}(u) + |Ku(L-) - u(0+)|, \quad (2.2)$$

where $u(0+)$ and $u(L-)$ have to be understood respectively as the right and left limits of the function u in $x = 0$ and $x = L$, respectively. Note that this quantity is well-defined for all $u \in BV(0, L)$.

2.1 On the quantity $TV_{[0,L]}^*$

We prove the following Lemma:

Lemma 2.1. *Under assumption (2.1), the quantity $TV^*(0, L)(\cdot)$ is a norm on $BV(0, L)$ which is equivalent to the norm $|\cdot|_{BV}$. Consequently, there exists a constant C such that for all $u \in BV(0, L)$,*

$$\|u\|_{L^\infty(0,L)} \leq C TV_{[0,L]}^*(u). \quad (2.3)$$

Proof. Clearly, there exists a positive constant $C > 0$ such that

$$TV_{[0,L]}^*(u) \leq C|u|_{BV} \quad \text{for all } u \in BV(0, L).$$

For this property, of course, (2.1) is not needed. To get the inequality in the other direction, we notice that for $u \in BV(0, L)$, one has

$$|u(L-) - u(0+)| \leq TV_{[0,L]}(u) \quad \text{and} \quad |Ku(L-) - u(0+)| \leq TV_{[0,L]}^*(u), \quad (2.4)$$

so

$$|u(L-) - Ku(L-)| \leq 2TV_{[0,L]}^*(u).$$

Expressing $u(L-)$ in the basis $(r_1(0), r_2(0))$:

$$u(L-) = a_1 r_1(0) + a_2 r_2(0)$$

and using

$$Kr_k(0) = (\ell_1(0) \cdot Kr_k(0))r_1(0) + (\ell_2(0) \cdot Kr_k(0))r_2(0),$$

we obtain that $u(L-) - Ku(L-)$ has the following coordinates

$$\begin{pmatrix} a_1 - a_1(\ell_1(0) \cdot Kr_1(0)) + a_2(\ell_1(0) \cdot Kr_2(0)) \\ a_2 - a_1(\ell_2(0) \cdot Kr_1(0)) + a_2(\ell_2(0) \cdot Kr_2(0)) \end{pmatrix}.$$

Due to (2.1), the matrix

$$\text{Id} - \begin{pmatrix} \ell_1(0) \cdot Kr_1(0) & \ell_1(0) \cdot Kr_2(0) \\ \ell_2(0) \cdot Kr_1(0) & \ell_2(0) \cdot Kr_2(0) \end{pmatrix}$$

is invertible and consequently for some positive $C > 0$:

$$|u(L-)| \leq C|u(L-) - Ku(L-)|.$$

The conclusion follows easily, using that for all $x \in (0, L)$, $|u(x) - u(L-)| \leq TV_{[0,L]}(u)$. \square

2.2 On the Riemann problem

2.2.1 Usual Riemann problem

Let us make some brief reminders on the Riemann problem for (1.1). Details can be found for instance in [5, Chapter 5].

Following [5, Section 5.2], we introduce the Lax curves as the curve obtained by gluing of the admissible part of the Hugoniot locus and of the rarefaction curves, i.e.

$$\Psi_k(\sigma, u) = \begin{cases} S_k(\sigma, u) & \text{if } \sigma < 0, \\ R_k(\sigma, u) & \text{if } \sigma \geq 0. \end{cases} \quad (2.5)$$

Here, $R_k(\sigma, u)$ corresponds to the rarefaction curves, that is, the orbits of the vector fields r_k :

$$\frac{dR_k(s, u)}{ds} = r_k(R_k(s, u)), \quad s \in [0, \sigma], \quad R_k(0, u) = u.$$

The part corresponding to $\sigma \geq 0$ (due to (1.7)) is composed of points u_+ which can be connected to u from left to right by a rarefaction wave:

$$u(t, x) = \begin{cases} u & \text{if } x < \lambda_k(u)t, \\ R_k(\sigma, u) & \text{if } x = \lambda_k(R_k(\sigma, u))t, \\ u_+ & \text{if } x > \lambda_k(u_+)t. \end{cases} \quad (2.6)$$

On the other side $S_k(\sigma, u)$ stands for the shock curve, which describes the k -th branch of the Hugoniot locus which gathers point u_+ satisfying for a fixed state u the Rankine-Hugoniot condition:

$$f(u_+) - f(u) = s(u_+ - u), \quad s \in \mathbb{R}. \quad (2.7)$$

To be slightly more precise, the k -th shock curve is defined for σ small as follows:

$$u_+ = S_k(\sigma, u)$$

if and only if u_+ and u satisfy

$$u_+ - u = \sigma r_k(u_+, u),$$

where $r_k(u_+, u)$ is the (suitably normalized) k -th eigenvector of the matrix

$$A(u_+, u) = \int_0^1 Df(u + t(u_+ - u)) dt.$$

For $\sigma < 0$, points $u_+ = S_k(\sigma, u)$ can be connected to u from left to right by an admissible shock wave:

$$u(t, x) = \begin{cases} u & \text{if } x < st, \\ u_+ & \text{if } x > st, \end{cases} \quad (2.8)$$

where the shock speed s is equal to $\lambda_k(u_+, u)$, the k -th eigenvalue of the above matrix $A(u_+, u)$.

We recall that, under suitable parameterization, the function $(\sigma, u) \mapsto \Psi_k(\sigma, u)$ is of class C^2 , see for example [5, p. 99].

Finally, following Lax [19], using the implicit function theorem, we see that there exists $\delta > 0$ such that for all u_- and u_+ with $|u_-| \leq \delta$ and $|u_+| \leq \delta$, there exists σ_1 and σ_2 small such that

$$u_+ = \Psi_2(\sigma_2, \Psi_1(\sigma_1, u_-)), \quad (2.9)$$

and besides, there exists a constant $C > 0$ such that

$$\frac{1}{C}|u_+ - u_-| \leq |\sigma_1| + |\sigma_2| \leq C|u_+ - u_-|.$$

See for instance [5, Theorem 5.3].

In the following, when considering two states u_- and u_+ satisfying $u_+ = \Psi_k(\sigma, u_-)$, we will say that u_- and u_+ are connected through a k -wave of strength $|\sigma|$. If $\sigma \geq 0$, this wave is a k -rarefaction, while it is a k -shock otherwise.

2.2.2 Boundary Riemann problem

Since we are considering solutions to an initial-boundary value problem, we also need to consider the Riemann problem on the boundaries. Due to the assumption (1.3), we only consider the problem on the left side of the interval. Actually, under this assumption, this problem is fairly simple and solve as in the usual case. Let us consider indeed on the boundary $x = 0$ the conditions:

$$u(t, 0) = \begin{cases} u_- & \text{for } t > t_0, \\ u_+ & \text{for } t < t_0, \end{cases}$$

with u_- and u_+ sufficiently small. Then writing again (2.9), this problem can be solved as in the usual case by a 1-wave followed by a 2-wave (from left to right, that is from top to bottom).

Note that under the constraint (1.4), boundary conditions u_0 at $x = 0^+$ and u_L at $x = L$ generate a Riemann problem between u_0 and Ku_L at $x = 0$.

3 Proof of Theorem 1.2

In the following, we work under the assumptions of Theorem 1.2, and in particular (1.3) and (2.1). Theorem 1.2 is based on a construction of solutions of (1.1)–(1.4) relying on a wave-front tracking algorithm. This algorithm generates approximations of a solution u of (1.1), which have a particular shape. To be more precise, for h small, we look for $u_h = u_h(t, x)$ defined for $t \geq 0$ and $x \in [0, L]$ such that:

- u_h is a piecewise constant function on $\mathbb{R}_+ \times [0, L]$, with a finite number of discontinuities (locally in time), which are straight lines (called *fronts*),
- each front is either a *rarefaction front* or a *shock*. In the former case, the states on the sides of the discontinuities are connected by a rarefaction wave, in the latter by a shock,
- the rarefaction fronts are of strength $\mathcal{O}(h)$,
- the boundary condition is satisfied for all times (taking left and right limits at discontinuity points),
- a quantity equivalent to the TV^* norm of $u(t)$ decays exponentially as time evolves.

3.1 Construction of front-tracking approximations

We mainly follow R. DiPerna's strategy [14] consisting, starting from a piecewise constant approximation of the initial data, in solving the generated Riemann problems and replacing rarefaction waves with piecewise constant approximations called *rarefaction fans*. When two

discontinuities meet, the process is iterated, with an important convention on rarefactions, that is *they are not re-split* across an interaction. See below for a more precise description. This means in particular that there are two ways to treat rarefaction waves in the process; let us describe these two methods.

Rarefaction fans. A k -rarefaction wave (u_-, u_+) with $u_+ = R_k(\sigma, u_-)$, $\sigma > 0$, centered at (\bar{t}, \bar{x}) can be approximated by a k -rarefaction fan (of accuracy h) as follows. Set

$$p = \lceil \sigma/h \rceil.$$

If $p > 1$, i.e. if $\sigma > h$, we define the intermediate states for $j \in \{1, \dots, p\}$:

$$u_j = R_k(\sigma/p, u_{j-1}), \quad u_0 = u_-,$$

and define

$$x_{j+1/2}(t) = \bar{x} + \lambda_k(u_{j+1})(t - \bar{t}).$$

In that case, the rarefaction wave is approximated locally by

$$\tilde{u}(t, x) = \begin{cases} u_0 = u_- & \text{for } x < x_{1/2}(t) \\ u_j & \text{for } x \in (x_{j-1/2}(t), x_{j+1/2}(t)), j \in \{1, \dots, p-1\}, \\ u_p = u_+ & \text{for } x > x_{p-1/2}(t). \end{cases} \quad (3.1)$$

If $p < 1$, i.e. if $\sigma \in (0, h]$, we simply set

$$\tilde{u}(t, x) = \begin{cases} u_- & \text{for } x < x_{1/2}(t) \\ u_+ & \text{for } x > x_{1/2}(t), \end{cases}$$

Approximating rarefaction waves by a single front. In this case, the above Riemann problem is solved using only one front, i.e. the approximate solution \tilde{u} is locally given by

$$\tilde{u}(t, x) = \begin{cases} u_- & \text{for } x < \bar{x} + \lambda_k(u_+)(t - \bar{t}), \\ u_+ & \text{for } x > \bar{x} + \lambda_k(u_+)(t - \bar{t}). \end{cases} \quad (3.2)$$

The wave-front tracking algorithm. We start from an approximate sequence $u_{0,h} \in BV(0, L)$, $h > 0$, of the initial condition, satisfying

$$\|u_{0,h}\|_\infty \leq \|u_0\|_\infty, \quad TV(u_{0,h}) \leq TV(u_0), \quad u_{0,h} \xrightarrow{h \rightarrow 0} u_0 \text{ in } L^1(0, L). \quad (3.3)$$

Now the construction of an approximate solution of (1.1) is then done as follows:

1. At time $t = 0$, we construct u_h as the solution of the Riemann problems for $u(t = 0)$ for which all rarefaction waves are replaced by rarefaction fans (with accuracy h). This includes the Riemann problem generated by $u_{0,h}(0^+)$ and $Ku_{0,h}(L^-)$. We extend the resulting discontinuities (called fronts) as straight lines, until two of them meet (at a point called an *interaction point*), or until one of them meets the boundary (which is necessary the right one, since all fronts have positive speeds under the assumption (1.3)).
2. When a front hits the (right) boundary, we solve the corresponding Riemann boundary problem between $Ku_h(t-, L-)$ and $u_h(t-, 0+)$. Again in that case we approximate all outgoing rarefaction waves by rarefaction fans.
3. When two fronts (say of family k and ℓ) interact at some time t in some point $x \in (0, L)$:

- if $k = \ell$, solve the resulting Riemann problem between the leftmost and rightmost states and approximate the outgoing rarefaction wave of the family k (if any) by a single front and the outgoing rarefaction wave of the other family (if any) by a rarefaction fan.
- if $k \neq \ell$, solve the Riemann problem and approximate each outgoing rarefaction wave (if any) by a single front.

4. If at some time, three fronts (or more) interact in the interior of the domain or two fronts (or more) interact at the boundary $x = L$, we slightly change the velocity of one of the incoming fronts so that there is only two fronts meeting at the same time. These changes of velocity are done so that the new (constant) velocity c belongs to an h -neighborhood of the expected velocity. We also modify the velocities similarly to avoid having several interactions at the same time. (This is classical in the context of front tracking approximation, see [5].)

Our construction works as long as the number of interactions of fronts is finite and u_h stays in the set where we can solve the Riemann problem. Let us therefore define

$$T_h^* = \sup\{t > 0, \text{ such that the number of fronts is finite in } (0, t) \times (0, L) \text{ and } \|u_h\|_{L^\infty((0,t) \times (0,L))} \leq \delta_0\}, \quad (3.4)$$

where $\delta_0 \in (0, \delta)$ is a positive parameter fixed below by (3.5)–(3.6), δ being the parameter in (2.9).

Remark 3.1. *Note that $T_h^* > 0$ as the fronts propagate at bounded velocities and since the initial data is piecewise constant. We will later show that T_h^* actually is infinite for all $h > 0$, see Lemma 3.5.*

3.2 A Glimm-type functional

In order to get estimates on this approximate solution u_h of (1.1)–(1.4), we will introduce a functional resembling Glimm's one [16] and adapted to our problem.

We choose $\delta_0 \in (0, \delta)$, $c_* > 0$, $\gamma > 0$, $\varepsilon > 0$ such that

$$c_* < \min_{|u| \leq \delta_0} \lambda_1(u) < \max_{|u| \leq \delta_0} \lambda_1(u) < \min_{|u| \leq \delta_0} \lambda_2(u), \quad (3.5)$$

$$\max_{|u| \leq \delta_0} \max_{k \in \{1,2\}} \{|\ell_1(Ku) \cdot Kr_k(u)| + |\ell_2(Ku) \cdot Kr_k(u)|\} < \exp(-\gamma L) - 2\varepsilon. \quad (3.6)$$

This can be done according to the assumptions (1.3) and (2.1).

Next, for a piecewise constant function U on $[0, L]$ with $TV^*(U)$ small enough:

- we denote $x_1 < x_2 < \dots < x_n$ the set of discontinuities in $(0, L)$,
- for each $i \in \{1, \dots, n\}$, we let $u_{i,-}$ and $u_{i,+}$ the limits of U at x_i from the left and from the right respectively,
- for each $i \in \{1, \dots, n\}$, we introduce the values $\sigma_{i,1}$ and $\sigma_{i,2}$ such that

$$u_{i,+} = \Psi_2(\sigma_{i,2}, \Psi_1(\sigma_{i,1}, u_{i,-})),$$

which are obtained by the solvability of the Riemann problem,

- finally, we quantify the “interaction” on the boundary between $Ku(L-)$ and $u(0+)$ by $\sigma_{0,1}, \sigma_{0,2}$ such that

$$u(0+) = \Psi_2(\sigma_{0,2}, \Psi_1(\sigma_{0,1}, Ku(L-))),$$

whose existence is granted by the solvability of the boundary Riemann problem.

Now for such a piecewise constant function U , we introduce the following functionals:

$$V(U) = \sum_{i=0}^n (|\sigma_{i,1}| + |\sigma_{i,2}|) e^{-\gamma x_i}, \quad (3.7)$$

$$Q(U) = \sum_{(x_i, \sigma_i)} |\sigma_i| e^{-\gamma x_i} \left(\sum_{(x_j, \sigma_j) \text{ approaching } (x_i, \sigma_i)} |\sigma_j| e^{-\gamma x_j} \right), \quad (3.8)$$

where a front (x_j, σ_j) of a family k_j is said to be approaching of a front (x_i, σ_i) of a family k_i if and only if one of the conditions is satisfied:

- $x_j < x_i$ and $(k_i, k_j) = (1, 2)$,
- $x_j < x_i$, $k_i = k_j$ and at least one of σ_j or σ_i is negative.

We underline that the difference with the usual functionals of total strength and of interaction potential lies in the exponentials, and on the fact that the boundary is taken into account at index $i = 0$, emphasizing the special role played by the feedback operator.

As seen from the construction, at all times except in a discrete set, namely the times of interaction of fronts, the discontinuities in the approximation u_h given by the front tracking algorithm are connected through a wave of family 1 or 2, that is, for almost all $t > 0$, for each $x_i(t)$, either $\sigma_{i,1}(t)$ or $\sigma_{i,2}(t)$ vanishes. In that case, we will say that $x_i(t)$ corresponds to a k -front, where $k \in \{1, 2\}$ is such that $\sigma_{i,k} \neq 0$.

For later use, let us also point out that two rarefaction fronts of the same family cannot meet.

Notations. For sake of simplicity, we will use the slight abuse of notation $V(t) = V(u_h(t))$ and $Q(t) = Q(u_h(t))$.

3.3 Decay of Glimm’s functional

Our goal is to prove the following result:

Lemma 3.2. *Under the setting of Theorem 1.2, there exist positive constants $\varepsilon_0 > 0$, $c_0 > 0$, $\nu > 0$ such that for any $h > 0$, if we define the approximate solution $u_h(t)$ of (1.1) using the front tracking approximation explained above, starting from $u_{0,h}$ satisfying*

$$V(u_{0,h}) \leq \varepsilon_0, \quad (3.9)$$

then the functional

$$J(t) = V(u_h(t)) + c_0 Q(u_h(t)) \quad (3.10)$$

satisfies

$$J(t) \leq \exp(-\nu t) J(0), \quad t \in [0, T_h^*]. \quad (3.11)$$

We recall that the approximation is well-defined till T_h^* , see (3.4).

Proof. We split the proof in two parts: first we study the evolution of V and Q in various cases, and then we come back to the functional J .

1. Behavior of V and Q .

We fix a time $t_0 \in (0, T_h^*)$. We recall that interactions in the domain only involve two fronts and interaction on the boundary only involve one. We discuss the evolution of the functionals V and Q locally around t_0 according to three cases (see Figures 1 and 2):

- **Case 1:** There is no interaction at $t = t_0$ in $(0, L)$, nor on the boundary.
- **Case 2:** There is a front interaction at $t = t_0$ in $(0, L)$.
- **Case 3:** There is a front hitting the boundary at $t = t_0$.




Figure 1: Left, Case 1: no interaction at time t_0 . Right, Case 2: two fronts interact at $t = t_0$.

Case 1. We suppose that there is no interaction at $t = t_0$. Since there is a finite number of fronts at time t_0 , there exists a neighborhood \mathcal{T} of t_0 on which there is no interaction.

As the approximation $u_h(t_0)$ is a piecewise constant function, we call as before $x_1(t) < x_2(t) < \dots < x_n(t)$ the discontinuities at time $t \in \mathcal{T}$ and $u_{i,-}$, $u_{i,+}$ the limits of $u_h(t, \cdot)$ at x_i from the left and from the right. Note that by construction, each trajectory $t \mapsto x_i(t)$ corresponds to a k_i -front and the states $u_{i,-}$ and $u_{i,+}$ on its sides do not depend on $t \in \mathcal{T}$. Therefore, each front has constant strength and velocity and moreover, due to condition (3.5), one has $x'_i(t) \geq c_*$. Therefore, we obtain, for all $t \in \mathcal{T}$,

$$\frac{dV}{dt}(t) \leq -c_*\gamma V(t), \quad \frac{dQ}{dt}(t) \leq -2c_*\gamma Q(t). \quad (3.12)$$

Case 2. We suppose that two fronts interact in $(0, L)$. Here it is necessary to relate the strength of the outgoing fronts with the incoming ones. Following [16] (see also [5, Lemma 7.2]), we have the following Glimm's interaction estimates:

Lemma 3.3 ([16]). *Consider an interaction between two incoming wave-fronts and assume that the left, middle and right states u_L, u_M and u_R satisfy $|u_L| \leq \delta$, $|u_M| \leq \delta$ and $|u_R| \leq \delta$.*

- **Distinct family:** Let $\hat{\sigma}_1$ and $\hat{\sigma}_2$ be the sizes of two incoming fronts corresponding respectively to the family 1 and 2. Then the outgoing fronts have strength σ_1, σ_2 satisfying:

$$|\sigma_1 - \hat{\sigma}_1| + |\sigma_2 - \hat{\sigma}_2| \leq C_\delta |\hat{\sigma}_1| |\hat{\sigma}_2|. \quad (3.13)$$

- **Same family:** Let $\tilde{\sigma}, \hat{\sigma}$ be the sizes of two incoming fronts both belonging to the k -th characteristic family. Then the outgoing fronts have strength σ_k for the outgoing fronts belonging to the k -th characteristic family and $\sigma_{k'}$ for the other family and satisfy:

$$|\sigma_k - (\tilde{\sigma} + \hat{\sigma})| + |\sigma_{k'}| \leq C_\delta |\tilde{\sigma}| |\hat{\sigma}| (|\tilde{\sigma}| + |\hat{\sigma}|). \quad (3.14)$$

We assume that the interaction takes place at $t = t_0, x = x_i$, and we denote by $\sigma_i(t_0-), \sigma_{i+1}(t_0-)$ the strength of the incoming waves, and by $\sigma_{i,1}(t_0+), \sigma_{i,2}(t_0+)$ the strength of the outgoing waves of families 1 and 2.

Using Lemma 3.3 and arguing as in [16] (see also [5, 14]), provided $e^{\gamma L} V(t_0-) \leq 1$ to guarantee that $|\sigma_i(t_0-)| + |\sigma_{i+1}(t_0-)| \leq 1$, we then obtain

$$V(t_0+) - V(t_0-) \leq C_\delta |\sigma_i(t_0-)| |\sigma_{i+1}(t_0-)| e^{-\gamma x_i} \leq C_\delta |\sigma_i(t_0-)| |\sigma_{i+1}(t_0-)|,$$

while

$$Q(t_0+) - Q(t_0-) \leq |\sigma_i(t_0-)| |\sigma_{i+1}(t_0-)| (-e^{-2\gamma x_i} + e^{-\gamma x_i} C_\delta V(t_0-)),$$

where $V(t_0-), V(t_0+), Q(t_0-), Q(t_0+)$ denote the respective limits of $V(t_0 - \tau), V(t_0 + \tau), Q(t_0 - \tau)$ and $Q(t_0 + \tau)$ as τ goes to 0^+ .

In particular, provided that

$$\max\{e^{\gamma L}, 2C_\delta e^{2\gamma L}\} V(t_0-) \leq 1, \quad (3.15)$$

we have

$$Q(t_0+) - Q(t_0-) \leq -\frac{1}{2} e^{-2\gamma L} |\sigma_i(t_0-)| |\sigma_{i+1}(t_0-)|, \quad (3.16)$$

and

$$V(t_0+) + c_0 Q(t_0+) \leq V(t_0-) + c_0 Q(t_0-), \quad (3.17)$$

for the choice

$$c_0 = 2C_\delta e^{2\gamma L}.$$

Case 3. Here we suppose that one front hits the boundary at time $t = t_0, x = L$. The following lemma helps us to describe the fronts created in $x = 0$.

Lemma 3.4. Assume that a k front of strength $\hat{\sigma}$ hits the boundary at $t = t_0-, i.e.$

$$u_R = \Psi_k(\hat{\sigma}, u_L). \quad (3.18)$$

Then it generates at $t = t_0+$ and $x = 0$ two waves σ_1, σ_2 such that

$$Ku_R = \Psi_2(\sigma_2, \Psi_1(\sigma_1, Ku_L)), \quad (3.19)$$

with the estimate

$$|\sigma_1 - \hat{\sigma} \ell_1(Ku_L) \cdot Kr_k(u_L)| + |\sigma_2 - \hat{\sigma} \ell_2(Ku_L) \cdot Kr_k(u_L)| \leq C_b |\hat{\sigma}|^2. \quad (3.20)$$




Figure 2: Case 3: A front hitting the boundary at $t = t_0$.

Proof. Since the front tracking solution satisfies the boundary condition at $t = t_0+$, it will generate the following Riemann problem from the left boundary

$$u(t, 0) = \begin{cases} Ku_L & \text{for } t = t_0+, \\ Ku_R & \text{for } t = t_0-. \end{cases}$$

Let us introduce the map

$$g(\sigma, \sigma_1, \sigma_2) = K\Psi_k(\sigma, u_L) - \Psi_2(\sigma_2, \Psi_1(\sigma_1, Ku_L)),$$

defined for $\sigma, \sigma_1, \sigma_2$ in a neighborhood of 0. Thanks to the C^2 regularity of Lax's curves, this map is C^2 . Besides,

$$\partial_{\sigma_1} g(0, 0, 0) = -r_1(Ku_L), \quad \partial_{\sigma_2} g(0, 0, 0) = -r_2(Ku_L).$$

As the eigenvectors $r_1(Ku_L)$ and $r_2(Ku_L)$ are linearly independent, by the implicit function theorem there exists a C^2 function Φ such that in a neighborhood of 0,

$$g(\sigma, \sigma_1, \sigma_2) = 0 \quad \text{if and only if} \quad (\sigma_1, \sigma_2) = \Phi(\sigma).$$

Close to $\sigma = 0$, this identity can be written as

$$K\Psi_k(\sigma, u_L) - \Psi_2(\Phi_2(\sigma), \Psi_1(\Phi_1(\sigma), Ku_L)) = 0.$$

Doing the Taylor-Young expansion close to $\sigma = 0$, we get the estimate

$$|\sigma Kr_k(u_L) - \sigma\Phi'_2(0)r_2(Ku_L) - \sigma\Phi'_1(0)r_1(Ku_L)| \leq C_b|\sigma|^2,$$

so that we immediately get

$$\Phi'_1(0) = \ell_1(Ku_L) \cdot Kr_k(u_L), \quad \Phi'_2(0) = \ell_2(Ku_L) \cdot Kr_k(u_L).$$

As $\Phi(0) = 0$, we deduce (3.20) for $\hat{\sigma}$ small enough. \square

Now according to (3.6), in the situation of Lemma 3.4, provided

$$C_b V(t_0-) e^{\gamma L} \leq \varepsilon, \quad (3.21)$$

we obtain

$$|\sigma_1| + |\sigma_2| \leq |\hat{\sigma}|(e^{-\gamma L} - \varepsilon), \quad (3.22)$$

so that

$$V(t_0+) - V(t_0-) \leq -\varepsilon|\hat{\sigma}|.$$

Besides, one easily gets that

$$Q(t_0+) - Q(t_0-) \leq V(t_0-)(|\sigma_1| + |\sigma_2|) \leq V(t_0-)|\hat{\sigma}|.$$

It follows that

$$(V(t_0+) + c_0 Q(t_0+)) - (V(t_0-) + c_0 Q(t_0-)) \leq |\hat{\sigma}|(-\varepsilon + c_0 V(t_0-)). \quad (3.23)$$

2. Decay of the functional $J(\mathbf{t}) = \mathbf{V}(\mathbf{t}) + c_0 \mathbf{Q}(\mathbf{t})$.

Let us now prove that if we assume

$$J(0) \leq \min \left\{ e^{-\gamma L}, \frac{1}{2C_\delta e^{2\gamma L}}, \frac{\varepsilon}{C_b e^{\gamma L}}, \frac{\varepsilon}{c_0} \right\}, \quad (3.24)$$

then the functional J is exponentially decaying on $[0, T_h^*)$.

Indeed, suppose that (3.24) is satisfied. We denote by $t_1 = 0$ and $\{t_k\}_{k=1}^\infty$ the increasing sequence of times such that at $t = t_k$ ($k \geq 2$), two fronts interact or some fronts hits the right boundary. We have proved the following:

- from Case 1 and estimate (3.12): for all $t \in (t_k, t_{k+1})$, one has $J(t) \leq e^{-c_* \gamma (t-t_k)} J(t_k)$;
- if two fronts interact at some time t_k corresponding to Case 2, then from estimates (3.17), $J(t_{k+}) \leq J(t_k-)$, provided (3.15) holds, i.e. $\max\{e^{\gamma L}, 2C_\delta e^{2\gamma L}\} V(t_k-) \leq 1$;
- if one front hits the boundary, from Case 3 and estimate (3.23), then $J(t_{k+}) \leq J(t_k-)$ provided $C_b e^{\gamma L} V(t_k) \leq \varepsilon$ and $c_0 V(t_k-) \leq \varepsilon$.

Therefore, if $J(0)$ satisfies (3.24), which can be ensured by choosing $V(0)$, i.e. ε_0 in (3.9), small enough since

$$J(0) \leq V(0) + c_0 V(0)^2,$$

then J decreases and $V = V(t)$ always satisfies

$$V(t) \leq \min \left\{ e^{-\gamma L}, \frac{1}{2C_\delta e^{2\gamma L}}, \frac{\varepsilon}{C_b e^{\gamma L}}, \frac{\varepsilon}{c_0} \right\}. \quad (3.25)$$

Combining the above estimates, we thus obtain

$$J(t) \leq e^{-c_* \gamma t} J(0), \quad \text{for all } t \in (0, T_h^*). \quad (3.26)$$

This concludes the proof of Lemma 3.2. \square

3.4 Global definiteness of the approximations

In this subsection, we prove the following.

Lemma 3.5. *There exists $\varepsilon_1 \in (0, \varepsilon_0]$, where ε_0 is the constant introduced in Lemma 3.2, such that for small $h > 0$, the front-tracking approximation u_h starting from an initial data $u_{0,h}$ satisfying*

$$V(u_{0,h}) \leq \varepsilon_1, \quad (3.27)$$

is defined globally in time, that is, $T_h^ = \infty$.*

Proof. We argue similarly as in [5, Section 7.3, item 4].

First, we notice that $TV^*(\cdot)$ in (2.2) and V are equivalent in the sense that for some $C > 0$,

$$C^{-1}V(U) \leq TV^*(U) \leq CV(U), \quad (3.28)$$

for all piecewise constant functions U on $[0, L]$ with suitably small BV norm (Let us recall that here V takes into account the strength of the fronts emanating from $x = 0$ so that it is not only a semi-norm but a norm under assumption (2.1)). From the estimate (2.3), it is clear that if $u_{0,h}$ satisfies (3.27) for some $\varepsilon_1 \in (0, \varepsilon_0]$ small enough, $\|u_{0,h}\|_{L^\infty(0,L)} \leq \delta_0$, so that $T_h^* > 0$. Then, using the decay estimates (3.11) and estimates (2.3) and (3.28), taking $\varepsilon_1 \in (0, \varepsilon_0]$ smaller if necessary, $J(0)$ can be made small enough to obtain $\|u_h(t)\|_{L^\infty(0,L)} \leq \delta_0/2$ for all $t \in (0, T_h^*)$.

Therefore, the only reason for T_h^* to be finite is the possible accumulation of interaction times. Let us show that this cannot happen. Set

$$M > \max_{|u| \leq \delta} \lambda_2(u), \quad (3.29)$$

and $T_M = L/M$, and define the domain

$$D(t_0) = \{(t, x) \in [0, t_0 + T_M] \times [0, L], \text{ with } M(t - t_0) \leq x\}.$$

Let us first show that in $D(0)$, there can be only finitely many occurrences of interactions from which more than two fronts exit. Indeed, the only fronts interacting in $D(0)$ are the ones coming from $t = 0$, $x \in (0, L)$ and their descendants. By that, we mean that tracing a front from this zone backward in time across interactions, we end up always at a point in $(0, L)$ at $t = 0$, not at the boundary.

Let us then consider the following functionals defined on piecewise constant functions U with suitably small BV norm:

$$\tilde{V}(U, X) = \sum_{\substack{i \in \{0, \dots, n\} \\ x_i \geq X}} (|\sigma_{i,1}| + |\sigma_{i,2}|), \quad (3.30)$$

$$\tilde{Q}(U, X) = \sum_{\substack{(x_i, \sigma_i) \\ x_i \geq X}} \left(|\sigma_i| \sum_{\substack{(x_j, \sigma_j) \text{ approaching } (x_i, \sigma_i) \\ \text{and } x_j \geq X}} |\sigma_j| \right). \quad (3.31)$$

First, similar estimates as before show that, taking C_δ as in Lemma 3.3, for all times $t \in (0, \min\{T_h^*, T_M\})$, the function $t \mapsto \tilde{V}(u_h(t), Mt) + 4C_\delta \tilde{Q}(u_h(t), Mt)$ is non-negative, non-increasing and bounded by $\tilde{V}(u_h(0), 0) + 4C_\delta \tilde{Q}(u_h(0), 0) \leq e^{2\gamma L}(V(0) + 4C_\delta Q(0))$. The only difference with the estimates obtained in Section 3.3 is that now, fronts can moreover leave the domain $D(0)$, in which case the decay of $t \mapsto \tilde{V}(u_h(t), Mt) + 4C_\delta \tilde{Q}(u_h(t), Mt)$ is obvious.

When more than two fronts exit from an interaction point in the domain $D(0) \cap \{t \in [0, T_h^*]\}$, we necessarily are in the situation of two interacting fronts of the same family and, according to Lemma 3.3, their strengths $\tilde{\sigma}$ and $\hat{\sigma}$ satisfy $|\tilde{\sigma}||\hat{\sigma}| \geq h/C_\delta$ (for small h). Indeed, according to the construction of our front tracking solution u_h (recall Section 3.2), this occurs only when two fronts of the same family interact, and the outgoing rarefaction wave of the other family is larger than h . According to (3.16), this can occur only if the product of the two incoming fronts $\tilde{\sigma}, \hat{\sigma}$ satisfies $C_\delta|\tilde{\sigma}||\hat{\sigma}| \geq h$ (recall that in our case $|\sigma| + |\hat{\sigma}|$ can be assumed to be always smaller than 1). But arguing as in the proof of (3.17), this induces a decay of the functional $\tilde{V} + 4C_\delta\tilde{Q}$ of $-h$ at least. Therefore, there can be only a finite number of interactions from which more than two fronts exit. Consequently, the number of fronts is finite in $D(0) \cap \{t \leq T_h^*\}$. This means that the construction cannot blow up in $D(0)$, and thus there are only a finite number of fronts in $D(0)$.

Accordingly, the number of fronts hitting the boundary $x = L$ during the time interval $[0, T_M]$ is finite. We now show that this implies that the number of newly created fronts in the set $[0, T_M] \times [0, L]$ is finite. Indeed, the functional $t \mapsto \tilde{J}(t) = \tilde{V}(u_h(t), 0) + 4C_\delta\tilde{Q}(u_h(t), 0)$ satisfies the following properties:

- When no front hits the boundary and fronts do not meet in the domain, the functional \tilde{J} is constant.
- When a front hits the boundary and additional fronts (in finite number) are possibly created, the functional \tilde{J} may increase. As this occurs a finite number of times, the increase of \tilde{J} on $[0, T_M]$ is necessarily bounded.
- When two fronts meet in the domain $(0, L)$, \tilde{J} decreases. Besides, when more than two fronts issue from the interaction, \tilde{J} decays from $-h$.

Consequently, there can be only a finite number of newly created fronts in the time interval $[0, T_M]$. We can then iterate this argument on the time intervals of the form $[jT_M, (j+1)T_M]$ for $j \in \mathbb{N}$, from which we conclude $T_h^* = \infty$. \square

As a corollary of Lemma 3.2, we have the following result.

Corollary 3.6 (Bounds on the total variation). *Under the setting of Theorem 1.2, there exist positive constants $\varepsilon_2 > 0$, $C > 0$ and $\nu > 0$ such that for any $h > 0$, if $u_{0,h}$ is a piecewise constant function satisfying $TV_{[0,L]}^*(u_{0,h}) \leq \varepsilon_2$, if we define the approximate solution $u_h(t)$ of (1.1)–(1.4) using the front tracking approximation explained in Section 3.2, for all $t \geq 0$,*

$$TV_{[0,L]}^*(u_h(t)) \leq Ce^{-\nu t}TV_{[0,L]}^*(u_{0,h}). \quad (3.32)$$

In particular, if we choose $u_{0,h}$ satisfying (3.3), we get the uniform estimate: for all $t \geq 0$,

$$TV_{[0,L]}^*(u_h(t)) \leq Ce^{-\nu t}TV_{[0,L]}^*(u_0). \quad (3.33)$$

Proof. We recall that the quantities $TV_{[0,L]}^*(\cdot)$ and $V(\cdot)$ are equivalent for u_h satisfying $\|u_h\|_{L^\infty(0,L)} \leq \delta$, see (3.28). Therefore, taking $\varepsilon_2 > 0$ small enough to guarantee (3.27), the front tracking approximation u_h is well-defined for all times according to Lemma 3.5, and we have the decay (3.11). This yields

$$V(t) \leq J(t) \leq \exp(-\nu t)J(0) \quad \text{with} \quad J(0) \leq V(0) + c_0V(0)^2 \leq V(0)(1 + c_0),$$

where we assumed in the last identity that $V(0) \leq 1$, which can be done by taking $\varepsilon_2 > 0$ small enough. This obviously implies (3.32). \square

3.5 Size of the rarefaction fronts

In this subsection, the term “rarefaction front” denotes any piecewise C^1 trajectory $t \mapsto x(t)$ corresponding to a rarefaction discontinuity. The goal of this section is to prove that the rarefaction fronts stay of strength $\mathcal{O}(h)$. This is central in the proof of the entropy inequality in the limit $h \rightarrow 0^+$.

Lemma 3.7. *Under the assumption (3.27), there exists $C > 0$ independent of h and of the number of discontinuities in $u_{0,h}$ such that all the rarefaction fronts in the front-tracking approximations u_h satisfy*

$$\sigma \leq Ch. \quad (3.34)$$

Proof. We argue similarly as in [5, Section 7.3, item 5]. We thus consider a rarefaction front $(t, x(t))$ of strength $\sigma(t) > 0$ and of family k traveling in the domain. Let us say that it is created at some time $t_0 \geq 0$ with $\sigma(t_0) \leq h$ and then evolves according to the construction introduced in Section 3.1 and then possibly ends in the domain $(0, L)$ or on the right boundary. Due to the construction, a rarefaction front hitting the boundary can only generate new rarefaction fronts in the domain with a strength smaller than h , and we can therefore limit ourselves to consider rarefaction fronts until they disappear or until the first time they hit the boundary. As both speeds are strictly positive, we note that the time of existence of a rarefaction front is necessarily limited by $t_0 + L/c_*$. We denote this final time of existence of the front $(t, x(t), \sigma(t))$ by $t_0 + t_*$.

Now for such a rarefaction front there are two possibilities of interactions:

- It may interact with some fronts of the same family: in this case, as rarefaction fronts of the same family never interact, it interacts necessarily with a shock of the same family. Then Glimm’s interaction estimate in Lemma 3.3 shows that its strength becomes smaller.
- It can meet fronts of the other family.

We will consequently focus on the second case. Our proof is based on an explicit bound on the maximal increase of the strength of a rarefaction front during a time interval of length L/M . A simple iteration argument implies then (3.34).

We introduce t_1 such that

$$x(t_0) + M(t_1 - t_0) = L,$$

where we recall that M was defined in (3.29). Next we define a quantity similar to V (but following the rarefaction front that we consider) by

$$V_{(x,\sigma)}(t) = \sum_{\substack{(x_i, \sigma_i) \text{ approaching } (x(t), \sigma(t)) \\ x_i > M(t-t_1)_+}} |\sigma_i|,$$

where the front (x_i, σ_i) of family k_i is said to be approaching the front $(x(t), \sigma(t))$ of family k if one of the following cases occurs:

- $k_i = k$ and $\sigma_i < 0$,
- $k_i < k$ and $x_i > x(t)$,
- $k_i > k$ and $x_i < x(t)$.

Concerning the fronts which may pass through the boundary, we introduce

$$V_{(x,\sigma),b}(t) = \sum_{(x_i,\sigma_i) \text{ with } x_i > x(t_0) + M(t-t_0)} |\sigma_i|.$$

Similarly, we introduce $Q_{(x,\sigma)}(t)$ and $Q_{(x,\sigma),b}(t)$

$$Q_{(x,\sigma)}(t) = \sum_{(x_i,\sigma_i) \text{ with } x_i \geq M(t-t_1)_+} |\sigma_i| \left(\sum_{\substack{(x_j,\sigma_j) \text{ approaching } (x_i,\sigma_i) \\ \text{with } x_j \geq M(t-t_1)_+}} |\sigma_j| \right),$$

$$Q_{(x,\sigma),b}(t) = \sum_{\substack{(x_i,\sigma_i) \\ x_i > x(t_0) + M(t-t_0)}} |\sigma_i| \left(\sum_{\substack{(x_j,\sigma_j) \text{ approaching } (x_i,\sigma_i) \\ \text{with } x_j \geq x(t_0) + M(t-t_0)}} |\sigma_j| \right).$$

We now discuss the evolution of the quantities σ , $V_{(x,\sigma)}$, $V_{(x,\sigma),b}$, $Q_{(x,\sigma)}$, $Q_{(x,\sigma),b}$ as time evolves. Obviously, when there are no interactions at time t , all the above quantities decay. We are therefore left to consider the following three cases:

- **Case 1:** At time t , there is an interaction inside $(0, L)$ that do not involve the rarefaction front $(x(t), \sigma(t))$.
- **Case 2:** At time t , there is an interaction (inside $(0, L)$) involving the rarefaction front $(x(t), \sigma(t))$.
- **Case 3:** At time t , a front hits the boundary.

Case 1. At times $t \in [t_0, t_0 + \min\{L/M, t_*\}]$ corresponding to interactions inside $(0, L)$ that do not involve the rarefaction front $(t, x(t), \sigma(t))$, as in [5, Section 7.3, item 5], we obtain from Lemma 3.3 that

$$\begin{cases} \sigma(t+) = \sigma(t-), \\ V_{(x,\sigma)}(t+) + c_0 Q_{(x,\sigma)}(t+) \leq V_{(x,\sigma)}(t-) + c_0 Q_{(x,\sigma)}(t-), \\ V_{(x,\sigma),b}(t+) + c_0 Q_{(x,\sigma),b}(t+) \leq V_{(x,\sigma),b}(t-) + c_0 Q_{(x,\sigma),b}(t-). \end{cases} \quad (3.35)$$

Case 2. At times $t \in [t_0, t_0 + \min\{L/M, t_*\}]$ corresponding to an interaction inside $(0, L)$ involving the rarefaction front $(t, x(t), \sigma(t))$ with a front of strength σ_α of another family, from Lemma 3.3,

$$\begin{cases} \sigma(t+) \leq \sigma(t-) + C_\delta |\sigma(t-)| |\sigma_\alpha|, \\ V_{(x,\sigma)}(t+) \leq V_{(x,\sigma)}(t-) - |\sigma_\alpha|, \\ Q_{(x,\sigma)}(t+) \leq Q_{(x,\sigma)}(t-), \\ V_{(x,\sigma),b}(t+) + c_0 Q_{(x,\sigma),b}(t+) \leq V_{(x,\sigma),b}(t-) + c_0 Q_{(x,\sigma),b}(t-). \end{cases} \quad (3.36)$$

When the rarefaction front $(t, x(t), \sigma(t))$ interacts with a front σ_α of the same family, then it necessarily corresponds to a shock and therefore $\sigma_\alpha < 0$. Using then Lemma 3.3 and the fact that $\sigma(t-) \leq e^{\gamma L} V(t-)$ is smaller than $1/C_\delta$, recall (3.25), one easily shows that

$$\begin{cases} \sigma(t+) \leq \sigma(t-) - |\sigma_\alpha| + C_\delta |\sigma(t-)| |\sigma_\alpha| \leq \sigma(t-), \\ V_{(x,\sigma)}(t+) + c_0 Q_{(x,\sigma)}(t+) \leq V_{(x,\sigma)}(t-) + c_0 Q_{(x,\sigma)}(t-), \\ V_{(x,\sigma),b}(t+) + c_0 Q_{(x,\sigma),b}(t+) \leq V_{(x,\sigma),b}(t-) + c_0 Q_{(x,\sigma),b}(t-). \end{cases} \quad (3.37)$$

Case 3. At times $t \in [t_0, t_0 + \min\{L/M, t_*\}]$ corresponding to a front hitting the boundary of strength $\hat{\sigma}$, from Lemma 3.4 (recall that the smallness condition (3.21) holds),

$$\begin{cases} \sigma(t+) = \sigma(t-), \\ V_{(x,\sigma)}(t+) + c_0 Q_{(x,\sigma)}(t+) \leq V_{(x,\sigma)}(t-) + c_0 Q_{(x,\sigma)}(t-) + (1 - \varepsilon)|\hat{\sigma}|, \\ V_{(x,\sigma),b}(t+) + c_0 Q_{(x,\sigma),b}(t+) \leq V_{(x,\sigma),b}(t-) + c_0 Q_{(x,\sigma),b}(t-) - |\hat{\sigma}|. \end{cases} \quad (3.38)$$

In particular, relying on the estimates (3.35)–(3.38), there exists some constant C large enough, depending only on C_δ in (3.36), such that

$$t \mapsto \sigma(t) \exp\left(C(V_{(x,\sigma)}(t) + c_0 Q_{(x,\sigma)}(t) + V_{(x,\sigma),b}(t) + c_0 Q_{(x,\sigma),b}(t))\right),$$

is non-increasing on $[t_0, t_0 + \min\{L/M, t_*\}]$. Introducing J defined in (3.10) and \tilde{V}, \tilde{Q} defined in (3.30)–(3.31), we immediately get that for all $t \in [t_0, t_0 + \min\{L/M, t_*\}]$,

$$V_{(x,\sigma)}(t) + c_0 Q_{(x,\sigma)}(t) + V_{(x,\sigma),b}(t) + c_0 Q_{(x,\sigma),b}(t) \leq 2\tilde{V}(u_h(t), 0) + 2c_0\tilde{Q}(u_h(t), 0) \leq 2e^{2\gamma L}J(t).$$

Therefore, the decay property (3.11) yields, for all $t \in [t_0, t_0 + \min\{L/M, t_*\}]$,

$$\sigma(t) \leq \sigma(t_0) \exp(CJ(0)).$$

Of course, this estimate on the maximal amplification of $\sigma(t)$ stays valid in each interval of time of length L/M in the time interval $[t_0, t_0 + t^*]$. As t_* is necessarily bounded by L/c_* , iterating this estimate at worst a finite number of times $\simeq M/c_*$, we get that for all t during which the rarefaction exists, $\sigma(t)$ always satisfies

$$\sigma(t) \leq \sigma(t_0) \exp\left(\frac{CM}{c_*}(V(0) + c_0(V(0))^2)\right) \leq Ch,$$

i.e. the last estimate (3.34) in Lemma 3.3. \square

3.6 Lipschitz in time estimates

In order to pass to the limit in the boundary conditions (1.4), we will also estimate the norm of the approximations in $W^{1,\infty}(0, L; L^1_{loc}(0, \infty))$. Our strategy is inspired by sideways energy estimates which are made possible by the fact that the velocities are bounded from below. These types of energy estimates consist in exchanging the role played by the time and space variables.

The main result of this section is the following one.

Lemma 3.8. *Under the assumptions of Lemma 3.2, there exists $\varepsilon_3 \in (0, \varepsilon_2)$ such that if*

$$TV_{[0,L]}^*(u_{0,h}) \leq \varepsilon_3, \quad (3.39)$$

the approximate solution u_h constructed in Section 3.1 satisfies the following property: for all $T > 0$, there exists C_T with

$$\sup_{x \in [0,L]} TV(u_h(\cdot, x)|_{t \in (0,T)}) \leq C_T. \quad (3.40)$$

Proof. Our main argument here consists in considering the front-tracking approximations u_h with x as a time variable, which is made simple by the fact that the velocities never vanish, recall (1.3). We thus introduce the functionals

$$V_\perp(x, T) = \sum_{\text{fronts } (x,t,\sigma) \text{ with } t < T} |\sigma|, \quad (3.41)$$

and

$$Q_{\perp}(x, T) = \sum_{\text{fronts } (x, t_i, \sigma_i) \text{ with } t_i < T} |\sigma_i| \left(\sum_{\substack{\text{fronts } (x, t_j, \sigma_j) \text{ with } t_j < T \\ (x, t_j, \sigma_j) \text{ approaching } (x, t_i, \sigma_i)}} |\sigma_j| \right), \quad (3.42)$$

where here a front at (x, t_j, σ_j) with strength σ_j and from family k_j is said to be approaching the front at (x, t_i, σ_i) with strength σ_i and from family k_i if

- $t_j < t_i$, $k_i = k_j$ and σ_i or σ_j is negative,
- $t_j < t_i$ and $(k_i, k_j) = (1, 2)$.

Using the functionals in (3.30)–(3.31), one can check that the map

$$x \mapsto \tilde{V}(u_h(x/M), x) + c_0 \tilde{Q}(u_h(x/M), x) + V_{\perp}(L, x/M)$$

decays. This implies in particular that

$$V_{\perp}(L, L/M) \leq \tilde{V}(u_{0,h}, 0) + c_0 \tilde{Q}(u_{0,h}, 0) \leq e^{2\gamma L} J(0).$$

Thanks to the conditions (2.1),

$$V_{\perp}(0, L/M) \leq V_{\perp}(L, L/M) \leq e^{2\gamma L} J(0). \quad (3.43)$$

Now, let us consider the functional

$$J_{\perp}(x) = V_{\perp}(x, L/M) + c_0 Q_{\perp}(x, L/M) + 2\tilde{V}(u_{0,h}, x), \quad (3.44)$$

where the functional \tilde{V} has been defined in (3.30).

With this choice, the same estimates as before show that J_{\perp} decays as x increases provided $J_{\perp}(0)$ is small enough. Indeed, according to Glimm's interaction estimate in Lemma 3.3, arguing as before, if two fronts meet in $(0, L/M) \times \{x\}$, this functional decays provided $V_{\perp}(x, L/M)$ is small enough. When a front appears at time $t = 0$ in the point x , the increase of $V_{\perp}(x, L/M) + c_0 Q_{\perp}(x, L/M)$ is compensated by the decrease of $\tilde{V}(u_{0,h}, x)$. Therefore, while $V_{\perp}(x, L/M)$ is small enough, J_{\perp} decays. Thus, using $V_{\perp}(x, L/M) \leq J_{\perp}(x)$ for all $x \in (0, 1)$, one only needs to guarantee that $J_{\perp}(0)$ is small enough to guarantee the decay for all $x \in (0, L)$. Using (3.43), we deduce that there exists $\varepsilon_{3,0} > 0$ such that if

$$J(0) \leq \varepsilon_{3,0}, \quad (3.45)$$

the functional J_{\perp} decays on $(0, L)$ and

$$\sup_x J_{\perp}(x) \leq C J(0).$$

Now, we obviously have that the semi-norm $TV(u_h(\cdot, x)|_{t \in (0, L/M)})$ is bounded up to a constant by $J_{\perp}(x)$, so that we obtain

$$\sup_{x \in [0, L]} TV(u_h(\cdot, x)|_{t \in (0, L/M)}) \leq C J(0).$$

Obviously, we can perform this argument on each time interval starting at time t_0 , so we can get the following: if

$$J(t_0) \leq \varepsilon_{3,0}, \quad (3.46)$$

one has the following estimate:

$$\sup_{x \in [0, L]} TV(u_h(\cdot, x)|_{t \in (t_0, t_0 + L/M)}) \leq CJ(t_0). \quad (3.47)$$

Using that J is a decaying function of time, recall Lemma 3.2, condition (3.46) is automatically satisfied once condition (3.45) holds. Therefore, provided condition (3.45), summing up the estimates (3.47) for $t_0 \in \frac{L}{2M}\mathbb{N}$, for all $T > 0$, we get the existence of a constant C_T such that (3.40) holds. The proof of Lemma 3.8 is then completed once we have remarked that Condition (3.45) can be enforced by choosing ε_3 small enough in (3.39). \square

3.7 Passing to the limit: end of the proof of Theorem 1.2

Let $u_0 \in BV(0, L)$, and consider a sequence of approximations $u_{0,h}$ converging to u_0 strongly in $L^1(0, L)$ and a.e. in $(0, L)$ as $h \rightarrow 0$ with $TV_{[0, L]}^*(u_{0,h}) \leq 2TV_{[0, L]}^*(u_0)$ such that $u_{0,h}$ is piecewise constant, and consider the corresponding sequence u_h constructed in Section 3.1.

We assume that $\epsilon_0 > 0$ in (1.13) coincides with ε_3 in Lemma 3.8, so that Corollary 3.6 and Lemma 3.8 apply.

Corollary 3.6 shows that u_h is bounded in $L^\infty(0, \infty; BV(0, L))$. Besides, it is easy to check that, due to the finite speed of propagation, there exists a constant $C > 0$ such that for all $t_1 > t_0 > 0$,

$$\|u_h(t_1) - u_h(t_0)\|_{L^1(0, L)} \leq C|t_1 - t_0| \max_{t \in [t_0, t_1]} TV_{[0, L]}^*(u_h(t)) \leq C|t_1 - t_0| TV_{[0, L]}^*(u_0). \quad (3.48)$$

Therefore, one can use Helly's theorem and a diagonal extraction argument (see [5, Theorem 2.4]) and obtain a limit function $u \in L^1_{loc}(0, \infty; BV(0, L))$ such that, up to a subsequence still denoted in the same way for simplicity, u_h strongly converges as $h \rightarrow 0$ to u in $L^1_{loc}((0, \infty) \times (0, L))$. Besides, $u \in L^\infty(0, \infty; BV(0, L)) \cap W^{1, \infty}(0, \infty; L^1(0, L))$ and satisfies

$$\|u(t_1) - u(t_0)\|_{L^1(0, L)} \leq C|t_1 - t_0| TV_{[0, L]}^*(u_0), \quad t_1 > t_0 > 0.$$

In particular, since $u_h(0) = u_{0,h}$ strongly converges to u_0 in $L^1(0, L)$, we immediately get $u(0) = u_0$.

Furthermore, using the semi-continuity of the $TV_{[0, L]}^*$ norm and passing to the limit in (3.32), we obtain, for all $t \geq 0$,

$$TV_{[0, L]}^*(u(t)) \leq Ce^{-\nu t} TV_{[0, L]}^*(u_0), \quad (3.49)$$

which proves (1.14).

To derive that u necessarily is a weak entropy solution of (1.1) in $(0, \infty) \times (0, L)$, we argue as in [5, Section 7.4]. That step mainly consists in measuring the errors done in approximating rarefaction waves by rarefaction fronts. This relies on the estimate (3.34) on the size of the rarefaction fronts. Details of the proof are left to the reader.

It remains to prove that the boundary conditions (1.4) are satisfied. According to Lemma 3.8, for all $T > 0$, the approximate solutions u_h are uniformly bounded in the space $L^\infty(0, L; BV(0, T))$. As the velocities are strictly positive, it follows, similarly as in (3.48), that for all $T > 0$, u_h are uniformly bounded in $W^{1, \infty}(0, L; L^1(0, T))$. Accordingly, using again [5, Theorem 2.4], u satisfies the boundary conditions (1.4) almost everywhere.

References

- [1] Debora Amadori. Initial-boundary value problems for nonlinear systems of conservation laws, *NoDEA Nonlinear Differential Equations Appl.* 4 (1):1–42, 1997.
- [2] Debora Amadori and Rinaldo M. Colombo. Continuous dependence for 2×2 conservation laws with boundary. *J. Differential Equations* 138:229–266, 1997.
- [3] Debora Amadori and Rinaldo M. Colombo. Viscosity solutions and Standard Riemann Semigroup for conservation laws with boundary. *Rend. Sem. Mat. Univ. Padova* 99:219–245, 1998.
- [4] Fabio Ancona and Andrea Marson. Asymptotic stabilization of systems of conservation laws by controls acting at a single boundary point. *Control methods in PDE-dynamical systems*, Contemp. Math. 426, 1–43, Amer. Math. Soc., Providence, RI, 2007.
- [5] Alberto Bressan. *Hyperbolic systems of conservation laws*, volume 20 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2000. The one-dimensional Cauchy problem.
- [6] Alberto Bressan and Giuseppe Maria Coclite. On the boundary control of systems of conservation laws. *SIAM J. Control Optim.* 41 (2):607–622, 2002.
- [7] Rinaldo M. Colombo and Graziano Guerra. On General Balance Laws with Boundary. *J. Differential Equations* 248 (5):1017–1043, 2010.
- [8] Jean-Michel Coron and Georges Bastin. Dissipative boundary conditions for one-dimensional quasi-linear hyperbolic systems: Lyapunov stability for the C^1 -norm. *SIAM J. Control Optim.*, 53(3):1464–1483, 2015.
- [9] Jean-Michel Coron, Georges Bastin, and Brigitte d’Andréa Novel. Dissipative boundary conditions for one-dimensional nonlinear hyperbolic systems. *SIAM J. Control Optim.*, 47(3):1460–1498, 2008.
- [10] Jean-Michel Coron, Brigitte d’Andréa Novel, and Georges Bastin. A strict Lyapunov function for boundary control of hyperbolic systems of conservation laws. *IEEE Trans. Automat. Control*, 52(1):2–11, 2007.
- [11] Jean-Michel Coron and Hoai-Minh Nguyen. Dissipative boundary conditions for nonlinear 1-D hyperbolic systems: sharp conditions through an approach via time-delay systems. *SIAM J. Math. Anal.*, 47(3):2220–2240, 2015.
- [12] Constantine M. Dafermos. *Hyperbolic conservation laws in continuum physics*. Grundlehren Math. Wissenschaften Series, Vol. 325, Springer Verlag, Third Edition, 2010.
- [13] Constantine M. Dafermos. Polygonal approximations of solutions of the initial value problem for a conservation law. *J. Math. Anal. Appl.* 38:33–41, 1972.
- [14] Ronald J. DiPerna. Global existence of solutions to nonlinear hyperbolic systems of conservation laws. *J. Differential Equations*, 20(1):187–212, 1976.
- [15] Carlotta Donadello and Andrea Marson. Stability of front tracking solutions to the initial and boundary value problem for systems of conservation laws. *NoDEA Nonlinear Differential Equations Appl.* 14:569–592, 2007.

- [16] James Glimm. Solutions in the large for nonlinear hyperbolic systems of equations. *Comm. Pure Appl. Math.*, 18:697–715, 1965.
- [17] Jack K. Hale and Sjoerd M. Verduyn Lunel. *Introduction to functional-differential equations*, volume 99 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1993.
- [18] Jonathan de Halleux, Christophe Prieur, Jean-Michel Coron, Brigitte d’Andrea Novel, and Georges Bastin. Boundary feedback control in networks of open channels. *Automatica J. IFAC*, 39(8):1365–1376, 2003.
- [19] Peter D. Lax. Hyperbolic Systems of Conservation Laws. *Comm. Pure Appl. Math.* 10:537–566, 1957.
- [20] Ta-Tsien Li. *Global classical solutions for quasilinear hyperbolic systems*, volume 32 of *RAM: Research in Applied Mathematics*. Masson, Paris, 1994.
- [21] Ta-Tsien Li and Bopeng Rao. Exact boundary controllability for quasi-linear hyperbolic systems. *SIAM J. Control Optim.*, 41(6):1748–1755 (electronic), 2003.
- [22] Vincent Perrollaz. Asymptotic stabilization of entropy solutions to scalar conservation laws through a stationary feedback law. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 30(5):879–915, 2013.
- [23] Tie Hu Qin. Global smooth solutions of dissipative boundary value problems for first order quasilinear hyperbolic systems. *Chinese Ann. Math. Ser. B*, 6(3):289–298, 1985. A Chinese summary appears in *Chinese Ann. Math. Ser. A* **6** (1985), no. 4, 514.
- [24] Monique Sablé-Tougeron. Méthode de Glimm et problème mixte. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(4):423–443, 1993.
- [25] Yan Chun Zhao. Classical solutions for quasilinear hyperbolic systems. *Thesis, Fudan University*, 1986. In Chinese.