

ON THE INSTABILITY OF EIGENVALUES

SYLVAIN GOLÉLIA

ABSTRACT. This is the proceeding of a talk given in Workshop on Differential Geometry and its applications at Alexandru Ioan Cuza University Iași, Romania, September 2–4, 2009. I explain how positive commutator estimates help in the analysis of embedded eigenvalues in a geometrical setting. Then, I will discuss the disappearance of eigenvalues in the perturbation theory and its relation with the Fermi golden rule.

1. INTRODUCTION

Let $\mathbb{H} := \{(x, y) \in \mathbb{R}^2, y > 0\}$ be the Poincaré half-plane and we endow it with the metric $g := y^{-2}(dx^2 + dy^2)$. Consider the group $\Gamma := PSL_2(\mathbb{Z})$. It acts faithfully on \mathbb{H} by homographies, from the left. The interior of a fundamental domain of the quotient \mathbb{H}/Γ is given by $X := \{(x, y) \in \mathbb{H}, |x| < 1, x^2 + y^2 > 1\}$. Let $\mathcal{H} := L^2(X, g)$ be the set of L^2 integrable function acting on X , with respect to the volume element $dx dy/y^2$. Let $\mathcal{C}_b^\infty(X)$ be the restriction to X of the smooth bounded functions acting on \mathbb{H} which are \mathbb{C} -valued and invariant under Γ . The (non-negative) Laplace operator is defined as the closure of

$$\Delta := -y^2(\partial_x^2 + \partial_y^2), \text{ on } \mathcal{C}_b^\infty(X).$$

It is a (unbounded) self-adjoint operator on $L^2(X)$. Using Eisenstein series, for instance, one sees that its essential spectrum is given by $[1/4, \infty)$ and that it has no singularly continuous spectrum, with respect to the Lebesgue measure. It is well-known that Δ has infinitely many eigenvalues accumulating at $+\infty$ and that every eigenspace is of finite dimension. We refer to [5] for an introduction to the subject.

We consider the Schrödinger operator $H_\lambda := \Delta + \lambda V$, where V is the multiplication by a bounded, real-valued function and $k \in \mathbb{R}$. We focus on an eigenvalue $k > 1/4$ of Δ and assume that the following hypothesis of *Fermi golden rule* holds true. Namely, there is $c_0 > 0$ so that:

$$(1.1) \quad \lim_{\varepsilon \rightarrow 0^+} PV\bar{P} \operatorname{Im}(H_0 - k + i\varepsilon)^{-1} \bar{P}VP \geq c_0 P,$$

in the form sense and where $P := P_k$, the projection on the eigenspace of k , and $\bar{P} := 1 - P$. As P is of finite dimension, the limit can be taken in the weak or in the strong sense. At least formally, $\bar{P} \operatorname{Im}(H_0 - k + i\varepsilon)^{-1} \bar{P}$ tends to the Dirac mass $\pi \delta_k(\bar{P}H_0)$. Therefore, the potential V couples the eigenspace of k and $\bar{P}H_0$ over k in a non-trivial way. This is a key assumption in the second-order perturbation theory of embedded eigenvalues, e.g., [13], and all the art is to prove that it implies there is $\lambda_0 > 0$ that H_λ has no eigenvalue in a neighborhood of k for $\lambda \in (0, |\lambda_0|)$.

In [4], one shows that generically the eigenvalues disappear under the perturbation of a potential (or of the metric) on a compact set. In this note, we are interested about the optimal decay at infinity of the perturbation given by a potential. Using the general result obtained in [3] and under a hypothesis of Fermi golden rule, one is only able to cover the assumption $VL^3 = o(1)$, as $y \rightarrow +\infty$, where L denotes the operator of multiplication by $L := (x, y) \mapsto 1 + \ln(y)$. We give the main result:

Theorem 1.1. *Let $k > 1/4$ be an L^2 -eigenvalue of Δ . Suppose that $VL = o(1)$, as $y \rightarrow +\infty$ and that the Fermi golden rule (1.1) holds true, then there is $\lambda_0 > 0$, so that H_λ has no eigenvalue in a neighborhood of k , for all $\lambda \in (0, |\lambda_0|)$. Moreover, if $VL^{1+\varepsilon} = o(1)$, as $y \rightarrow +\infty$ for some $\varepsilon > 0$, then H_λ has no singularly continuous spectrum.*

We believe that the hypothesis $VL = o(1)$ is optimal in the scale of L . In our approach, we use the Mourre theory, see [1, 12] and establish a positive commutator estimate.

2. IDEA OF THE PROOF

Standardly, for y large enough and up to some isometry \mathcal{U} , see for instance [6, 9, 10] the Laplace operator can be written as

$$(2.2) \quad \tilde{\Delta} = (-\partial_r^2 + 1/4) \otimes P_0 + \tilde{\Delta}(1 \otimes P_0^\perp)$$

on $\mathcal{C}_c^\infty((c, \infty), dr) \otimes \mathcal{C}^\infty(S^1)$, for some $c > 0$ and where P_0 is the projection on constant functions and $P_0^\perp := 1 - P_0$. The Friedrichs extension of the operator $\tilde{\Delta}(1 \otimes P_0^\perp)$ has compact resolvent.

Then, as in [9, 10], we construct a conjugate operator. One chooses $\Phi \in \mathcal{C}_c^\infty(\mathbb{R})$ with $\Phi(x) = x$ on $[-1, 1]$, and sets $\Phi_\Upsilon(x) := \Upsilon\Phi(x/\Upsilon)$, for $\Upsilon \geq 1$. Let $\tilde{\chi}$ be a smooth cut-off function being 1 for r big enough and 0 for r being close to c . We define on $\mathcal{C}_c^\infty((c, \infty) \times S^1)$ a micro-localized version of the generator of dilations:

$$(2.3) \quad S_{\Upsilon,0} := \tilde{\chi} \left((\Phi_\Upsilon(-i\partial_r)r + r\Phi_\Upsilon(-i\partial_r)) \otimes P_0 \right) \tilde{\chi}.$$

The operator $\Phi_\Upsilon(-i\partial_r)$ is defined on the real line by $\mathcal{F}^{-1}\Phi_\Upsilon(\cdot)\mathcal{F}$, where \mathcal{F} is the unitary Fourier transform. We also denote its closure by $S_{\Upsilon,0}$ and it is self-adjoint. In [6] for instance, one does not use a micro-localization and one is not able to deal with really singular perturbation of the metric as in [9, 10].

Now, one obtains

$$[\partial_r^2, \tilde{\chi}(\Phi_\Upsilon r + r\Phi_\Upsilon)\tilde{\chi}] = 4\tilde{\chi}\partial_r\Phi_\Upsilon\tilde{\chi} + \text{remainder}.$$

Using a cut-off function $\tilde{\mu}$ being 1 on the cusp and 0 for $y \leq 2$, we set

$$(2.4) \quad S_\Upsilon := \mathcal{U}^{-1}S_{\Upsilon,0}\mathcal{U}\tilde{\mu}$$

This is self-adjoint in $L^2(X)$. Now by taking Υ big enough, one can show, as in [9, 10] that given an interval \mathcal{J} around k , there exist $\varepsilon_\Upsilon > 0$ and a compact operator K_Υ such that the inequality

$$(2.5) \quad E_{\mathcal{J}}(\Delta)[\Delta, iS_\Upsilon]E_{\mathcal{J}}(\Delta) \geq (4\inf(\mathcal{J}) - \varepsilon_\Upsilon)E_{\mathcal{J}}(\Delta) + E_{\mathcal{J}}(\Delta)K_\Upsilon E_{\mathcal{J}}(\Delta)$$

holds in the sense of forms, and such that ε_Υ tends to 0 as Υ goes to infinity. Here, $E_{\mathcal{J}}(\cdot)$ denotes the spectral measure above the interval \mathcal{J} .

Now, we apply \overline{P} to the left and right of (2.5). Easily one has $\overline{P}E_{\mathcal{J}}(\Delta) = \overline{P}E_{\mathcal{J}}(\Delta\overline{P})$. We get:

$$\begin{aligned} \overline{P}E_{\mathcal{J}}(\overline{P}\Delta)[\overline{P}\Delta, i\overline{P}S_{\Upsilon}\overline{P}]E_{\mathcal{J}}(\overline{P}\Delta)\overline{P} &\geq (4\inf(\mathcal{J}) - \varepsilon_{\Upsilon})\overline{P}E_{\mathcal{J}}(\Delta\overline{P})\overline{P} \\ &\quad + \overline{P}E_{\mathcal{J}}(\overline{P}\Delta)K_{\Upsilon}E_{\mathcal{J}}(\overline{P}\Delta)\overline{P} \end{aligned}$$

One can show that $\overline{P}S_{\Upsilon}\overline{P}$ is self-adjoint in $\overline{P}L^2(X)$ and that $[\overline{P}\Delta, \overline{P}S_{\Upsilon}\overline{P}]$ extends to a bounded operator.

We now shrink the size of the interval \mathcal{J} . As $\overline{P}\Delta$ has no eigenvalue in \mathcal{J} , then the operator $\overline{P}E_{\mathcal{J}}(\overline{P}\Delta)K_{\Upsilon}E_{\mathcal{J}}(\overline{P}\Delta)\overline{P}$ tends to 0 in norm. Therefore, by shrinking enough, one obtains a smaller interval \mathcal{J} containing k and a constant $c > 0$ so that

$$(2.6) \quad \overline{P}E_{\mathcal{J}}(\overline{P}\Delta)[\overline{P}\Delta, i\overline{P}S_{\Upsilon}\overline{P}]E_{\mathcal{J}}(\overline{P}\Delta)\overline{P} \geq c\overline{P}E_{\mathcal{J}}(\Delta\overline{P})\overline{P}$$

holds true in the form sense on $\overline{P}L^2(X)$. At least formally, the positivity on $\overline{P}L^2(X)$ of the commutator $[H_{\lambda}, i\overline{P}S_{\Upsilon}\overline{P}]$, up to some spectral measure and to some small λ , should be a general fact and should not rely on the Fermi golden rule hypothesis.

We now try to extract some positivity on $PL^2(X)$. First, we set

$$(2.7) \quad R_{\varepsilon} := ((H_0 - k)^2 + \varepsilon^2)^{-1/2}, \quad \overline{R}_{\varepsilon} := \overline{P}R_{\varepsilon} \text{ and } F_{\varepsilon} := \overline{R}_{\varepsilon}^2.$$

Note that $\varepsilon R_{\varepsilon}^2 = \text{Im}(H_0 - k + i\varepsilon)^{-1}$ and that R_{ε} commutes with P . Using (1.1), we get:

$$(2.8) \quad (c_1/\varepsilon)P \geq PV\overline{P}F_{\varepsilon}\overline{P}VP \geq (c_2/\varepsilon)P,$$

for $\varepsilon_0 > \varepsilon > 0$.

We follow an idea of [2], which was successfully used in [8, 11] and set

$$B_{\varepsilon} := \text{Im}(\overline{R}_{\varepsilon}^2VP).$$

It is a finite rank operator. Observe now that we gain some positivity as soon as $\lambda \neq 0$:

$$(2.9) \quad P[H_{\lambda}, i\lambda B_{\varepsilon}]P = \lambda^2PVF_{\varepsilon}VP \geq (c_2\lambda^2/\varepsilon)P.$$

It is therefore natural to modify the conjugate operator S_{Υ} to obtain some positivity on $PL^2(X)$. We set

$$(2.10) \quad \hat{S}_{\Upsilon} := \overline{P}S_{\Upsilon}\overline{P} + \lambda\theta B_{\varepsilon}.$$

It is self-adjoint on $\mathcal{D}(S_{\Upsilon})$ and is diagonal with respect to the decomposition $\overline{P}L^2(X) \oplus PL^2(X)$.

Here $\theta > 0$ is a technical parameter. We choose ε and θ , depending on λ , so that $\lambda = o(\varepsilon)$, $\varepsilon = o(\theta)$ and $\theta = o(1)$ as λ tends to 0. We summarize this into:

$$(2.11) \quad |\lambda| \ll \varepsilon \ll \theta \ll 1, \text{ as } \lambda \text{ tends to } 0.$$

With respect to the decomposition $\overline{P}E_{\mathcal{J}}(\Delta) \oplus PE_{\mathcal{J}}(\Delta)$, as λ goes to 0, we have

$$\begin{aligned} E_{\mathcal{J}}(\Delta) [\lambda V, i\overline{P}S_{\Upsilon}\overline{P}] E_{\mathcal{J}}(\Delta) &= \begin{pmatrix} O(\lambda) & O(\lambda) \\ O(\lambda) & 0 \end{pmatrix}, \\ E_{\mathcal{J}}(\Delta) [\Delta, i\lambda\theta B_{\varepsilon}] E_{\mathcal{J}}(\Delta) &= \begin{pmatrix} 0 & O(\lambda\theta\varepsilon^{-1/2}) \\ O(\lambda\theta\varepsilon^{-1/2}) & 0 \end{pmatrix}, \\ \text{and } E_{\mathcal{J}}(\Delta) [\lambda V, i\lambda\theta B_{\varepsilon}] E_{\mathcal{J}}(\Delta) &= \begin{pmatrix} O(\lambda^2\theta\varepsilon^{-3/2}) & O(\lambda^2\theta\varepsilon^{-3/2}) \\ O(\lambda^2\theta\varepsilon^{-3/2}) & \lambda^2\theta F_{\varepsilon} \end{pmatrix}. \end{aligned}$$

Now comes the delicate point. Under the condition (2.11) and by choosing \mathcal{I} , slightly smaller than \mathcal{J} , we use the previous estimates and a Schur Lemma to deduce:

$$(2.12) \quad E_{\mathcal{I}}(H_{\lambda}) [H_{\lambda}, i\hat{S}_{\Upsilon}] E_{\mathcal{I}}(H_{\lambda}) \geq \frac{c\lambda^2\theta}{\varepsilon} E_{\mathcal{I}}(H_{\lambda}),$$

for some positive c and as λ tends to 0.

We mention that only the decay of VL is used to establish the last estimate. In fact, one uses that $[V, i\hat{S}_{\Upsilon}](\Delta + 1)^{-1}$ is a compact operator.

Now it is a standard use of the Mourre theory to deduce Theorem 1.1 and refer to [1], see [9, 10] for some similar application of the theory. For the absence of eigenvalue, one relies on the fact that given an eigenfunction f of H_{λ} w.r.t. an eigenvalue $\kappa \in \mathcal{I}$, one has:

$$(2.13) \quad \langle f, [H_{\lambda}, i\hat{S}_{\Upsilon}]f \rangle = \langle f, [H_{\lambda} - \kappa, i\hat{S}_{\Upsilon}]f \rangle = 0.$$

Then, one applies f on the right and on the left of (2.12) and infers that $f = 0$ thanks to the fact that the constant $c\lambda^2\theta$ is non-zero.

In [9, 10], we prove that the C_0 -group $(e^{iS_{\Upsilon}t})_{t \in \mathbb{R}}$ stabilizes the domain $\mathcal{D}(H_{\lambda}) = \mathcal{D}(\Delta)$. By perturbation, we prove that this is also the case for $(e^{i\hat{S}_{\Upsilon}t})_{t \in \mathbb{R}}$. Thanks to this property, we can expand the commutator of (2.13) in a legal way. This is known as the Virial theorem in the Mourre Theory, see [1, 12].

REFERENCES

- [1] W. Amrein, A. Boutet de Monvel and V. Georgescu, *C₀-Groups, commutator methods and spectral theory of N-body Hamiltonians*, Birkhäuser, Basel-Boston-Berlin, 1996.
- [2] V. Bach, J. Fröhlich, I.M. Sigal and A. Soffer: *Positive commutators and Spectrum of Pauli-Fierz Hamiltonian of Atoms and Molecules*, Commun. Math. Phys. **207** (3), 1999, 557–587.
- [3] L. Cattaneo, G.M. Graf, and W. Hunziker *A general resonance theory based on Mourre's inequality* Ann. Henri Poincaré 7, No. 3, 583–601 (2006).
- [4] Y. Colin de Verdière, *Pseudo-laplaciens II*, Ann. Inst. Fourier 33, No.2, 87–113 (1983).
- [5] J. Fischer, *An approach to the Selberg trace formula via the Selberg zeta-function*, Lecture Notes in Mathematics, **1253**. Springer-Verlag, Berlin, 1987.
- [6] R. Froese and P. D. Hislop, *Spectral analysis of second-order elliptic operators on noncompact manifolds*, Duke Math. J. **58** (1989), No. 1, 103–129.
- [7] V. Georgescu and S. Golénia, *Decay Preserving Operators and stability of the essential spectrum*, J. Oper. Theory. **59** (2008), no. 1, 115–155.
- [8] S. Golénia: *Positive commutators, Fermi golden rule and the spectrum of the zero temperature Pauli-Fierz hamiltonians*. J. of Funct. Anal., Vol. 256, Issue 8, April 2009, 2587–2620.

- [9] S. Golénia and S. Moroianu, *Spectral analysis of magnetic Laplacians on conformally cusp manifolds*, Ann. H. Poincaré **9** (2008), 131–179.
- [10] S. Golénia and S. Moroianu, *The spectrum of Schrödinger operators and Hodge Laplacians on conformally cusp manifolds*. To appear in Trans. A.M.S.
- [11] M. Merlki, *Positive Commutators in Non-Equilibrium Statistical Mechanics*, Comm. Math. Phys. **223**, 327–362 (2001).
- [12] E. Mourre, *Absence of singular continuous spectrum for certain self-adjoint operators*, Comm. Math. Phys. **91** (1981) 391–408.
- [13] M. Reed and B. Simon, *Methods of modern mathematical physics I-IV: Fourier analysis, self-adjointness*, Academic Press, New York – San Francisco – London, 1975.

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UNIVERSITÉ BORDEAUX 1, 351, COURS DE LA LIBRA-
TION, 33405 TALENCE CEDEX, FRANCE

E-mail address: sylvain.golenia@u-bordeaux1.fr