

# Theoretical and numerical study of a quasilinear Zakharov system describing Landau damping

R. Belaouar<sup>1,2</sup>, T. Colin<sup>2</sup>, G. Gallice<sup>1</sup>, C. Galusinski<sup>2</sup>

<sup>1</sup> *SIS, CEA CESTA, BP 2 33114 Le barp, France.*

<sup>2</sup> *Mathématiques Appliquées de Bordeaux, UMR CNRS 5466 et CEA LRC M03,  
Université Bordeaux 1, 351 cours de la Libération, 33405 Talence, France*

## Abstract

In this paper, we study a Zakharov system coupled to an electron diffusion equation describing laser-plasma interactions. Starting from the Vlasov-Maxwell system, and using a transformation that decomposes slowly varying components on the plasma frequency time scale, we derive a nonlinear Schrödinger like system which takes into account the energy exchanged between the plasma waves and the electrons via Landau damping. Two existence theorems are established in a subsonic regime. Then, by using a time-splitting spectral discretizations for the Zakharov system and a finite difference scheme for the electron diffusion equation, we investigate numerical simulations and show how Landau damping works.

## 1 Introduction and physical situation

The interaction of an intense laser pulse with a plasma is a complex physical phenomenon where numerical simulation plays a key role in its understanding. One of the main goal is to simulate nuclear fusion by inertial confinement in a laboratory. We therefore need some accurate and reliable numerical models of laser-plasma interactions. Vlasov or particle-in-cell (PIC) simulations have been used for a more complete description of the problem. However, these kinetic simulations have difficulties in studying weak instabilities and long time behaviors because they need to resolve very small spatial and temporal scales. For the same reasons, it is not possible to use Euler-Maxwell equations. At the beginning of the 70's, Zakharov and its collaborators introduced the so-called Zakharov's equations in order to describe the non-linear interactions between the high-frequency electronic plasma waves and the low-frequency ion-acoustic waves. Basically, the slowly varying envelope of the electric field  $E = \nabla\psi$  is coupled to the low-frequency variation of the density of the ions  $\delta n$  by the following equations written in a dimensionless form [23] :

$$\begin{cases} i\partial_t \nabla\psi + \Delta(\nabla\psi) = \nabla\Delta^{-1}\text{div}(\delta n \nabla\psi), \\ \partial_t^2 \delta n - \Delta\delta n = \Delta(|\nabla\psi|^2). \end{cases} \quad (1.1)$$

Of course, variations of this systems exist (see [20] for example). For laser propagation, one uses the paraxial approximation and the Zakharov system reads

$$\begin{cases} i(\partial_t + \partial_y)E + \Delta_{\perp}E = nE, \\ (\partial_t^2 - \Delta_{\perp})n = \Delta_{\perp}(|E|^2), \end{cases} \quad (1.2)$$

where  $\Delta_{\perp} = \partial_x^2 + \partial_z^2$ . (See [16] for a systematic use of this kind of models for numerical simulation). Concerning the system (1.2), F. Linares, G. Ponce and J.C. Saut (see [13])

have shown that the Cauchy problem is well-posed on  $\mathbb{R}^n$  but T. Colin and G. Metivier (see [3]) have shown that it is ill-posed in the periodic framework.

Recently, M. Colin and T. Colin [4], starting from [17], derived a complete set of quasilinear Zakharov equations describing the interactions between the laser fields, the stimulated Raman and Brillouin processes, the electronic plasma waves and the low-frequency variations of density of the ions. The system involves four Schrödinger equations coupled by quasilinear terms and a wave equation. It reads :

$$i \left( \partial_t + \frac{k_0 c^2}{\omega_0} \partial_y \right) A_0 + \frac{c^2}{2\omega_0} \Delta A_0 - \frac{k_0^2 c^4}{2\omega_0^3} \partial_y^2 A_0 = \frac{\omega_{pe}^2}{2\omega_0} \langle \delta n \rangle (A_0 + e^{-2ik_0 y} A_B) - \frac{e}{2m_e \omega_0} (\nabla \cdot E_0) A_R e^{-i(k_1 y - \omega_1 t)}, \quad (1.3)$$

$$i \left( \partial_t - \frac{k_0 c^2}{\omega_0} \partial_y \right) A_B + \frac{c^2}{2\omega_0} \Delta A_B - \frac{k_0^2 c^4}{2\omega_0^3} \partial_y^2 A_B = \frac{\omega_{pe}^2}{2\omega_0} \langle \delta n \rangle (A_0 + e^{-2ik_0 y} A_B) - \frac{e}{2m_e \omega_0} (\nabla \cdot E_0) A_R e^{i((2k_0 - k_1)y - \omega_1 t)}, \quad (1.4)$$

$$i \left( \partial_t + \frac{k_R c^2}{\omega_R} \partial_y \right) A_R + \frac{c^2}{2\omega_R} \Delta A_R - \frac{k_R^2 c^4}{2\omega_R^3} \partial_y^2 A_R = \frac{\omega_{pe}^2}{2\omega_R} \langle \delta n \rangle A_R - \frac{e}{2m_e \omega_R} (\nabla \cdot E_0^*) (A_0 + e^{-2ik_0 y} A_B) e^{i(k_1 y - \omega_1 t)}, \quad (1.5)$$

$$i \partial_t E_0 + \frac{v_{th}^2}{2\omega_{pe}} \Delta E_0 = \frac{\omega_{pe}}{2} \nabla \Delta^{-1} \text{div} (\langle \delta n \rangle E_0) + \frac{e \omega_{pe}}{2c^2 m_e} \nabla \left( A_R^* (A_0 + e^{-2ik_0 y} A_B) e^{i(k_1 y - \omega_1 t)} \right), \quad (1.6)$$

$$(\partial_t^2 - c_s^2 \Delta) \langle \delta n \rangle = \frac{1}{4\pi n_0 m_i} \Delta (|E_0|^2 + \frac{\omega_{pe}^2}{c^2} (|A_0 + e^{-2ik_0 y} A_B|^2 + |A_R|^2)). \quad (1.7)$$

Here  $A_0$  is the incident laser field,  $A_B$  is the Brillouin component,  $A_R$  is the Raman field,  $E_0$  the electronic-plasma field and  $\langle \delta n \rangle$  the low-frequency variation of the density of the ions. See [4] for the precise definition of all the constants involved in (1.3)-(1.7).

However these various fluid models do not take into account the kinetic effects such that Landau damping (wave-particle process). This process is especially important in the context of fusion by inertial confinement by lasers because electrons are accelerated to high energy and this induces a preheat of the fusion fuel and reduces the target gain. In order to obtain a system describing this wave-particle process we will derive a new set of equations starting from the Vlasov-Maxwell system. This will be done in section 2. The system involves the Zakharov equations coupled with a quasilinear diffusion equations for

the electron distribution function. In one dimension of space, the dimensionless system reads :

$$2i(\partial_t E + \nu \star E) + 3\partial_x^2 E = \delta n E + E_p(x)e^{i(k_1 x - \omega_1 t)}, \quad (1.8)$$

$$\partial_t^2 \delta n - \mu \partial_x^2 \delta n = \frac{\mu}{4} \partial_x^2 (|E|^2), \quad (1.9)$$

$$\widehat{\nu}(t, \xi) = -\frac{\pi}{2\xi|\xi|} \partial_v F_e(t, v = \frac{1}{\xi}), \quad \xi \in \Omega_\xi, \quad (1.10)$$

where

$$\partial_t F_e = \partial_v (D(t, v) \partial_v F_e), \quad D(t, v) = \frac{1}{4|v|} \left| \widehat{E} \left( t, \xi = \frac{1}{v} \right) \right|^2, \quad v \in \Omega_v. \quad (1.11)$$

Here  $E$  is the slowly varying amplitude of the high-frequency electronic plasma waves,  $\delta n$  the low-frequency variation of the density of the ions,  $F_e$  the spatially average electron distribution function and  $\widehat{\nu}$  the spatial fourier transform of  $\nu$  corresponding to the Landau damping rate. This kind of model is valid for bounded velocity that are also bounded away from zero.  $\Omega_v$  is the velocity domain on which equation (1.11) has to be satisfied and  $\Omega_\xi = \{\xi \in \mathbb{R} \text{ s.t. } \exists v \in \Omega_v, v = \frac{1}{\xi}\}$ . The domain  $\Omega_\xi$  will therefore be taken under the form,  $\Omega_\xi = [-A, -a] \cup [a, A]$  with  $0 < a < A$  (see section 2).

Note the term  $\widehat{\nu}$  is only defined on  $\Omega_\xi$  by (1.10) and is extended by 0 outside the domain  $\Omega_\xi$ .

The term  $E_p(x)e^{i(k_1 x - \omega_1 t)}$  is the pump wave. In this work, it is a given function.  $(k_1, \omega_1)$  satisfies the dispersion relation of the Schödinger equation :  $\omega_1 = \frac{3}{2}k_1^2$ . The energy is brought to the system through this term. In fact, in a more complete model, this term will be given by the Raman interaction given by system (1.3)-(1.7) and corresponds to  $\nabla (A_R^*(A_0 + e^{-2ik_0 y} A_B) e^{i(k_1 y - \omega_1 t)})$ . We postponed the study of this completed system to a futur work.

**Notation :** We define the Sobolev space  $H^s(\mathbb{R}^d)$  as follows

$$H^s(\mathbb{R}^d) = \left\{ u \in \mathcal{S}'(\mathbb{R}^d) \mid \|u\|_{H^s(\mathbb{R}^d)}^2 = \int_{\mathbb{R}^d} (1 + |\xi|^2)^s |\widehat{u}(\xi)|^2 d\xi < +\infty \right\}$$

where  $\widehat{u}(\xi) = \mathcal{F}(u)$  is the Fourier transform of  $u$ .

## 1.1 Statements of the results.

The local in time Cauchy problem for the usual Zakharov equations (1.1) is now well understood in the context of regular solutions (see [1], [14], [19] [21] for local models, see [2] for the non-local case (1.1)). For weak solutions, one can see [6]. For finite-time blow-up see [7, 8]. For system (1.3)-(1.6), M. Colin and T. Colin have shown local existence in time for strong solutions [4].

Unfortunately, we are not able at this point to give an existence result for (1.8)-(1.11). We will restrict ourself to the case  $\mu = +\infty$  corresponding to a “subsonic regime”. In this case, introducing  $H_e(t, \xi) = F_e(t, \frac{1}{\xi})$  and devoting by  $\Omega = \Omega_\xi$  the frequency domain,

system (1.8)-(1.11) becomes

$$\begin{aligned}
i(\partial_t E + \nu * E) + \partial_x^2 E &= |E|^2 E + f, \\
\partial_t H_e - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 \partial_\xi H_e) &= 0, \quad \forall \xi \in \Omega, \\
\hat{\nu}(t, \xi) &= \text{sgn}(\xi) \partial_\xi H_e(t, \xi) 1_\Omega, \\
H_e(0, \cdot) &= H_{e0}(\cdot), E(0, \cdot) = E_0(\cdot).
\end{aligned} \tag{1.12}$$

The second equation of system (1.12) has to be endowed with boundary conditions. Since  $\hat{\nu}$  is extended by zero outside of  $\Omega$ , and since  $\hat{\nu}(t, \xi) = \text{sgn}(\xi) \partial_\xi H_e(t, \xi)$  for  $\xi \in \Omega$ , it is natural to impose  $\partial_\xi H_e|_{\partial\Omega} = 0$ .

In order to construct local in time solutions for (1.12), the main problem is to deal with a nonlinear coupling between the electric field, which is a function of the space position, and the electronic distribution, which is function of the frequency. Due to this spatio-frequential coupling as well as the nonlinear terms, we need to simultaneously consider the problem in space and frequency variables for the electric field.

We obtain two kinds of result. The first one concerns the local existence and uniqueness of solutions to (1.12). Moreover, one shows that if  $\hat{\nu}(0, \xi) \geq 0$  then for all  $t$ ,  $\hat{\nu}(t, \xi) \geq 0$ . The term  $\nu * E$  is therefore a damping term. The second result is a global existence result but only in the case where the nonlinear term  $|E|^2 E$  and the source term  $f$  in the right-hand-side of the first equation of (1.12) are replaced by 0. Note that even in this case, the system is far away of being linear!

Let us denote by  $H_n^2(\Omega) = \{g \in H^2(\Omega) \text{ s.t. } \frac{\partial g}{\partial x} = 0 \text{ on } \partial\Omega\}$ .

**Theorem 1.1.** (*Local strong solutions.*) Let  $E_0$  and  $H_{e0}$  such that  $E_0 \in H^1(\mathbb{R})$ ,  $\hat{E}_0 \in H^2(\Omega) \cap H^1(\mathbb{R})$  and  $H_{e0} \in H_n^2(\Omega)$ .

Take  $f \in L^\infty(\mathbb{R}^+; H^1)$  such that  $\hat{f} \in L^\infty(\mathbb{R}^+; H^1(\mathbb{R}) \cap H^2(\Omega))$ .

Then there exists  $T^* > 0$  and a unique solution  $(E, H_e)$  of (1.12) satisfying

$$\begin{aligned}
(E, \hat{E}, H_e) &\in L^\infty([0, T^*]; H^1(\mathbb{R})) \times L^\infty([0, T^*]; H^2(\Omega) \cap H^1(\mathbb{R})) \times L^\infty([0, T^*]; H_n^2(\Omega)), \\
(E, \hat{E}, H_e) &\in C^0([0, T^*]; H^{1-\eta}(\mathbb{R})) \times C^0([0, T^*]; H^{2-\eta}(\Omega) \cap H^{1-\eta}(\mathbb{R})) \times C^0([0, T^*]; H^{2-\eta}(\Omega)), \quad \forall \eta > 0.
\end{aligned}$$

Moreover if  $\hat{\nu}(0, \xi) \geq 0 \quad \forall \xi \in \mathbb{R}$  then  $\forall t \in [0, T^*]$ ,  $\hat{\nu}(t, \xi) \geq 0$ .

**Theorem 1.2.** (*Global solution in the homogeneous case.*) Let  $E_0, H_0$ , such that  $\hat{E}_0 \in H^1(\Omega)$   $H_0 \in H^1(\Omega)$ . One moreover assume that

$$\hat{\nu}_{e0}(\xi) = \text{sgn}(\xi) \partial_\xi H_{e0}(\xi) \geq 0, \quad \forall \xi \in \Omega,$$

and  $(H_0 + \xi^2 \partial_\xi (|\xi|^3 |\hat{E}_0|^2))^+ \in L^\infty(\Omega)$ . Then there exists  $E, H_e$  such that  $\hat{E} \in L_{loc}^\infty(\mathbb{R}_+, H^1)$ ,  $H_e \in L_{loc}^\infty(\mathbb{R}_+, H^1)$  satisfying

$$i(\partial_t E + \nu * E) + \partial_x^2 E = 0, \tag{1.13}$$

$$\begin{aligned}
\hat{\nu}(t, \xi) &= \text{sgn}(\xi) \partial_\xi H_e(t, \xi), \quad \forall \xi \in \Omega, \\
\partial_t H_e - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 \partial_\xi H_e) &= 0, \quad \forall \xi \in \Omega,
\end{aligned} \tag{1.14}$$

$$\begin{aligned}
E(0, x) &= E_0(x), \\
H_e(0, \xi) &= H_0(\xi),
\end{aligned}$$

and

$$\partial_\xi H_e = 0 \text{ on } \partial\Omega$$

in the weak sense.

Moreover  $\forall t, \xi, \hat{v}(t, \xi) \geq 0$ .

It follows that for all  $t \in \mathbb{R}_+$ ,

$$\int |E(t, x)|^2 dx \leq \int |E_0(x)|^2 dx.$$

The paper is organized as follows. In section 2, we formally derive the nonlinear model (1.8)-(1.11). In section 3, we introduce the dimensionless form of the system, and give the proofs of the main results. In section 4, we study a numerical scheme for (1.8)-(1.11) and present numerical results.

## 2 Formal derivation of the system.

The aim of this section is to present a formal derivation of system (1.8)-(1.11). For physical considerations, we refer to textbooks [5].

We consider here a plasma where collisions between the particles and the gravitational field are neglected. In this context, the Vlasov equation describes the evolution of the distribution function for each particle species  $\alpha$  of the collisionless plasma in the phase space  $(x, v)$  and time  $t$ . Denote by  $q_\alpha$  the particle charge of specy  $\alpha$  and by  $m_\alpha$  the mass of specy  $\alpha$ . The Vlasov equation reads :

$$\partial_t f_\alpha(x, v, t) + v \cdot \nabla_x f_\alpha(x, v, t) + \frac{q_\alpha}{m_\alpha} \left( E + \frac{1}{c} v \times B \right) \cdot \nabla_v f_\alpha(x, v, t) = 0, \quad (2.1)$$

$\alpha = e$  denotes the electrons and  $\alpha = i$  the ions. The fields  $E$  and  $B$  correspond to the electric and magnetic fields respectively and are given by the Maxwell equations :

$$\nabla \times B = \frac{4\pi}{c} j + \frac{1}{c} \partial_t E, \quad (2.2)$$

$$\nabla \times E = -\frac{1}{c} \partial_t B, \quad (2.3)$$

$$\nabla \cdot E = 4\pi \rho, \quad (2.4)$$

$$\nabla \cdot B = 0. \quad (2.5)$$

where  $\rho = -e \left( \int f_e(x, v, t) dv - \int f_i(x, v, t) dv \right)$  and  $j = -e \left( \int v f_e(x, v, t) dv - \int v f_i(x, v, t) dv \right)$  are the density of the total charge and total current respectively. The constant  $c$  is the velocity of the light in the vacuum. For a precise description of this kind of model, see classical textbooks [5]. Equations (2.2)-(2.3) yield :

$$\frac{1}{c^2} \partial_t^2 E + \nabla \times \nabla \times E = -\frac{4\pi}{c^2} \partial_t j, \quad (2.6)$$

and since the mass of the electrons is very small compared to the mass of the ions ( $m_e \ll m_i$ ) and the Lorentz force is the same for electrons and ions, the contribution of the ions

in the current  $j$  can be neglected, so the electric field satisfies :

$$\frac{1}{c^2} \partial_t^2 E + \nabla \times \nabla \times E = \frac{4\pi e}{c^2} \int v \partial_t f_e dv. \quad (2.7)$$

In a situation where an electromagnetic wave is injected into a collisionless plasma, one can identify one high-frequency time scale for the evolution of the electronic plasma wave and the high-frequency electromagnetic wave. In nuclear fusion by inertial confinement, the time scale is the order  $T = \frac{1}{\omega_{pe}} = 10^{-15} s$  where  $\omega_{pe}$  is the electron plasma frequency.

## 2.1 The high-frequency electron motion

The goal of this part is to obtain an equation for the slowly varying amplitude  $\mathcal{E}$  of the high frequency longitudinal electric field and its dependence on the slowly varying density fluctuation. The electric field is assumed to be decomposed as :

$$E = \frac{1}{2} (\mathcal{E} e^{-i\omega_{pe}t} + c.c.), \quad (2.8)$$

and we make the time envelope approximation assuming  $\partial_t \mathcal{E} \ll \omega_{pe} \mathcal{E}$ . Here *c.c.* denotes the complex conjugate.

For any function  $f$ , we define its average over the fast time scale by

$$\langle f \rangle = \frac{\omega_{pe}}{2\pi} \int_t^{t+\frac{2\pi}{\omega_{pe}}} f(x, s) ds$$

which gives the slowly varying components only.

We also define its average over the slow time scale by

$$\langle f e^{i\omega_{pe}t} \rangle e^{-i\omega_{pe}t}$$

which gives the slowly varying amplitude of the high-frequency component.

Plugging (2.8) in (2.7) gives :

$$\partial_t^2 \mathcal{E} - 2i\omega_{pe} \partial_t \mathcal{E} - \omega_{pe}^2 \mathcal{E} + c^2 \nabla \times \nabla \times \mathcal{E} = 4\pi e \langle e^{i\omega_{pe}t} \int v \partial_t f_e(x, v, t) \rangle. \quad (2.9)$$

We therefore have to found the contribution of the right-hand side of (2.9), that is the contribution of  $f_e$ .

We separate the behavior of the cold fluid from kinetic effects. We perform on the Vlasov equation (2.1) for electrons a change of the velocity reference frame. We introduce the jitter velocity  $u_e$  of the total high-frequency electric field defined by

$$m_e \frac{du_e}{dt} = q_e \left( E + \frac{1}{c} u_e \times B \right). \quad (2.10)$$

From now on, we work in the one dimension phase's space  $(x, v)$ . Let us define a distribution function  $F_e$  in the oscillation frame by  $F_e(x, w, t) = f_e(x, w + u_e, t)$ , the equation satisfied by  $F_e$  is

$$\partial_t F_e + w \partial_x F_e - u_e \partial_x u_e \partial_w F_e = w \partial_x u_e \partial_w F_e - u_e \partial_x F_e. \quad (2.11)$$

This formulation is convenient because it provides a clear separation of kinetic effects and cold fluid effects. For example the current for the electrons can be written as :

$$\begin{aligned} j(x, t) &= q_e u_e(x, t) \int F_e(x, w, t) dw + q_e \int w F_e(x, w, t) dw, \\ &= q_e u_e(x, t) n_e(x, t) + q_e \int w F_e(x, w, t) dw, \end{aligned}$$

where  $n_e u_e$  is the fluid contribution and the integral contains kinetic effects. Nevertheless, the distribution  $F_e$  contains high and low frequency components, so that we can introduce the following decomposition :

$$F_e(x, w, t) = F_0(x, w, t) + \frac{1}{2} (F_1 e^{-i\omega_{pe}t} + c.c.), \quad (2.12)$$

$$u_e = \frac{1}{2} (u_1 e^{-i\omega_{pe}t} + c.c.). \quad (2.13)$$

with  $\partial_t (F_0, F_1, u_1) \ll \omega_{pe} (F_0, F_1, u_1)$ .

Then plugging (2.12)-(2.13) in the right-hand side of (2.9) gives :

$$\partial_t^2 \mathcal{E} - 2i\omega_{pe} \partial_t \mathcal{E} - \omega_{pe}^2 \mathcal{E} = -i\omega_{pe} \left( 4\pi e u_1 \langle n_e \rangle + 4\pi e \int w F_1(x, w, t) dw \right). \quad (2.14)$$

By using (2.10), it follows that at the leading order:

$$u_1 = -i \frac{e\mathcal{E}}{m_e \omega_{pe}}. \quad (2.15)$$

Using the definition of  $\omega_{pe}$ ,  $\omega_{pe}^2 = \frac{4\pi e^2 n_0}{m_e}$ , where  $n_0$  is the background density, the slowly varying amplitude  $\mathcal{E}$  satisfies

$$\partial_t^2 \mathcal{E} - 2i\omega_{pe} \partial_t \mathcal{E} = \frac{\omega_{pe}^2}{n_0} (n_0 - \langle n_e \rangle) \mathcal{E} - i4\pi e \omega_{pe} \int w F_1(x, w, t) dw.$$

Now, we have to determine the contribution of the integral  $\int w F_1(x, w, t) dw$ . To this aim, we have to think of the solution of (2.11) under the form (2.12)-(2.13).

Keeping only at most quadratic terms, we find that  $(F_0, F_1)$  are given by

$$\partial_t F_0 + w \partial_x F_0 - \frac{1}{2} \partial_x |u_1|^2 \partial_w F_0 = \frac{1}{2} (w \partial_x u_1 \partial_w \overline{F_1} - u_1 \partial_x \overline{F_1}) + c.c., \quad (2.16)$$

$$\partial_t F_1 - i\omega_{pe} F_1 + w \partial_x F_1 = w \partial_x u_1 \partial_w F_0 - u_1 \partial_x F_0. \quad (2.17)$$

and since we have  $\partial_t F_1 \ll \omega_{pe} F_1$ ,  $F_1$  satisfies

$$-i\omega_{pe} F_1 + w \partial_x F_1 = w \partial_x u_1 \partial_w F_0 - u_1 \partial_x F_0. \quad (2.18)$$

Now, we want to compute  $\int w F_1(x, w, t) dw$ . Using the fact that for any distribution  $f$ , we have  $\widehat{\partial_x f} = i\hat{\xi} \hat{f}$ , the solution of (2.18) can be written as

$$F_1 = G *_{x,w} S(F_0)$$

where  $S(F_0)$  denotes the operator  $w\partial_x u_1 \partial_w F_0 - u_1 \partial_x F_0$  and  $G$  the Green's function associated with the equation (2.18)

$$G(x - x', v - v') = \frac{1}{2i\pi} \lim_{\varepsilon \rightarrow 0^+} \int e^{i\xi(x-x')} \frac{\delta(v-v')}{\xi \cdot v - \omega_{pe} + i\varepsilon} d\xi.$$

Then by using equation (2.18),  $F_1(x, w, t)$  can be written as

$$F_1 = -\frac{1}{i\omega_{pe}} (-w\partial_x F_1 + S(F_0)). \quad (2.19)$$

Once again, we use this relation in (2.19), which gives

$$F_1 = -\frac{1}{i\omega_{pe}} \left( -w\partial_x \left[ -\frac{1}{i\omega_{pe}} (-w\partial_x F_1 + S(F_0)) \right] + S(F_0) \right),$$

and finally, with  $F_1 = G *_{x,w} S(F_0)$ , we obtain

$$wF_1 = \frac{i}{\omega_{pe}} wS(F_0) + \frac{1}{\omega_{pe}^2} w^2 \partial_x S(F_0) - \frac{w^3}{\omega_{pe}^2} \partial_x^2 G * S(F_0). \quad (2.20)$$

Computing  $\int wF_1(x, w, t)dw$  thanks to (2.20) give us three terms.

- The contribution of the first term to the current is 0 since  $F_0(w) = F_0(-w)$ .

For the second and third term, we suppose that  $F_0$  can be decomposed as

$$F_0(x, w, t) = F_{00}(w, t) + \delta F_0(x, w, t). \quad (2.21)$$

where  $F_{00}$  is the spatially averaged distribution with  $|\delta F_0| \ll |F_{00}|$ . We also define, the electronic thermal velocity  $v_e$  and the local fluctuation of velocity  $\delta v_e(x, t)$  ( $\delta v_e \ll v_e$ ) by

$$n_0 v_e^2 = \int w^2 F_{00} dw, \quad n_0 \delta v_e^2 = \int w^2 \delta F_0 dw.$$

After some integrations by parts in  $w$ , the second term exactly yields

$$\int w^2 \partial_x S(F_0) dw = -3n_0 v_e^2 \partial_x^2 u_1 - 3n_0 \partial_x (\delta v_e^2 \partial_x u_1) - n_0 \partial_x (u_1 \partial_x \delta v_e^2). \quad (2.22)$$

Neglecting the two last terms in (2.22) one obtains

$$\int w^2 \partial_x S(F_0) dw \approx -3n_0 v_e^2 \partial_x^2 u_1. \quad (2.23)$$

The contribution of the third term gives in (2.20)

$$\partial_x^2 \int_{\mathbb{R}_v} v^3 G * S(F_0) dv = \partial_x^2 \int_{\mathbb{R}_v} v^3 \left( \int_{\mathbb{R}_w} \int_{\mathbb{R}_y} G(x-y, v-w) S(F_0)(y, w) dy dw \right) dv \quad (2.24)$$

Then plugging the expression of  $G$  in (2.24) gives

$$\partial_x^2 \int_{\mathbb{R}_v} v^3 G * S(F_0) dv = -\frac{1}{2\pi} \frac{1}{i} \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} e^{i\xi(x-y)} I(y, \xi, t) d\xi dy,$$



where  $I(y, v, t)$  denotes the integral

$$I(y, \xi, t) = \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} \frac{\beta(y, v, t)}{\xi v - \omega_{pe} + i\varepsilon} dv$$

which value is given by

$$I(y, \xi, t) = \frac{1}{\xi} \mathcal{P} \left( \int_{\mathbb{R}_v} \frac{\beta(y, v, t)}{v - \frac{\omega_{pe}}{\xi}} dv \right) + \frac{1}{|\xi|} i\pi \beta \left( y, \frac{\omega_{pe}}{\xi}, t \right)$$

where  $\mathcal{P}$  denotes the Cauchy Principal value and

$$\beta(y, v, t) = (\xi v)^2 \left( v^2 \partial_y u_1(y, t) \partial_v F_0(y, v, t) - u_1(y, t) v \partial_y F_0(y, v, t) \right).$$

The more important part of  $I(y, \xi, t)$  is the second one ( see [5]) and one gets

$$\partial_x^2 \int_{\mathbb{R}_v} v^3 G * S(F_0) dv = -\pi \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \beta(y, \frac{\omega_{pe}}{\xi}, t) d\xi dy. \quad (2.25)$$

We will therefore focus on the second term in (2.25) for which  $F_{00}$  will play a role.

With

$$\beta \left( y, \frac{\omega_{pe}}{\xi}, t \right) = \left( \frac{\omega_{pe}^4}{\xi^2} \partial_y u_1(y, t) \partial_v F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) - u_1(y, t) \frac{\omega_{pe}^3}{\xi} \partial_y F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) \right),$$

we write :

$$\begin{aligned} \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \beta(y, \frac{\omega_{pe}}{\xi}, t) d\xi dy &= \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \frac{\omega_{pe}^4}{\xi^2} \partial_y u_1 \partial_v F_{00} \left( \frac{\omega_{pe}}{\xi}, t \right) d\xi dy \\ &+ \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \frac{\omega_{pe}^4}{\xi^2} \partial_y u_1 \partial_v \delta F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) d\xi dy \\ &+ \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \frac{\omega_{pe}^3}{\xi} u_1(y, t) \partial_y \delta F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) d\xi dy. \end{aligned}$$

Integrating by part with respect to  $y$  the first integral, give us

$$\begin{aligned} \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \beta(y, \frac{\omega_{pe}}{\xi}, t) d\xi dy &= i\omega_{pe}^4 \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|\xi} u_1 \partial_v F_{00} \left( \frac{\omega_{pe}}{\xi}, t \right) d\xi dy \\ &+ \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \frac{\omega_{pe}^4}{\xi^2} \partial_y u_1 \partial_v \delta F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) d\xi dy \\ &+ \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|} \frac{\omega_{pe}^3}{\xi} u_1(y, t) \partial_y \delta F_0 \left( y, \frac{\omega_{pe}}{\xi}, t \right) d\xi dy. \end{aligned}$$

Neglecting the two last terms (since  $\delta F_0 \ll F_{00}$ ), we obtain that the contribution of this term in the current is :

$$\begin{aligned} \frac{4i\pi e}{\omega_{pe}^2} \partial_x^2 \int v^3 G * S(F_0) dv &= \frac{4i\pi e}{\omega_{pe}^2} i\omega_{pe}^4 \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|\xi} u_1(t, y) \partial_v F_{00} \left( t, \frac{\omega_{pe}}{\xi} \right) d\xi dy \\ &= \frac{4\pi^2 e}{\omega_{pe}^2} \omega_{pe}^4 \frac{ie}{m_e \omega_{pe}} \int_{\mathbb{R}_y} \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi|\xi|\xi} \mathcal{E}(t, y) \partial_v F_{00} \left( t, \frac{\omega_{pe}}{\xi} \right) d\xi dy \end{aligned}$$

where (2.15) has been used.

So, we have

$$\begin{aligned} \frac{4i\pi e}{\omega_{pe}^2} \partial_x^2 \int v^3 G * S(F_0) dv &= 2i\omega_{pe} \int_{\mathbb{R}_y} \mathcal{E}(t, y) \left( \int_{\mathbb{R}_\xi} \frac{e^{i\xi(x-y)}}{2\pi} \frac{-\pi\omega_{pe}^3}{2n_0|\xi|\xi} \partial_v F_{00} \left( t, \frac{\omega_{pe}}{\xi} \right) d\xi \right) dy \\ &= 2i\omega_{pe} \int_{\mathbb{R}_y} \mathcal{E}(t, y) \nu(t, x - y) dy \\ &= 2i\omega_{pe} \mathcal{E} * \nu \end{aligned}$$

with

$$\nu(t, x - y) = \frac{1}{2\pi} \int_{\mathbb{R}_\xi} e^{i\xi(x-y)} \hat{\nu}(t, \xi) d\xi$$

where

$$\hat{\nu}(t, \xi) = -\frac{\pi\omega_{pe}^3}{2n_0|\xi|\xi} \partial_v F_{00} \left( t, \frac{\omega_{pe}}{\xi} \right). \quad (2.26)$$

Finally, at the first order, the slowly varying envelope satisfies the nonlinear Schrödinger equation

$$2i\omega_{pe} \partial_t \mathcal{E} + 2i\omega_{pe} \mathcal{E} * \nu + 3v_e^2 \partial_x^2 \mathcal{E} = \frac{\omega_{pe}^2}{n_0} (< n_e > - n_0) \mathcal{E}$$

with  $\hat{\nu}$  given by (2.26).

In order to close the system, we have to find an equation evolving  $F_{00}$  and an equation evolving the low-frequency  $\delta n_e = < n_e > - n_0$ .

The equation involving the spatially averaged distribution  $F_{00}$  is recovered by substitute the Green's function solution of (2.18) into the right-hand side of (2.16) and taking the spatial average we find the usual quasilinear diffusion equation (see [18])

$$\partial_t F_{00} - \partial_v (D(t, v) \partial_v F_{00}) = 0, \text{ with } D(t, v) = \frac{e^2}{m_e^2 |v|} \left| \hat{\mathcal{E}} \left( t, \frac{\omega_{pe}}{v} \right) \right|^2,$$

where the diffusion coefficient  $D(t, v)$  depends on the density spectral energy  $\left| \hat{\mathcal{E}} \left( \frac{\omega_{pe}}{v} \right) \right|^2$ . As usual, the plasma is assumed to be quasi-neutral on the slow ion acoustic time scale, that is  $\delta n_e = \delta n_i$  with  $\delta n_i = n_i - n_0$ .

Now we look for an equation involving  $\delta n_i$ . Since  $m_i \gg m_e$ , the study is much simpler and we assume that the ion distribution function is Maxwellian.

In this context, one can see in [4], [17], [22] and [23], that  $\delta n_i$  satisfies the wave equation

$$\partial_t^2 \delta n_i - c_s^2 \partial_x^2 \delta n_i = \frac{1}{16\pi m_i} \partial_x^2 (|\mathcal{E}|^2),$$

where  $c_s = \sqrt{\frac{T_e}{m_i}}$  is the ion acoustic velocity.

Finally, we have to deal with the following system

$$2i\omega_{pe} (\partial_t \mathcal{E} + \nu * \mathcal{E}) + 3v_e^2 \partial_x^2 \mathcal{E} = \frac{\omega_{pe}^2}{n_0} \delta n \mathcal{E}, \quad (2.27)$$

$$\partial_t^2 \delta n - c_s^2 \partial_x^2 \delta n = \frac{1}{16\pi m_i} \partial_x^2 (|\mathcal{E}|^2), \quad (2.28)$$

$$\widehat{\nu}(\xi, t) = -\frac{\pi \omega_{pe}^3}{2\xi n_0 |\xi|} \partial_v F_e\left(\frac{\omega_{pe}}{\xi}\right), \quad (2.29)$$

$$\partial_t F_0 = \partial_v (D(v, t) \partial_v F_0), \quad D(v, t) = \frac{e^2}{m_e^2 |v|} \left| \widehat{\mathcal{E}}\left(\xi = \frac{\omega_{pe}}{v}, t\right) \right|^2. \quad (2.30)$$

It describes the interaction between the high-frequency envelope of the longitudinal electric field and the low-frequency density fluctuation (wave-wave process) and the resonant interaction between the electrons and the longitudinal electron plasma waves (wave-particle process).

Usually the electron plasma waves are created by a pump wave obtained by the stimulated Raman scattering, acting as a source term in (2.27) (see [4]). However, it is possible to add a given pump wave in the derivation of the Schrödinger equation by changing the envelope approximation (2.8) by

$$E(x, t) = \frac{1}{2} (E_p(t, x) e^{i(kx - \omega t)} + \mathcal{E}(t, x) e^{-i\omega_{pe} t} + c.c.),$$

and we can replace (2.27) with

$$2i\omega_{pe} (\partial_t \mathcal{E} + \nu * \mathcal{E}) + 3v_e^2 \partial_x^2 \mathcal{E} = \frac{\omega_{pe}^2}{n_0} \delta n \mathcal{E} + \omega_{pe}^2 E_p e^{i(k_1 x - \omega_1 t)}. \quad (2.31)$$

Since the dispersion relation of the linear part of (2.31) is  $\omega = \frac{3v_e^2 k^2}{2\omega_{pe}}$ , we choose  $(k_1, \omega_1)$  satisfying this relation.

## 2.2 Dimensionless form

We now introduce a dimensionless form of (2.27)-(2.30).

We use  $T = \frac{1}{\omega_{pe}}$  as time scale and  $L = \lambda_{De}$  as space scale (where  $\lambda_{De} = \frac{v_e}{\omega_{pe}}$  is the Debye's length) and introduce

$$\begin{aligned} \widetilde{E} &= \frac{e}{m_e v_e \omega_{pe}} \mathcal{E}, & \widetilde{\nu} &= \frac{1}{\omega_{pe}} \widehat{\nu}, \\ \widetilde{k}_1 &= k_1 \lambda_{De}, & \widetilde{\omega}_1 &= \frac{\omega_1}{\omega_{pe}}, \\ \widetilde{F}_0 &= \frac{v_e}{n_0} F_0, & \widetilde{\delta n} &= \frac{1}{n_0} \delta n. \end{aligned}$$

Dropping the tildes, we get the following system :

$$2i(\partial_t E + \nu * E) + 3\partial_x^2 E = \delta n E + E_p(x)e^{i(k_1 x - \omega_1 t)}, \quad (2.32)$$

$$\partial_t^2 \delta n - \mu \partial_x^2 \delta n = \frac{\mu}{4} \partial_x^2 (|E|^2), \quad (2.33)$$

$$\hat{\nu}(t, \xi) = -\frac{\pi}{2\xi|\xi|} \partial_v F_e(t, \frac{1}{\xi}), \quad (2.34)$$

$$\partial_t F_e = \partial_v (D(t, v) \partial_v F_e), \quad D(t, v) = \frac{1}{4|v|} \left| \hat{E}\left(t, \frac{1}{v}\right) \right|^2, \quad (2.35)$$

where  $\mu = \frac{c_s^2}{v_e^2} = \frac{m_e}{m_i}$ .

Usually, (2.32)-(2.33) are satisfied on the whole space  $\mathbb{R}$ . But (2.34)-(2.35) are valid only for bounded speeds and far away from zero. The velocity domain that we consider is

$$v \in \Omega_v = [-A, -a] \cup [a, A], \quad (A > a > 0),$$

which gives frequency domain of the form,

$$\xi \in \Omega_\xi = [-a^{-1}, -A^{-1}] \cup [A^{-1}, a^{-1}], \quad (A > a > 0).$$

Apart from this set,  $\hat{\nu}_e(., \xi)$  is extended to 0.

Moreover, in order to study the system (2.32)-(2.35), it is more convenient to write it by using the variable  $\xi$  rather than the variable  $v$  on equations (2.34)-(2.35). Then denoting

$$H_e(t, \xi) = F_0(t, \frac{1}{\xi}),$$

the system that we study is (we set all coefficients equal to one excepted  $\mu$ )

$$\begin{aligned} i(\partial_t E + \nu * E) + \partial_x^2 E &= \delta n E + E_p(x)e^{i(k_1 x - \omega_1 t)}, \quad x \in \mathbb{R}, t \geq 0, \\ \frac{1}{\mu} \partial_t^2 \delta n - \partial_x^2 \delta n &= \partial_x^2 (|E|^2), \quad x \in \mathbb{R}, t \geq 0, \end{aligned} \quad (2.36)$$

$$\begin{aligned} \partial_t H_e - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}(t, \xi)|^2 \partial_\xi H_e) &= 0, \quad \xi \in \Omega_\xi, t \geq 0, \\ \hat{\nu}(t, \xi) &= \text{sgn}(\xi) \partial_\xi H_e(t, \xi) 1_{\Omega_\xi}. \end{aligned} \quad (2.37)$$

The boundary conditions are

$$\partial_\xi H_e|_{\partial\Omega_\xi} = 0, \quad \hat{\nu}|_{\partial\Omega_\xi} = 0. \quad (2.38)$$

### 3 Proofs of the main results.

Let us consider the system (2.36)-(2.37). In the following, we will replace  $\Omega_\xi$  by  $\Omega$ . Unfortunately, we are not able to handle the general case. So we will consider only the case  $\mu = +\infty$  corresponding to a subsonic regime. Therefore, the system (2.36)-(2.37) to

be solved becomes

$$\begin{aligned}
i(\partial_t E + \nu * E) + \partial_x^2 E &= |E|^2 E + S(t, x), \\
\partial_t H_e - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 \partial_\xi H_e) &= 0, \quad \forall \xi \in \Omega, \\
\partial_\xi H_e|_\Omega &= 0, \\
\hat{\nu}(\xi, \cdot) &= \text{sgn}(\xi) \partial_\xi H_e 1_\Omega, \\
H_e(\cdot, 0) &= H_{e0}(\cdot), E(\cdot, 0) = E_0(\cdot).
\end{aligned} \tag{3.1}$$

The functional space for  $H_e$  is

$$H_n^2(\Omega) = \{u \in H^2(\Omega) \text{ such that } \partial_n u|_{\partial\Omega} = 0\}.$$

In order to simplify computations, and without loss of generality, we take  $\Omega = ]-2, -1[ \cup ]1, 2[$ . The first step of the proof is the construction of solutions given by theorem (1.1) for a regularized problem. The regularization is obtained by taking a non degenerate dissipation on equation (2.37), namely one replaces  $|\xi|^3 |\hat{E}|^2$  by  $|\xi|^3 |\hat{E}|^2 + \varepsilon$  with  $\varepsilon > 0$ .

Moreover the left hand side of the first equation of (3.1) is replaced by a source term  $f$ . One then obtains local existence for solutions whose time of existence depends on  $\varepsilon$  and  $f$ . We obtain a solution satisfying

$$(E, \hat{E}, H_e) \in L^\infty(0, T^*; H^1(\mathbb{R})) \times L^\infty(0, T^*; H^2(\Omega) \cap H^1(\mathbb{R})) \times L^\infty(0, T^*; H_n^2(\Omega)).$$

The next step consists in replacing  $f$  by  $|E|^2 E$  by using a fixed point method. The important point is that  $H^1$  is an algebra and  $E, \hat{E}$  are in  $H^1$  so that

$$\widehat{|E|^2 E} \in W^{2,\infty}(\mathbb{R})$$

and therefore

$$\widehat{|E|^2 E} \in H^2(\Omega).$$

The last step consists in obtaining estimates that are uniform with respect to  $\varepsilon$ . In physical situation the term  $\nu * E$  is a damping term. The positivity of  $\hat{\nu}(t, \xi)$  is then obtained by using the maximum principle on equation (2.34). In order to prove theorem (1.2), we use an algebraic cancellation between the left hand side of (2.32) and the dissipative part of (2.35). We now prove both theorems.

### 3.1 Local existence of weak solutions

In order to prove theorems 1.1 and 1.2, we consider a problem with non degenerate dissipation on the diffusion equation.

#### 3.1.1 Non degenerated simplified problem.

For  $\varepsilon > 0$ , we consider

$$\begin{aligned}
i(\partial_t E^\varepsilon + \nu^\varepsilon * E^\varepsilon) + \partial_x^2 E^\varepsilon &= f, \\
\partial_t H_e^\varepsilon - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 + \varepsilon) \partial_\xi H_e^\varepsilon &= 0, \quad \forall \xi \in \Omega, \\
\partial_\xi H_e^\varepsilon|_\Omega &= 0, \\
\hat{\nu}_e^\varepsilon &= \text{sgn}(\xi) \partial_\xi H_e^\varepsilon 1_\Omega, \\
H_e^\varepsilon(0, \cdot) &= H_{e0}(\cdot), E^\varepsilon(0, \cdot) = E_0(\cdot),
\end{aligned} \tag{3.2}$$

with  $f \in L^\infty(0, T; H^1(\mathbb{R}))$ ,  $\hat{f} \in L^\infty(0, T; H^2(\Omega) \cap H^1(\mathbb{R}))$ .

**Proposition 3.1.** *Let  $E_0 \in H^1(\mathbb{R})$  such that  $\hat{E}_0 \in H^1(\mathbb{R})$ , and let  $H_{e0} \in H^1(\Omega)$ , then there exist  $T^* > 0$  and a unique solution  $(E^\varepsilon, H_e^\varepsilon)$  of (3.2) such that*

$$(E^\varepsilon, \hat{E}^\varepsilon, H_e^\varepsilon) \in C^0([0, T^*]; H^1(\mathbb{R})) \times C^0([0, T^*]; H^1(\mathbb{R})) \times C^0([0, T^*]; H^1(\Omega)).$$

*If moreover  $\hat{E}_0 \in H^2(\Omega) \cap H^1(\mathbb{R})$ , and  $H_{e0} \in H_n^2(\Omega)$  then*

$$(E^\varepsilon, \hat{E}^\varepsilon, H_e^\varepsilon) \in C^0([0, T^*]; H^1(\mathbb{R})) \times C^0([0, T^*]; H^2(\Omega) \cap H^1(\mathbb{R})) \times C^0([0, T^*]; H_n^2(\Omega)).$$

The proof of this proposition is obtained with a fixed point method.

Let us consider a function  $G_e : (x, t) \in \mathbb{R} \times \mathbb{R}^+ \rightarrow \mathbb{R}$  such that  $G_e \in L^\infty(\mathbb{R}^+, H_n^2(\Omega))$ , and define  $\mu_e = \text{sgn}(\xi) \partial_\xi G_e 1_\Omega$ . Consider  $E^\varepsilon$ ,  $H_e^\varepsilon$  and  $\nu^\varepsilon$  solution of the following system,

$$\begin{aligned} i(\partial_t E^\varepsilon + \mu_e * E^\varepsilon) + \partial_x^2 E^\varepsilon &= f, \\ \partial_t H_e^\varepsilon - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 + \varepsilon) \partial_\xi H_e^\varepsilon &= 0, \quad \forall \xi \in \Omega, \\ \partial_\xi H_e^\varepsilon|_\Omega &= 0, \\ \hat{\nu}_e^\varepsilon &= \text{sgn}(\xi) \partial_\xi H_e^\varepsilon 1_\Omega, \\ H_e^\varepsilon(., 0) &= H_{e0}(.), E(., 0) = E_0(.). \end{aligned} \tag{3.3}$$

We want to show that the map  $\tau : G_e \mapsto H_e^\varepsilon$  is a contraction on a suitable metric spaces. For any  $R > 0$ , we denote by  $B_R(H)$  the ball of the space  $H$  centered on 0 with radius  $R$ . For  $R$  large enough and  $T$  small enough, we show that  $\tau$  maps

$$B_R(L^\infty(0, T; H^1(\mathbb{R}) \cap \mathcal{F}(H^1(\mathbb{R})) \cap H_n^2(\Omega)) \cap L^2(0, T, H_n^3(\Omega)))$$

into itself and maps

$$B_R(L^\infty(0, T; H^1(\mathbb{R}) \cap \mathcal{F}(H^1(\mathbb{R}))) \cap L^2(0, T, H_n^2(\Omega)))$$

into itself.

Let us prove this last result.

Performing an  $L^2$  estimate on the first and second equation of (3.3) yields

$$\begin{aligned} \frac{d}{dt} |\hat{E}|_{L^2(\mathbb{R})}^2 &\leq 2 \int_\Omega |\hat{\mu}_e| |\hat{E}|^2 + 2 \int_{\mathbb{R}} |\hat{f}| |\hat{E}| \\ &\leq c |\hat{\mu}_e|_{H^1(\Omega)} |\hat{E}|_{L^2(\Omega)}^2 + |\hat{f}|_{L^2(\mathbb{R})}^2 + |\hat{E}|_{L^2(\mathbb{R})}^2, \end{aligned}$$

and

$$\frac{d}{dt} |\xi^{-1} H_e^\varepsilon|_{L^2(\Omega)}^2 + 2 \int_\Omega (|\xi|^3 |\hat{E}|^2 + \varepsilon) |\partial_\xi H_e^\varepsilon|^2 = 0.$$

We now perform an  $H^1$  estimate in the  $\xi$  space

$$\frac{d}{dt} |\partial_\xi \hat{E}|_{L^2(\mathbb{R})}^2 + 2 \int_\Omega \hat{\mu}_e |\partial_\xi \hat{E}|^2 + 2\Im \left( \int_\Omega \partial_\xi \hat{\mu}_e \hat{E} \partial_\xi \bar{\hat{E}} \right) = 2\Im \left( \int_{\mathbb{R}} \partial_\xi \hat{f} \partial_\xi \bar{\hat{E}} \right) + 4\Im \left( \int_{\mathbb{R}} \xi \hat{E} \partial_\xi \bar{\hat{E}} \right).$$

This gives,

$$\frac{d}{dt} |\partial_\xi \hat{E}|_{L^2(\mathbb{R})}^2 \leq c |\hat{\mu}_e|_{H^1(\Omega)} |\hat{E}|_{H^1(\Omega)}^2 + |\partial_\xi \hat{f}|_{L^2(\mathbb{R})}^2 + 2 |\partial_\xi \hat{E}|_{L^2(\mathbb{R})}^2 + C |\hat{E}|_{H^1(\mathbb{R})}^2 + c |E|_{H^1(\mathbb{R})}^2.$$

An  $H^1$  estimate on  $H_e^\varepsilon$  leads to

$$\frac{d}{dt} |\partial_\xi H_e^\varepsilon|_{L^2(\Omega)}^2 + 2 \int_\Omega \xi^2 (|\xi|^3 |\hat{E}|^2 + \varepsilon) |\partial_\xi H_e^\varepsilon|^2 \leq \varepsilon |\partial_\xi H_e^\varepsilon|_{L^2(\Omega)}^2 + \varepsilon^{-1} |\xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2)|_{L^1(\Omega)}.$$

Summing theses estimates, it follows that,

$$\begin{aligned} \frac{d}{dt} \left( |\hat{E}|_{H^1(\mathbb{R})}^2 + |\xi^{-1} H_e^\varepsilon|_{L^2(\Omega)}^2 + |\xi^{-1} \partial_\xi H_e^\varepsilon|_{L^2(\Omega)}^2 \right) + \int_\Omega (|\xi|^3 |\hat{E}| + \varepsilon) (\xi^2 |\partial_\xi H_e^\varepsilon|^2 + |H_e^\varepsilon|^2) \leq \\ c(1 + \varepsilon^{-1} + |\hat{\mu}_e|_{H^1}) |\hat{E}|_{H^1(\mathbb{R})}^2 + |\hat{f}|_{H^1(\Omega)}^2 + c|E|_{H^1(\mathbb{R})}^2. \end{aligned}$$

We still have to obtain an  $H^1$  estimate for  $E^\varepsilon$  in the  $x$  space. We apply  $\partial_x$  on the first equation of (3.3), multiplying by  $\partial_x \overline{E}$  and take the imaginary part gives :

$$\frac{d}{dt} \int |\partial_x E|^2 \leq 2|\hat{\mu}_e|_{L^\infty} |\partial_x E|_{L^2}^2 + 2|\partial_x f|_{L^2} |\partial_x E|_{L^2}.$$

That leads to

$$\frac{d}{dt} \int |\partial_x E|^2 \leq (2|\hat{\mu}_e|_{L^\infty} + 1) |\partial_x E|_{L^2}^2 + |\partial_x f|_{L^2}^2.$$

Thanks to the non degenerate diffusion term, we recover the estimate of  $\hat{\nu}$  in  $L^2(0, T; H^1(\Omega))$ . By choosing  $T$  small enough (depending on  $\varepsilon$ ) and  $R$  large enough (depending on the size of initial data), the map  $\tau$  which associated  $G_e$  with  $H_e^\varepsilon$  maps

$$B_R \left( L^\infty(0, T; H^1(\mathbb{R}) \cap \mathcal{F}(H^1(\mathbb{R}))) \cap L^2(0, T, H_n^2(\Omega)) \right)$$

into itself.

The contraction property is obtained in  $L^\infty(0, T; L^2(\mathbb{R}) \cap \mathcal{F}(H^1(\mathbb{R}))) \cap L^2(0, T, H^1(\Omega))$ . Let  $G_e^1, G_e^2$  belonging to  $B_R(L^\infty(0, T; H^1(\mathbb{R}) \cap \mathcal{F}(H^1(\mathbb{R}))) \cap L^2(0, T, H_n^2(\Omega)))$ . Denote by  $(E^1, H_e^{\varepsilon,1}, \nu^{\varepsilon,1}), (E^2, H_e^{\varepsilon,2}, \nu^{\varepsilon,2})$  the solutions of (3.3) associated with  $G_e^1$  and  $G_e^2$  respectively. Denote by  $E = E^1 - E^2, H_e^\varepsilon = H_e^{\varepsilon,1} - H_e^{\varepsilon,2}, \nu_e^\varepsilon = \nu^{\varepsilon,1} - \nu^{\varepsilon,2}$ . Then for all  $\eta > 0$ , we have,

$$\frac{d}{dt} |\hat{E}|_{L^2(\Omega)}^2 \leq K(\eta, R) |\hat{E}|_{L^2(\Omega)}^2 + \eta |\mu_e|_{L^2(\Omega)}^2,$$

where  $K$  depends of  $\eta$  and  $R$ .

$$\frac{d}{dt} |H_e^\varepsilon|_{L^2(\Omega)}^2 + \int_\Omega (|\hat{E}|^2 + \varepsilon) |\partial_\xi H_e^\varepsilon|^2 \leq C(R, \varepsilon) |\hat{E}|_{L^2(\Omega)}^2.$$

According to Gronwall's lemma, we easily conclude that

$$|H_e^\varepsilon|_{L^\infty(0, T; L^2(\Omega))}^2 + \varepsilon |\partial_\xi H_e^\varepsilon|_{L^2(0, T; L^2(\Omega))}^2 \leq C(R, \varepsilon) \exp(KT) \eta |\mu_e|_{L^2(0, T; L^2(\Omega))}^2.$$

The contraction property is obtained by choosing  $\eta$  and  $T$  small enough. ■

### 3.1.2 The complete non degenerated problem

For  $\varepsilon > 0$ , we consider

$$\begin{aligned}
i(\partial_t E^\varepsilon + \nu^\varepsilon * E^\varepsilon) + \partial_x^2 E^\varepsilon &= |E^\varepsilon|^2 E^\varepsilon + S, \\
\partial_t H_e^\varepsilon - \xi^2 \partial_\xi((|\xi|^3 |\hat{E}^\varepsilon|^2 + \varepsilon) \partial_\xi H_e^\varepsilon) &= 0, \quad \forall \xi \in \Omega, \\
\partial_\xi H_e^\varepsilon|_\Omega &= 0, \\
\hat{\nu}^\varepsilon(., \xi) &= \text{sgn}(\xi) \partial_\xi H_e^\varepsilon 1_\Omega, \\
H_e^\varepsilon(., 0) &= H_{e0}(.), E^\varepsilon(., 0) = E_0(.).
\end{aligned} \tag{3.4}$$

**Proposition 3.2.** *Let  $S \in C^0(\mathbb{R}^+; H^1(\mathbb{R}))$  such that  $\hat{S} \in C^0(\mathbb{R}^+; W^{2,\infty}(\mathbb{R}))$ . Let  $E_0 \in H^1(\mathbb{R})$  such that  $\hat{E}_0 \in H^1(\mathbb{R})$ , and let  $H_{e0} \in H^1(\Omega)$ . Then there exists  $T^* > 0$  and a unique solution  $(E^\varepsilon, H_e^\varepsilon)$  of (3.4) such that*

$$(E^\varepsilon, \hat{E}^\varepsilon, H_e^\varepsilon) \in C^0(0, T^*; H^1(\mathbb{R})) \times C^0(0, T^*; H^1(\mathbb{R})) \times C^0(0, T^*; H^1(\Omega)).$$

If moreover  $\hat{E}_0 \in H^1(\mathbb{R}) \cap H^2(\Omega)$ , and  $H_{e0} \in H_n^2(\Omega)$ , one has

$$(E^\varepsilon, \hat{E}^\varepsilon, H_e^\varepsilon) \in C^0(0, T^*; H^1(\mathbb{R})) \times C^0(0, T^*; H^1(\mathbb{R}) \cap H^2(\Omega)) \times C^0(0, T^*; H_n^2(\Omega)).$$

According to proposition 3.1, problem (3.2) has a solution. Denote by  $f = |D|^2 D + S$  where  $D$  belongs to  $C^0([0, T^*[, H^1(\mathbb{R}))$  and  $\hat{D}$  belongs to  $C^0([0, T^*[, H^1(\mathbb{R}))$ . It follows that  $f$  is in  $C^0([0, T^*[, H^1(\mathbb{R}))$  because  $H^1(\mathbb{R})$  is an algebra in one dimension and moreover  $\hat{f} = \hat{D} \star \hat{\bar{D}} \star \hat{D} + \hat{S}$  belongs to  $C^0([0, T^*[, H^1(\mathbb{R}))$  because  $\hat{D}$  is in  $C^0([0, T^*[, L^1(\mathbb{R}))$ .

In the same way,

$$\partial_\xi^2 \hat{f} = \hat{D} \star \partial_\xi \hat{\bar{D}} \star \partial_\xi \hat{D} + \partial_\xi^2 \hat{S}$$

belongs to  $C^0([0, T^*[, L^\infty(\mathbb{R}))$  resulting from the convolution between a  $L^1(\mathbb{R})$ -function and two  $L^2(\mathbb{R})$ -functions.

We will show the above proposition by using a fixed point method on the map  $\tau$  which associates  $D$  with  $E^\varepsilon$ ,  $H_e^\varepsilon$  and  $\nu^\varepsilon$  solutions of

$$\begin{aligned}
i(\partial_t E^\varepsilon + \nu^\varepsilon * E^\varepsilon) + \partial_x^2 E^\varepsilon &= |D|^2 D + S, \\
\partial_t H_e^\varepsilon - \xi^2 \partial_\xi((|\xi|^3 |\hat{E}^\varepsilon|^2 + \varepsilon) \partial_\xi H_e^\varepsilon) &= 0, \quad \forall \xi \in \Omega, \\
\partial_\xi H_e^\varepsilon|_\Omega &= 0, \\
\hat{\nu}^\varepsilon(., \xi) &= \text{sgn}(\xi) \partial_\xi H_e^\varepsilon 1_\Omega, \\
H_e^\varepsilon(., 0) &= H_{e0}(.), E^\varepsilon(., 0) = E_0(.).
\end{aligned} \tag{3.5}$$

If  $H_{e0} \in H^1(\Omega)$ ,  $E_0 \in H^1(\mathbb{R})$  and  $\hat{E}_0 \in H^1(\mathbb{R})$ , we show that  $\tau$  maps  $B_R(H_T)$  into itself, for  $R$  large enough and  $T$  small enough, where  $H_T$  is

$$H_T = \{e \in L^\infty(0, T; H^1(\mathbb{R})) \text{ s.t. } \hat{e} \in L^\infty(0, T; H^1(\mathbb{R})) \text{ and } e \in L^\infty(0, T; H^2(\mathbb{R}))\}.$$

The contraction property is obtained in  $L^\infty(0, T; L^2(\mathbb{R}))$ .

If  $H_{e0} \in H_n^2(\Omega)$ ,  $E_0 \in H^1(\mathbb{R})$  and  $\hat{E}_0 \in H^2(\Omega) \cap H^1(\mathbb{R})$ , we also show that  $\tau$  maps  $B_R(V_T)$  into itself, for  $R$  large enough and  $T$  small enough, where  $V_T$  is

$$V_T = \{e \in L^\infty(0, T, H^1(\mathbb{R})) \text{ s.t. } \hat{e} \in L^\infty(0, T, H^1(\mathbb{R}) \cap H^2(\Omega))\}.$$

■



### 3.1.3 Uniform estimates in $\varepsilon$

At this point, we have a solution to problem (3.4). We will obtain uniform estimates in  $\varepsilon$  for the solutions of (3.4) in order to let  $\varepsilon$  tends to 0.

**Proposition 3.3.** *The solution of (3.4) satisfies for all  $t \leq T^*$*

$$\begin{aligned} & |E^\varepsilon(t)|_{H^1(\mathbb{R})}^2 + |\hat{E}^\varepsilon(t)|_{H^1(\mathbb{R})}^2 + |\partial_\xi^2 \hat{E}^\varepsilon(t)|_{L^2(\Omega)}^2 + |H_e(t)|_{L^2(\Omega)}^2 + |\partial_\xi^2 H_e(t)|_{L^2(\Omega)}^2 \\ & \quad + \int_0^t \int_\Omega (|\hat{E}|^2 + \varepsilon) |\partial_\xi^3 H_e|^2 \\ & \leq C(t) \left( |E_0|_{H^1(\mathbb{R})}^2 + |\hat{E}_0|_{H^1(\mathbb{R})}^2 + |\partial_\xi^2 \hat{E}_0|_{L^2(\Omega)}^2 + |H_{e0}|_{L^2(\Omega)}^2 + |\partial_\xi^2 H_{e0}|_{L^2(\Omega)}^2 \right) \\ & \quad + |S|_{H^1(\mathbb{R})}^2 + |\hat{S}|_{H^1(\mathbb{R})}^2 + |\partial_\xi^2 \hat{S}|_{L^2(\Omega)}^2, \end{aligned} \quad (3.6)$$

where the function  $C(t)$  does not depend on  $\varepsilon$ .

The  $L^2$  estimates on  $\hat{E}^\varepsilon$  and  $H_e^\varepsilon$  are

$$\begin{aligned} \frac{d}{dt} |\hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2 & \leq 2|\hat{\nu}^\varepsilon \hat{E}^\varepsilon|_{L^2(\Omega)} |\hat{E}^\varepsilon|_{L^2(\Omega)} + |\hat{S}|_{L^2(\mathbb{R})}^2 + |\hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2, \\ \frac{d}{dt} |\xi^{-1} H_e^\varepsilon|_{L^2(\Omega)}^2 & + 2 \int_\Omega (|\xi|^3 |\hat{E}^\varepsilon|^2 + \varepsilon) |\partial_\xi H_e^\varepsilon|^2 \leq 0, \end{aligned}$$

while the  $H^1$  estimates give

$$\frac{d}{dt} |\partial_x E^\varepsilon|_{L^2(\mathbb{R})}^2 + 2 \int_\Omega \hat{\nu}_e^\varepsilon \hat{E}^\varepsilon \xi^2 \overline{\hat{E}^\varepsilon} \leq 6 \int_\mathbb{R} |\partial_x E^\varepsilon|^2 |E^\varepsilon|^2, \quad (3.7)$$

$$\frac{d}{dt} |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2 + 2 \int_\Omega \hat{\nu}^\varepsilon |\partial_\xi \hat{E}^\varepsilon|^2 + 2 \int_\Omega \partial_\xi \hat{\nu}^\varepsilon \hat{E}^\varepsilon \partial_\xi \overline{\hat{E}^\varepsilon} = 2 \int_\mathbb{R} \left( \partial_\xi \hat{E}^\varepsilon * \hat{E} * \overline{\hat{E}^\varepsilon} \right) \partial_\xi \overline{\hat{E}^\varepsilon}, \quad (3.8)$$

$$\frac{d}{dt} |\partial_\xi H_e^\varepsilon|_{L^2(\Omega)}^2 + 2 \int_\Omega \xi^2 (|\xi|^3 |\hat{E}^\varepsilon|^2 + \varepsilon) |\partial_\xi^2 H_e^\varepsilon|^2 \leq \int \xi^3 |\hat{E}^\varepsilon|^2 |\partial_\xi H_e^\varepsilon|^2 + |\partial_\xi^2 H_e^\varepsilon|_{L^2}^2 |\partial_\xi \hat{E}^\varepsilon|_{L^2}^2. \quad (3.9)$$

The different terms in (3.8) are controlled in the following way :

$$- \int_\Omega \partial_\xi \hat{\nu}^\varepsilon \hat{E}^\varepsilon \partial_\xi \overline{\hat{E}^\varepsilon} \leq \int_\Omega |\xi|^5 |\hat{E}^\varepsilon|^2 |\partial_\xi^2 \hat{H}_e^\varepsilon|^2 + |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2, \quad (3.10)$$

since  $\hat{\nu}^\varepsilon(\xi) = \text{sgn}(\xi) \partial_\xi H_e^\varepsilon$ , while the convolution is estimated as usual by

$$\int_\mathbb{R} \left( \partial_\xi \hat{E}^\varepsilon * \hat{E} * \overline{\hat{E}^\varepsilon} \right) \partial_\xi \overline{\hat{E}^\varepsilon} \leq |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2 |\hat{E}^\varepsilon|_{L^1(\mathbb{R})}^2 \leq c |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2 |E^\varepsilon|_{H^1(\mathbb{R})}^2. \quad (3.11)$$

We remark that the estimate (3.10) is uniform in  $\varepsilon$  thanks to the inequality (3.9).

Without an assumption on the sign of  $\hat{\nu}_e^\varepsilon$ , we get

$$- \int_\Omega \hat{\nu}^\varepsilon |\partial_\xi \hat{E}^\varepsilon|^2 \leq c |\hat{\nu}^\varepsilon|_{H^1(\Omega)} |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2.$$

The latter term needs to establish a control in  $H^2(\Omega)$  in order to obtain uniform estimates in  $\varepsilon$  of  $\hat{\nu}^\varepsilon$ . We obtain

$$\begin{aligned} & \frac{d}{dt} |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^2 + 2 \int_\Omega \hat{\nu}^\varepsilon |\partial_\xi^2 \hat{E}^\varepsilon|^2 + 4 \int_\Omega \partial_\xi \hat{\nu}^\varepsilon \partial_\xi \hat{E}^\varepsilon \partial_\xi^2 \overline{\hat{E}^\varepsilon} \\ & + 2 \int_\Omega \partial_\xi^2 \hat{\nu}^\varepsilon \hat{E}^\varepsilon \partial_\xi^2 \overline{\hat{E}^\varepsilon} = 2 \int_\Omega (\partial_\xi \hat{E}^\varepsilon * \partial_\xi \hat{E}^\varepsilon * \overline{\hat{E}^\varepsilon}) \partial_\xi^2 \overline{\hat{E}^\varepsilon} + 2 \int_\Omega \partial_\xi^2 \hat{S} \partial_\xi^2 \overline{\hat{E}^\varepsilon}, \end{aligned} \quad (3.12)$$

and

$$\begin{aligned} & \frac{d}{dt} |\partial_\xi^2 H_e^\varepsilon|_{L^2(\Omega)}^2 + 2 \int_\Omega \xi^2 (|\xi|^3 |\hat{E}|^2 + \varepsilon) |\partial_\xi^3 H_e^\varepsilon|^2 + |\partial_\xi^2 H_e^\varepsilon|_{L^2(\Omega)}^2 \leq \\ & 4 \int_\Omega |(1 + \partial_\xi)(|\xi|^3 |\hat{E}|^2 + \varepsilon)| |\partial_\xi^2 H_e^\varepsilon| |\partial_\xi^3 H_e^\varepsilon| + 4 \int_\Omega |\partial_\xi (|\xi|^3 |\hat{E}|^2 + \varepsilon)| |\partial_\xi H_e^\varepsilon| |\partial_\xi^3 H_e^\varepsilon| \\ & + |\partial_\xi^2 H_{e0}|_{L^2(\Omega)}^2. \end{aligned} \quad (3.13)$$

The control of the various terms of (3.12) is done in the following way :

$$\int_\Omega \hat{\nu}^\varepsilon |\partial_\xi^2 \hat{E}^\varepsilon|^2 \leq c |\partial_\xi H_e^\varepsilon|_{H^1(\Omega)} |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^2,$$

since  $\hat{\nu}^\varepsilon(\xi) = \text{sgn}(\xi) \partial_\xi H_e^\varepsilon$ . Moreover

$$\int_\Omega \partial_\xi \hat{\nu}^\varepsilon \partial_\xi \hat{E}^\varepsilon \partial_\xi^2 \overline{\hat{E}^\varepsilon} \leq |\partial_\xi^2 H_e^\varepsilon|_{L^2(\Omega)}^2 + c |\partial_\xi \hat{E}^\varepsilon|_{H^1(\Omega)}^2 |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^2,$$

and

$$\int_\Omega \partial_\xi^2 \hat{\nu}^\varepsilon \hat{E}^\varepsilon \partial_\xi^2 \overline{\hat{E}^\varepsilon} \leq \int_\Omega |\xi|^5 |\hat{E}^\varepsilon|^2 |\partial_\xi^3 H_e^\varepsilon|^2 + |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^2.$$

In order to control the last term of (3.12), we use the fact that the domain  $\Omega$  is bounded :

$$\begin{aligned} & \int_\Omega (\partial_\xi \hat{E}^\varepsilon * \partial_\xi \hat{E}^\varepsilon * \overline{\hat{E}^\varepsilon}) \partial_\xi^2 \overline{\hat{E}^\varepsilon} \leq |\partial_\xi \hat{E}^\varepsilon * \partial_\xi \hat{E}^\varepsilon * \overline{\hat{E}^\varepsilon}|_{L^2(\Omega)} |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)} \\ & \leq |\Omega|^{\frac{1}{2}} |\partial_\xi \hat{E}^\varepsilon * \partial_\xi \hat{E}^\varepsilon * \overline{\hat{E}^\varepsilon}|_{L^\infty(\mathbb{R})} |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)} \\ & \leq |\Omega|^{\frac{1}{2}} |\partial_\xi \hat{E}^\varepsilon|_{L^2(\mathbb{R})}^2 |\hat{E}^\varepsilon|_{L^1(\mathbb{R})} |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)} \end{aligned}$$

Moreover,

$$|\hat{E}^\varepsilon|_{L^1(\mathbb{R})} \leq c |E^\varepsilon|_{H^1(\mathbb{R})}.$$

This finishes the control of the terms of (3.12).

Using (3.13) it follows that,

$$\begin{aligned} & \int_\Omega |(1 + \partial_\xi)(|\xi|^3 |\hat{E}|^2)| |\partial_\xi^2 H_e^\varepsilon| |\partial_\xi^3 H_e^\varepsilon| + \int_\Omega |\partial_\xi (|\xi|^3 |\hat{E}|^2)| |\partial_\xi H_e^\varepsilon| |\partial_\xi^3 H_e^\varepsilon|, \\ & \leq 8 \int_\Omega (|\hat{E}^\varepsilon| + |\partial \hat{E}^\varepsilon|) (|\partial_\xi H_e^\varepsilon| + |\partial_\xi^2 H_e^\varepsilon|) (|\hat{E}^\varepsilon| |\partial_\xi^3 H_e^\varepsilon|), \\ & \leq c |\hat{E}^\varepsilon|_{H^2(\Omega)}^4 + |H_e^\varepsilon|_{H^2(\Omega)}^4 + \frac{1}{4} \int_\Omega |\xi|^5 |\hat{E}^\varepsilon|^2 |\partial_\xi^3 H_e^\varepsilon|^2. \end{aligned}$$

Thus, summing the various estimates we have obtained, we get

$$\begin{aligned} & \frac{d}{dt} \left( |E^\varepsilon|_{H^1(\mathbb{R})}^2 + |\hat{E}^\varepsilon|_{H^1(\mathbb{R})}^2 + |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^2 + |\xi^{-1} H_e^\varepsilon|_{L^2(\Omega)}^2 + |\partial_\xi H_e^\varepsilon|_{H^1(\Omega)}^2 \right), \\ & + \int_\Omega (|\hat{E}^\varepsilon|^2 + \varepsilon) (|\partial_\xi^3 H_e^\varepsilon|^2 + |\partial_\xi^2 H_e^\varepsilon|^2 + |\partial_\xi H_e^\varepsilon|^2), \\ & \leq c \left( |\hat{E}^\varepsilon|_{H^1(\mathbb{R})}^4 + |\partial_\xi^2 \hat{E}^\varepsilon|_{L^2(\Omega)}^4 + |H_e^\varepsilon|_{H^2(\Omega)}^4 + |E^\varepsilon|_{H^1(\mathbb{R})}^4 + |S|_{H^1(\mathbb{R})}^2 + |\hat{S}|_{H^1(\mathbb{R})}^2 + |\partial_\xi^2 \hat{S}|_{L^2(\Omega)}^2 \right). \end{aligned}$$

One easily deduces the estimate (3.6). ■

Passing to the limit  $\varepsilon$  to 0, we obtain the local existence result of Theorem 1.1.

We still have to prove the uniqueness result.

Let  $(E_1, \hat{E}_1, H_{e1}, \nu_1)$  and  $(E_2, \hat{E}_2, H_{e2}, \nu_2)$  be two solutions of (3.1) and denote by  $(E, \hat{E}, H_e, \nu)$  the difference of these solutions. Then  $(E, \hat{E}, H_e, \nu)$  satisfies :

$$\begin{aligned} i(\partial_t E + \nu_1 * E + \nu * E_2) + \partial_x^2 E &= |E_1|^2 E + (E_1 \overline{E} + E \overline{E_2}) E_2, \\ \partial_t H_e - \frac{1}{2} \xi^2 \partial_\xi (|\xi|^3 |\hat{E}_2|^2 \partial_\xi H_e + |\xi|^3 (|\hat{E}_1|^2 - |\hat{E}_2|^2) \partial_\xi H_{e1}) \\ - \frac{1}{2} \xi^2 \partial_\xi (|\xi|^3 |\hat{E}_1|^2 \partial_\xi H_e + |\xi|^3 (|\hat{E}_2|^2 - |\hat{E}_1|^2) \partial_\xi H_{e2}) &= 0, \quad \forall \xi \in \Omega, \\ \partial_\xi H_e|_{\partial\Omega} &= 0, \\ H_e(0, \cdot) = 0, E(0, \cdot) &= 0. \end{aligned} \tag{3.14}$$

Performing an  $L^2$  estimate of  $(E, \hat{E}, H_e, \nu)$  yields

$$\begin{aligned} \frac{d}{dt} |\hat{E}|_{L^2(\mathbb{R})}^2 &\leq 2 \int_{\Omega} |\hat{\nu}_1| |\hat{E}|^2 + |\hat{\nu}| |\hat{E}_2| |\hat{E}| + |E_1|_{L^\infty(\mathbb{R})} |\hat{E}|_{L^2(\mathbb{R})}^2 \leq C |\hat{E}|_{L^2(\mathbb{R})}^2 + \int_{\Omega} |\hat{E}_2|^2 |\hat{\nu}|^2, \\ \frac{d}{dt} |\xi^{-1} H_e|_{L^2(\mathbb{R})}^2 + \int_{\Omega} \xi^3 (|\hat{E}_1|^2 + |\hat{E}_2|^2) |\partial_\xi H_e|^2 + 2 |H_e|_{L^2(\Omega)}^2 &\leq, \\ \int_{\Omega} \xi^3 (|\hat{E}_1|^2 - |\hat{E}_2|^2) (|\partial_\xi H_{e1}| + |\partial_\xi H_{e2}|) |\partial_\xi H_e|, \\ &\leq C |\hat{E}|_{L^2(\Omega)}^2 + \frac{1}{2} \int_{\Omega} \xi^3 (|\hat{E}_1|^2 + |\hat{E}_2|^2) |\partial_\xi H_e|^2. \end{aligned}$$

Thus, by making the sum of these estimates, one gets

$$\frac{d}{dt} \left( |\hat{E}|_{L^2(\mathbb{R})}^2 + |\xi^{-1} H_e|_{L^2(\mathbb{R})}^2 \right) + \int_{\Omega} \xi^3 |\hat{E}_2|^2 |\partial_\xi H_e|^2 \leq C |\hat{E}|_{L^2(\Omega)}^2,$$

where  $C$  depends on the solutions  $(E_i, \hat{E}_i, H_{ei})$  ( $i = 1, 2$ ), in the norm of  $H^1(\mathbb{R}) \times H^2(\Omega) \cap H^1(\mathbb{R}) \times H^2(\Omega)$ .

The uniqueness result is then straightforward. ■

### 3.2 Global existence of strong solutions for the homogeneous problem

The aim of this section is to prove Theorem 1.2. We just perform a priori estimates (these estimates can be done on the approximate solutions constructed in the preceding section for example).

• **Step 1:** Maximum principle.

**Proposition 3.4.** *If  $H$  is a solution to*

$$\partial_t H - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 \partial_\xi H) = 0, \quad \forall \xi \in \Omega,$$

*and  $\text{sgn}(\xi) \partial_\xi H(0, \xi) \geq 0$ , then for all  $t \geq 0$ ,  $\xi \in \Omega$*

$$\hat{\nu}(t, \xi) = \text{sgn}(\xi) \partial_\xi H(t, \xi) \geq 0.$$

Proof : the equation satisfied by  $\widehat{\nu}$  is

$$\partial_t \widehat{\nu} - \xi^2 \partial_\xi \left( |\xi|^3 |\widehat{E}|^2 \partial_\xi \widehat{\nu} \right) = \xi^2 \partial_\xi \left( (\partial_\xi (\xi^3 |\widehat{E}|^2) \widehat{\nu} \right) + 2\xi \partial_\xi (\xi^3 |\widehat{E}|^2 \widehat{\nu}).$$

By substituting  $\widehat{\nu}$  by  $\widehat{\nu} = \widehat{\nu}^+ - \widehat{\nu}^-$ , where  $\widehat{\nu}^+$ ,  $\widehat{\nu}^-$  are the positive part and negative part of  $\widehat{\nu}$  respectively, multiplying the equation by  $-\xi^{-2} \widehat{\nu}^-$  and integrating on  $\Omega$  yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |\xi^{-1} \widehat{\nu}^-|_{L^2}^2 + \int \xi^3 |\widehat{E}|^2 |\partial_\xi \widehat{\nu}^-|^2 d\xi &= \int \widehat{\nu}^- \partial_\xi \widehat{\nu}^- \partial_\xi (\xi^3 |\widehat{E}|^2) d\xi \\ &\quad + 2 \int \widehat{\nu}^- \partial_\xi (\xi^{-1} \widehat{\nu}^-) (\xi^3 |\widehat{E}|^2) d\xi. \end{aligned}$$

This gives

$$\frac{1}{2} \frac{d}{dt} |\xi^{-1} \widehat{\nu}^-|_{L^2}^2 + \int \xi^3 |\widehat{E}|^2 |\partial_\xi \widehat{\nu}^-|^2 d\xi \leq C(\Omega) \max(|\widehat{E}|_{L^\infty}, |\partial_\xi \widehat{E}|_{L^\infty}) |\widehat{\nu}^-|_{L^2}^2.$$

Then it follows

$$\frac{d}{dt} |\widehat{\nu}^-|_{L^2}^2 \leq \alpha(\Omega) |\widehat{\nu}^-|_{L^2}^2.$$

According to the Gronwall lemma with  $\widehat{\nu}^-(., 0) = 0$ , one gets the result. ■

• **Step 2 :** We establish a  $L_{loc}^\infty(\mathbb{R}^+, H^1)$  bound for  $\widehat{E}$ .

If  $E$  satisfies

$$i(\partial_t E + \nu * E) + \partial_x^2 E = 0,$$

taking the Fourier transform yields

$$i \left( \partial_t \widehat{E} + \widehat{\nu} \widehat{E} \right) - \xi^2 \widehat{E} = 0. \quad (3.15)$$

Since  $\widehat{\nu} \geq 0$ , we have

$$|\widehat{E}|_{L^2}(t) \leq |\widehat{E}|_{L^2}(0).$$

Applying  $\partial_\xi$  on (3.15), multiplying by  $\partial_\xi \overline{\widehat{E}}$  and taking the imaginary part yields

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi \widehat{E}|^2 + \int \widehat{\nu} |\partial_\xi \widehat{E}|^2 + \Re \int \partial_\xi \widehat{\nu} \widehat{E} \partial_\xi \overline{\widehat{E}} = 0,$$

which implies

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi \widehat{E}|^2 \leq \alpha \int |\partial_\xi \widehat{\nu}|^2 |\widehat{E}|^2 + \frac{1}{\alpha} |\partial_\xi \widehat{E}|^2, \quad (3.16)$$

since  $\int \widehat{\nu} |\partial_\xi \widehat{E}|^2 \geq 0$  and  $\alpha$  will be fixed latter one.

Remark that the term  $\int |\partial_\xi \widehat{\nu}|^2 |\widehat{E}|^2$  will be controlled thanks to the dissipative term on the equation on  $H$ .

• **Step 3:**  $L_{loc}^\infty(\mathbb{R}^+, L^2)$  estimate for  $H$ .

Multiplying (1.14) by  $\frac{H}{\xi^2}$  and integrating on  $\Omega$  gives

$$\frac{1}{2} \frac{d}{dt} \int \frac{H^2}{\xi^2} + \int \xi^3 |\widehat{E}|^2 |\partial_\xi H|^2 = 0. \quad (3.17)$$

• **Step 4:**  $L_{loc}^\infty(\mathbb{R}^+, H^1)$  estimate for  $H$ .

We apply  $\partial_\xi$  on (1.14) and multiply the result by  $\partial_\xi H$ . This leads to

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 - \int \partial_\xi \left( \xi^2 \partial_\xi \left( \xi^3 |\widehat{E}|^2 \partial_\xi H \right) \right) \partial_\xi H = 0, \quad (3.18)$$

that is

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 + \int \xi^2 \partial_\xi \left( \xi^3 |\hat{E}|^2 \partial_\xi H \right) \partial_\xi^2 H = 0,$$

or

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 + \int \xi^5 |\hat{E}|^2 |\partial_\xi^2 H|^2 + \int \xi^2 \partial_\xi \left( \xi^3 |\hat{E}|^2 \right) \partial_\xi H \partial_\xi^2 H = 0.$$

The crucial point is to deal with the term  $\int \xi^2 \partial_\xi \left( \xi^3 |\hat{E}|^2 \right) \partial_\xi H \partial_\xi^2 H$ . We strongly use the structure of the system (1.13)-(1.14).

Taking the Fourier transform of (1.13), multiplying by  $\overline{\hat{E}}$  and taking the imaginary part gives

$$\frac{1}{2} \frac{d}{dt} |\hat{E}|^2 + \partial_\xi H |\hat{E}|^2 = 0. \quad (3.19)$$

We consider  $\xi^2 \partial_\xi \left( \xi^3 (3.17) \right)$  and obtain

$$\frac{1}{2} \frac{d}{dt} \left( \xi^2 \partial_\xi \left( \xi^3 (|\hat{E}|^2) \right) \right) + \xi^2 \partial_\xi \left( \xi^3 \partial_\xi H |\hat{E}|^2 \right) = 0, \quad (3.20)$$

and using (1.14), one gets

$$\frac{1}{2} \frac{d}{dt} \left( \xi^2 \partial_\xi \left( \xi^3 (|\hat{E}|^2) \right) \right) + \partial_t H = 0. \quad (3.21)$$

It follows that

$$\xi^2 \partial_\xi \left( \xi^3 (|\hat{E}|^2) \right) + 2H = f_0, \quad (3.22)$$

where  $f_0 = \xi^2 \partial_\xi \left( \xi^3 (|\hat{E}_0|^2) \right) + 2H_0$ .

Plugging (3.22) in (3.18) leads to

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 + \int \xi^5 |\hat{E}|^2 |\partial_\xi^2 H|^2 - 2 \int H \partial_\xi H \partial_\xi^2 H + \int f_0 \partial_\xi H \partial_\xi^2 H = 0.$$

Integration by parts gives

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 + \int \xi^5 |\hat{E}|^2 |\partial_\xi^2 H|^2 + \int (\partial_\xi H)^3 - \frac{1}{2} \int \partial_\xi f_0 |\partial_\xi H|^2 = 0. \quad (3.23)$$

Recall now that thanks to step 1,  $\partial_\xi H \geq 0$ , and one gets

$$\frac{1}{2} \frac{d}{dt} \int |\partial_\xi H|^2 + \int \xi^5 |\hat{E}|^2 |\partial_\xi^2 H|^2 \leq \frac{1}{2} \int (\partial_\xi f_0)^+ |\partial_\xi H|^2. \quad (3.24)$$

Inequalities (3.16)-(3.24) give

$$\frac{d}{dt} \left( |\hat{E}|^2 + |\partial_\xi H|^2 \right) + \int \xi^5 |\hat{E}|^2 |\partial_\xi^2 H|^2 \leq \alpha \int |\partial_\xi \hat{v}|^2 |\hat{E}|^2 + \frac{1}{\alpha} |\partial_\xi \hat{E}|^2 + \frac{1}{2} \int (\partial_\xi f_0)^+ |\partial_\xi H|^2.$$

Since  $\xi \in [1, 2]$ , we can choose  $\alpha = 1$ . One gets bounds for  $\hat{E}, H$  in  $L_{loc}^\infty(\mathbb{R}^+, H^1)$  and the result follows.  $\blacksquare$

## 4 Numerical approximation.

In this section, using a time-splitting spectral discretization for the Zakharov system and a finite difference scheme for the electron diffusion equation, we present numerical simulations and we show how Landau damping works.

We consider the full system with periodic boundary conditions :

$$i(\partial_t E + \nu * E) + \partial_x^2 E = \delta n E + E_p(x) e^{i(k_1 x - \omega_1 t)}, \quad 0 < x < L, \quad t > 0, \quad (4.1)$$

$$\partial_t^2 \delta n - \mu \partial_x^2 \delta n = \mu \partial_x^2 (|E|^2), \quad 0 < x < L, \quad t > 0, \quad (4.2)$$

$$\hat{\nu}(t, \xi) = \text{sgn}(\xi) \partial_\xi H_e(t, \xi), \quad \xi \in \Omega, \quad t > 0, \quad (4.3)$$

$$\partial_t H_e - \xi^2 \partial_\xi (|\xi|^3 |\hat{E}|^2 \partial_\xi H_e) = 0, \quad \xi \in \Omega, \quad (4.4)$$

$$E(t, 0) = E(t, L), \quad t > 0, \quad (4.5)$$

$$\delta n(t, 0) = \delta n(t, L), \quad t > 0. \quad (4.6)$$

System (4.1)-(4.4) is endowed with the following initial conditions :

$$E(0, x) = E_0(x), \quad \delta n(0, x) = \delta n_0(x), \quad \partial_t \delta n(0, x) = \delta n_1(x), \quad H_e(0, \xi) = H_{e,0}(\xi).$$

Moreover we suppose that

$$E_0(0) = E_0(L), \quad \delta n_0(0) = \delta n_0(L), \quad \delta n_1(0) = \delta n_1(L).$$

It is easy to show that the above system satisfies the following

**Proposition 4.1.** *If  $\int_0^L \delta n_1(x) dx = 0$ , any regular solution of (4.1)-(4.6) satisfies :*

$$\int_0^L \delta n(t, x) dx = \int_0^L \delta n_0(x) dx, \quad \forall t \geq 0, \quad (4.7)$$

$$\int \frac{1}{\xi^2} H_e(t, \xi) d\xi = \int \frac{1}{\xi^2} H_e(0, \xi) d\xi, \quad \forall t \geq 0 \quad (4.8)$$

Moreover if  $E_p = 0$ , then

$$\frac{1}{2} \frac{d}{dt} \int \frac{1}{\xi^4} H_e(t, \xi) d\xi = \int \hat{\nu}(t, \xi) |\hat{E}|^2 d\xi, \quad \forall t \geq 0. \quad (4.9)$$

Below, we construct a numerical approximation of (4.1)-(4.4) which is consistent with the above properties.

We choose the spatial mesh size  $h = \Delta x$  with  $h = L/N$  for  $N = 2M$  being an even number, the time step being  $\Delta t > 0$  and let the grid points and the time step be

$$x_j = jh, \quad t_k = k\Delta t, \quad j = 1, \dots, N, \quad k = 0, 1, 2, \dots$$

We use the sequence  $(\xi_j = \frac{2\pi j}{L})_{j=-\frac{N}{2}, \dots, 0, \dots, \frac{N}{2}-1}$  as the regular mesh grid in frequency with  $\Delta \xi = \frac{2\pi}{L}$ . Furthermore, let  $E_j^k$ ,  $\delta n_j^k$ , and  $\hat{\nu}_j^k$  be the approximations of  $E(t_k, x_j)$ ,  $\delta n(t_k, x_j)$ ,  $\hat{\nu}(t_k, \xi_j)$ . Then, in order to be consistent with the evaluation of  $\hat{\nu}(t_k, \xi_j) = \text{sgn}(\xi_j) \frac{\pi}{2} \partial_\xi H(t_k, \xi_j)$ , we approximate  $H_e$  on the grid  $(\xi_{j+\frac{1}{2}})_j$  defined by  $\xi_{j+\frac{1}{2}} = \frac{2\pi(j+1/2)}{L}$ .

#### 4.1 Finite difference scheme for the diffusion equation.

In order to evaluate the approximation of  $H(t_k, \xi_{j+\frac{1}{2}})$ , we use an implicit difference scheme for the diffusion equation :

$$\partial_t H_e - \xi^2 \partial_\xi \left( |\xi|^3 |\hat{E}|^2 \partial_\xi H_e \right) = 0, \quad \xi \in \Omega,$$

where  $\Omega = [-\xi_2, -\xi_1] \cup [\xi_1, \xi_2]$ , ( $\xi_2 > \xi_1 > 0$ ), with  $\xi_1 = \frac{2\pi(j_1+1/2)}{L} > \frac{2\pi}{L}$ ,  $\xi_2 = \frac{2\pi(j_2+1/2)}{L} < \frac{2\pi(M-1)}{L}$ .

The initial electron distribution function is assumed to be a Maxwellian distribution :

$$H_{e,0}(\xi) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2\xi^2}\right).$$

The boundary conditions are

$$\begin{aligned} \partial_\xi H_e(., \pm \xi_1) &= 0, \\ H_e(., \xi_2) &= H_e(., -\xi_2). \end{aligned}$$

The scheme reads :

$$\frac{1}{\Delta t} (H_{j+\frac{1}{2}}^{k+1} - H_{j+\frac{1}{2}}^k) + (AH)_{j+\frac{1}{2}}^{k+1} = 0, \quad (4.10)$$

where  $(AH)_{j+\frac{1}{2}}^{k+1}$  is a discretization of  $-\xi^2 \partial_\xi (\beta(t, \xi) \partial_\xi H_e)$  in a conservative form at the point  $\xi_{j+\frac{1}{2}}$  and time  $t_{k+1}$ . Here  $\beta(t, \xi) = |\xi|^3 |\hat{E}|^2$ . We choose  $A$  such that :

$$(AH)_{j+\frac{1}{2}}^k = -\frac{\xi_{j+\frac{1}{2}}^2}{\xi_{j+1} - \xi_j} \left[ \beta_{j+1}^{k-1} \frac{H_{j+\frac{3}{2}}^k - H_{j+\frac{1}{2}}^k}{\xi_{j+\frac{3}{2}} - \xi_{j+\frac{1}{2}}} - \beta_j^{k-1} \frac{H_{j+\frac{1}{2}}^k - H_{j-\frac{1}{2}}^k}{\xi_{j+\frac{1}{2}} - \xi_{j-\frac{1}{2}}} \right], \quad (4.11)$$

where  $\beta_j^k$  is the discretization of  $\beta(t, \xi)$  given in proposition 4.2. Then we can evaluate  $\hat{\nu}(\xi_j, t^{k+1})$  with the centered finite difference scheme :

$$\hat{\nu}_j^{k+1} = \text{sgn}(\xi_j) \frac{H_{j+\frac{1}{2}}^{k+1} - H_{j-\frac{1}{2}}^{k+1}}{\Delta \xi}. \quad (4.12)$$

The next proposition shows that there exists some approximate values of  $\beta(t, \xi)$  such that one obtains discrete equivalent of (4.8)-(4.9).

**Proposition 4.2.** (*Energy and density conservation*). *Take*

$$\beta_l^k = \frac{\xi_{l+\frac{1}{2}}^2 \xi_{l-\frac{1}{2}}^2}{|\xi_{l+\frac{1}{2}} + \xi_{l-\frac{1}{2}}|} |\hat{E}_l^k|^2.$$

*Then the following identities hold*

$$\sum_l \frac{1}{\xi_{l+\frac{1}{2}}^2} H_{l+\frac{1}{2}}^{k+1} = \sum_l \frac{1}{\xi_{l+\frac{1}{2}}^2} H_{l+\frac{1}{2}}^k, \quad (4.13)$$

$$\sum_l \frac{1}{\xi_{l+\frac{1}{2}}^4} H_{l+\frac{1}{2}}^{k+1} = \sum_l \frac{1}{\xi_{l+\frac{1}{2}}^4} H_{l+\frac{1}{2}}^k + \Delta t \sum_l \hat{\nu}_l^{k+1} |\hat{E}_l^k|^2. \quad (4.14)$$

Proof : The first identity follows from the conservativity of the discretization of  $\partial_\xi (\beta(\xi, t) \partial_\xi H_e)$  given by (4.11).

Below, we will prove the second conservation law.

According to the scheme (4.10)-(4.12), we have

$$\frac{H_{j+\frac{1}{2}}^{k+1}}{\xi_{j+\frac{1}{2}}^4} = \frac{H_{j+\frac{1}{2}}^k}{\xi_{j+\frac{1}{2}}^4} + \Delta t \left( \frac{\beta_{j+1}^k}{\xi_{j+\frac{1}{2}}^2} \frac{H_{j+\frac{3}{2}}^{k+1} - H_{j+\frac{1}{2}}^{k+1}}{\Delta \xi^2} - \frac{\beta_j^k}{\xi_{j+\frac{1}{2}}^2} \frac{H_{j+\frac{1}{2}}^{k+1} - H_{j-\frac{1}{2}}^{k+1}}{\Delta \xi^2} \right).$$

Taking into account boundary conditions, it follows that

$$\sum_l \frac{H_{l+\frac{1}{2}}^{k+1}}{\xi_{l+\frac{1}{2}}^4} = \sum_l \frac{H_{l+\frac{1}{2}}^k}{\xi_{l+\frac{1}{2}}^4} + \Delta t \left( \sum_l \beta_l^k \left( \frac{1}{\xi_{l-\frac{1}{2}}^2} - \frac{1}{\xi_{l+\frac{1}{2}}^2} \right) \frac{H_{l+\frac{1}{2}}^{k+1} - H_{l-\frac{1}{2}}^{k+1}}{\Delta \xi^2} \right). \quad (4.15)$$

Plugging (4.12) in (4.15) yields

$$\sum_l \frac{H_{l+\frac{1}{2}}^{k+1}}{\xi_{l+\frac{1}{2}}^4} = \sum_l \frac{H_{l+\frac{1}{2}}^k}{\xi_{l+\frac{1}{2}}^4} + \Delta t \left( \sum_l \beta_l^k \operatorname{sgn}(\xi_l) \left( \frac{\xi_{l+\frac{1}{2}}^2 \xi_{l-\frac{1}{2}}^2}{\xi_{l+\frac{1}{2}} + \xi_{l-\frac{1}{2}}} \right)^{-1} \widehat{\nu}_l^{k+1} \right), \quad (4.16)$$

and with the choice of discretization of  $\beta(t, \xi)$ , the result follows. ■

**Proposition 4.3.** (*Maximum principle*). *Let  $\widehat{\nu}_0 = \widehat{\nu}(t=0)$ . If  $\widehat{\nu}_0$  satisfies*

$$\widehat{\nu}_0(\xi_l) \geq 0, \quad l = 0, \dots, N-1,$$

*then for all  $k > 0$*

$$\widehat{\nu}_l^k \geq 0, \quad l = 0, \dots, N-1.$$

Proof : Here, we restrict ourselves to the case where the sequence  $(\xi_l)_l$  belongs to  $[\xi_1, \xi_2]$ . The discretization of  $\widehat{\nu}$  is given by

$$\widehat{\nu}_l^{k+1} = \operatorname{sgn}(\xi_l) \frac{H_{l+\frac{1}{2}}^{k+1} - H_{l-\frac{1}{2}}^{k+1}}{\Delta \xi}.$$

The implicit difference scheme used for  $H$  implies that the sequence  $(\widehat{\nu}_j^k)_{(j,k)}$  satisfies

$$-\frac{\Delta t}{\Delta \xi^2} \xi_{l+\frac{1}{2}}^2 \beta_{l+1}^k \widehat{\nu}_{l+1}^{k+1} + \left(1 + \frac{\Delta t}{\Delta \xi^2} \beta_l^k (\xi_{l+\frac{1}{2}}^2 + \xi_{l-\frac{1}{2}}^2)\right) \widehat{\nu}_l^{k+1} - \frac{\Delta t}{\Delta \xi^2} \xi_{l-\frac{1}{2}}^2 \beta_{l-1}^k \widehat{\nu}_{l-1}^{k+1} = \widehat{\nu}_l^k. \quad (4.17)$$

Suppose that  $\widehat{\nu}_l^k \geq 0$  for all  $l$  and introduce the sequence  $\mu_l^{k+1}$  such that  $\mu_l^{k+1} = \beta_l^k \widehat{\nu}_l^{k+1}$  and  $l_0$  such that  $\mu_{l_0}^{k+1} = \min_l(\mu_l^{k+1})$ .

Firstly, suppose that  $\mu_{l_0}^{k+1} < 0$  then  $\beta_{l_0}^k \neq 0$ . It follows that

$$\left( \frac{1}{\beta_{l_0}^k} + \frac{\Delta t}{\Delta \xi^2} (\xi_{l_0+\frac{1}{2}}^2 + \xi_{l_0-\frac{1}{2}}^2) \right) \mu_{l_0}^{k+1} = \widehat{\nu}_{l_0}^k + \frac{\Delta t}{\Delta \xi^2} \xi_{l_0+\frac{1}{2}}^2 \mu_{l_0+1}^{k+1} + \frac{\Delta t}{\Delta \xi^2} \xi_{l_0-\frac{1}{2}}^2 \mu_{l_0-1}^{k+1}. \quad (4.18)$$



Since  $\mu_{l_0+1}^{k+1} \geq \mu_{l_0}^{k+1}$  and  $\mu_{l_0-1}^{k+1} \geq \mu_{l_0}^{k+1}$ , (4.18) implies that

$$\frac{\mu_{l_0}^{k+1}}{\beta_{l_0}^k} \geq \widehat{\nu}_{l_0}^k \geq 0,$$

that is  $\mu_{l_0}^{k+1}$  is positive which is a contradiction. So the sequence  $(\mu_l^k)_l$  is positive for all  $l$ . Therefore the sequence  $(\widehat{\nu}_l^{k+1})_l$  is positive for  $l$  not included in the set  $J = \{j \text{ s.t. } \beta_j^k = 0\}$ . Let be  $j$  such that  $\beta_j^k = 0$ , then for all  $k \geq 0$

$$\widehat{\nu}_j^{k+1} = \widehat{\nu}_j^k + \frac{\Delta t}{\Delta \xi^2} \xi_{j+\frac{1}{2}}^2 \mu_{j+1}^{k+1} + \frac{\Delta t}{\Delta \xi^2} \xi_{j-\frac{1}{2}}^2 \mu_{j-1}^{k+1}, \quad (4.19)$$

which shows that  $\widehat{\nu}_j^{k+1}$  is positive and the result follows.  $\blacksquare$

## 4.2 Time-splitting spectral discretizations for the Zakharov system.

Many numerical methods have been proposed for the Zakharov part of the system. For example Payne *et al.* [15] have introduced a spectral method. They used a truncated Fourier expansion in their algorithm to eliminate aliasing errors. Glassey [9] presented an energy-preserving finite difference scheme for the Zakharov system in one dimension. Here we describe a time-splitting spectral discretization. The first equation (4.1) is solved in two splitting steps. One first solves

$$i(\partial_t E + \nu * E) + \partial_x^2 E = E_p(x) e^{i(k_1 x - \omega_1 t)}, \quad (4.20)$$

for the time step of length  $\Delta t$ . In the second step, one solves

$$i\partial_t E = \frac{1}{2} \delta n E, \quad (4.21)$$

$$\partial_t^2 \delta n - \mu \partial_x^2 \delta n = \mu \partial_x^2 (|E|^2), \quad (4.22)$$

$$E_0 = E^*(.), \quad (4.23)$$

$$\delta n_0 = \delta n(t^k, .), \quad (4.24)$$

$$\partial_t \delta n_0 = \partial_t \delta n(t^k, .). \quad (4.25)$$

Equation (4.20) will be discretized in space by the Fourier spectral method and integrated exactly in time. Let  $\overline{E}^*$  be the solution of (4.20) at time  $t = t_{k+1}$ . For  $t \in [t_k, t_{k+1}]$ , multiplying (4.21) by  $\overline{E}$ , the conjugate of  $E$  and taking the imaginary part, we get :

$$\frac{d}{dt} |E(t, x)|^2 = 0.$$

With this conservation law, in order to solve (4.21)-(4.25), we have to solve :

$$i\partial_t E = \frac{1}{2} n E \quad (4.26)$$

$$\partial_t^2 P - \mu \partial_x^2 P = 0 \quad (4.27)$$

$$E_0 = E^*(.) \quad (4.28)$$

$$P_0 = P(t^k, .) \quad (4.29)$$

$$\partial_t P_0 = \partial_t n(t^k, .) = P_1. \quad (4.30)$$

with  $P = \delta n + |E^*|^2$ .

Finally, from  $t = t_k$  to  $t = t_{k+1}$ , using trapezoidal rule, we combine the splitting steps via the standard Strang splitting.

For the first step of the splitting, we compute

$$\begin{aligned}\widehat{E}_l^k &= \frac{1}{N} \sum_{j=1}^N E_j^k e^{-i\xi_l x_j}, \quad l = -M, \dots, M-1, \\ \widehat{E}_l^* &= \widehat{E}_l^k \exp \left( -(\widehat{\nu}_l^k + \widehat{\nu}_l^{k+1}) \frac{\Delta t}{2} + \frac{3i\xi_l^2}{2} \Delta t \right) \\ &+ \frac{\delta t}{2} \left( \widehat{f}_l^k \exp \left( -(\widehat{\nu}_l^k + \widehat{\nu}_l^{k+1}) \frac{\Delta t}{2} + \frac{3i\xi_l^2}{2} \Delta t \right) + \widehat{f}_l^{k+1} \right), \quad l = -M, \dots, M-1, \\ E_j^* &= \sum_{l=-M}^{M-1} \widehat{E}_l^* e^{i\xi_l x_j}, \quad j = 1, \dots, N,\end{aligned}$$

where  $f = E_p(x) e^{i(k_1 x - \omega_1 t)}$  and  $(\widehat{U})_l$  denote the Fourier coefficients of the periodic vector  $U = (U_0, U_1, \dots, U_N)^T$ .

For the second step, we compute

$$\widehat{P}_l^{k+1} = \widehat{P}_0(\xi_l) \cos(\sqrt{\mu} \xi_l \Delta t) + \widehat{P}_1(\xi_l) \frac{\sin(\sqrt{\mu} \xi_l \Delta t)}{\sqrt{\mu} \xi_l}, \quad (4.31)$$

$$\partial_t \widehat{P}_l^{k+1} = \widehat{P}_1(\xi_l) \cos(\sqrt{\mu} \xi_l \Delta t) - \sqrt{\mu} \xi_l \widehat{P}_0(\xi_l) \sin(\sqrt{\mu} \xi_l \Delta t), \quad (4.32)$$

$$P_j^{k+1} = \sum_{l=-M}^{M-1} \widehat{P}_l^{k+1} e^{i\xi_l x_j}, \quad j = 1..N, \quad (4.33)$$

$$\delta n_j(t^{k+1}) = P_j^{k+1} - |E_j^*|^2 \quad j = 1..N, \quad (4.34)$$

$$E_j^{k+1} = E_j^* \exp \left( -\frac{i}{2} \int_{t^k}^{t^{k+1}} \delta n_j(s) ds \right) \simeq E_j^* \exp \left( -\frac{i}{2} (\delta n_j^k + \delta n_j^{k+1}) \frac{\Delta t}{2} \right). \quad (4.35)$$

Now let  $\|\cdot\|_{l^2}$  be the usual discrete  $l^2$  norm on the interval  $(0, L)$  :

$$\|U\|_{l^2} = \sqrt{\frac{L}{N} \sum_{j=1}^L |U_j|^2},$$

then, we have

**Proposition 4.4.** ( $l^2$  stability). *If the initial data  $\widehat{\nu}_0$  and  $n_1(\cdot) = \partial_t \delta n(\cdot, 0)$  satisfy*

$$\widehat{\nu}_0(\xi_j) \geq 0 \quad j = 0, \dots, N-1, \quad \text{and} \quad \int n_1(x) dx = 0$$

*then the time-splitting spectral discretization of the Zakharov system have the following properties :*

$$\left\| E^{k+1} \right\|_{l^2}^2 \leq \left\| E^k \right\|_{l^2}^2 + \Delta t^2 \|E_p\|_{l^2}^2 \quad \forall k > 0, \quad (4.36)$$

$$\frac{b-a}{N} \sum_{j=1}^N \delta n_j^k = \frac{b-a}{N} \sum_{j=1}^N \delta n_j^0 = \frac{b-a}{N} \sum_{j=1}^N \delta n_0(x_j), \quad k = 1, 2, \dots$$

Proof :

1.  $l_2$  estimate (4.36) :

We multiply equation (4.35) by  $\overline{E}^{k+1}$  and obtain :

$$\begin{aligned} \frac{N}{L} \|E^{k+1}\|_{l_2}^2 &= \sum_{j=1}^N |E_j^{k+1}|^2 = \sum_{j=1}^N \left| E_j^* \exp(-i(\delta n_j^k + \delta n_j^{k+1}) \frac{\delta t}{2}) \right|^2, \\ &= \sum_{j=1}^N |E_j^*|^2 = \sum_{j=1}^N \left| \sum_{l=-M}^{M-1} \widehat{E}_l^* e^{i\xi_l x_j} \right|^2, \\ &= \sum_{j=1}^N \left( \sum_{-M \leq l, k \leq M-1} (\widehat{E}^*)_l \overline{(\widehat{E}^*)_k} e^{i(\xi_l - \xi_k)x_j} \right), \\ &= \sum_{-M \leq l, k \leq M-1} \left( \sum_{j=1}^N e^{i(\xi_l - \xi_k)x_j} \right) (\widehat{E}^*)_l \overline{(\widehat{E}^*)_k}. \end{aligned}$$

Then with  $\sum_{j=1}^N e^{i(\xi_l - \xi_k)x_j} = \sum_{j=1}^N e^{\frac{2i\pi}{N}(l-k)(j-1)} = \begin{cases} N & \text{if } k-l = pN \\ 0 & \text{if } k-l \neq pN \end{cases}$ ,

we have

$$\frac{N}{L} \|E^{k+1}\|_{l_2}^2 = N \sum_{l=-M}^{M-1} |(\widehat{E}^*)_l|^2 = N \sum_{l=-M}^{M-1} \left| \exp\left(\frac{3}{2}i\xi_l^2 \delta t - (\hat{\nu}_l^k + \hat{\nu}_l^{k+1}) \frac{\delta t}{2}\right) (\widehat{E}^*)_l \right|^2.$$

Using proposition 4.3, we obtain that

$$\frac{N}{L} \|E^{k+1}\|_{l_2}^2 \leq N \sum_{l=-M}^{M-1} |(\widehat{E}^*)_l|^2,$$

so

$$\frac{N}{L} \|E^{k+1}\|_{l_2}^2 \leq \sum_{j=1}^N |E_j^k|^2 = \frac{N}{L} \|E^k\|_{l_2}^2.$$

2. Low-frequency density conservation.

Since we have the relation

$$\widehat{\delta n}(\xi_0) = \frac{1}{N} \sum_{j=1}^N \delta n_j,$$

with  $\xi_0 = 0$ , it is enough to show that  $\widehat{\delta n}(\xi_0)$  is preserved.

For that, one uses the expressions given previously for  $\widehat{P}_j^{k+1}$  and  $\partial_t \widehat{P}_j^{k+1}$  with  $P_j =$

$$\delta n_j + \frac{1}{4}|E_j^*|^2.$$

Taking (4.31) for  $\xi = \xi_0$  yields

$$\partial_t \widehat{P}^{k+1}(\xi_0) = \partial_t \widehat{P}^k(\xi_0).$$

Then, by using the hypothesis on the initial data  $\delta n_1(\cdot)$ , we have

$$\partial_t \widehat{P}^k(\xi_0) = 0 \quad \forall k > 0.$$

Taking (4.32) for  $\xi = \xi_0$  yields

$$\widehat{P}^{k+1}(\xi_0) = \widehat{P}_0(\xi_0) = \widehat{P}^k(\xi_0),$$

so

$$\widehat{\delta n}^{k+1}(\xi_0) + \frac{1}{4}\mathcal{F}_d(|E^*|^2)(\xi_0) = \widehat{\delta n}^k(\xi_0) + \frac{1}{4}\mathcal{F}_d(|E^*|^2)(\xi_0),$$

where  $\mathcal{F}_d$  denotes the discret Fourier transform.

Therefore we have

$$\sum_{j=1}^N \delta n_j^{k+1} = \sum_{j=1}^N \delta n_j^k$$

and the result follows. ■

### 4.3 Numerical results.

#### 4.3.1 The quasilinear diffusion

In order to illustrate how the quasilinear diffusion works, we computed the diffusion equation with a diffusion coefficient given by a fixed electric field

$$E(x) = e^{ik_1 x} e^{-\frac{(x - \frac{L}{2})^2}{2\beta^2}},$$

with  $L = 2000$ ,  $\beta = 50$  and  $k_1 = 0.45$ .

The initial electron distribution function is assumed to be a Maxwellian,

$$F_{e0}(v) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{v^2}{2}\right).$$

We can remark that the electron function is flattened near the phase velocity  $v_\phi = \frac{1}{k_1} = 2.22$  and since the Landau damping rate depends on the slope of the electron distribution, we can see that  $\widehat{\nu}$  tends toward zero near  $\xi = k_1$ .




Figure 1: The left plot corresponds to the evolution of the electron function distribution and the right plot to the evolution of the Landau damping rate.

#### 4.3.2 The full system

In this part, simulation results using the above quasilinear Zakharov (4.1)-(4.4) model is presented. The mass ratio is  $\sqrt{m_e/m_i} = 1/50$  and the simulation box length is  $L = 2000\lambda_{De}$  (where  $\lambda_{De}$  denotes the Debye length).

The initial electric field and low frequency density in the quasilinear Zakharov simulation are equal to zero. The initial electron distribution function is assumed to be a Maxwellian. The system is driven with an external periodic force localized in the center of the box and we use a 1024 grid point with  $\Delta\xi = 2\pi/L$ .

We present the results for  $k_1\lambda_{De} = 0.09$  and  $|E_p| = 8.10^{-3}$  which corresponds to a weakly driven case.

Figure 1 shows the time history of the spatially averaged Langmuir wave energy :  $W_E = \int |E|^2 dx$ . We can see two stages in this simulation : a linear stage where the external pump excited primary electron plasma wave at  $k_p$  and a nonlinear stage from  $\omega_{pe}t = 3000$  where the Langmuir waves have lost energy. Then the system saturates near  $\omega_{pe}t = 6000$ , due to a combinaison of wave-wave and wave-particle processes. The time history of electrons's energy

$W_e(t) = \int v^2 F_e(v, t) dv$  versus time in Figure 3 shows us that wave-wave saturation and wave-particle saturation occur simultaneously. Moreover we can estimate that the fraction of pump's energy which heats electrons is 5%. This heating implies that hot electrons have been created. Figure 4 shows the spatially averaged electron distribution at the final time of the simulation. The log of the distribution is plotted with respect to the half square of the velocity. We can see a plateau in the electron distribution function with a slope equal to 8. We see significant heating of the electrons : a hot electron tail is produced




Figure 2: The time history of the spatially averaged Langmuir wave energy  $|E|^2$  with  $|E_p| = 8.10^{-3}$  and  $k_1 \lambda_{De} = 0.09$

with approximately 8 times the initial temperature.

Concerning the wave-wave process, the Langmuir wave envelope and ion acoustic spectra are shown in Figure 4.

Finally, in figure 5 the electric field amplitude  $|E|$  and density perturbation  $\delta n$  are spread out spatially and contains many localized intense regions.




Figure 3: The time history  $W_e(t)$  of electron energy with  $|E_p| = 8.10^{-3}$  and  $k_1 \lambda_{De} = 0.09$ .




Figure 4: The solid line corresponds to the Log of the spatially averaged electron distribution versus the electron energy  $m_e v^2 / 2 v_{the}^2$  at the final time  $\omega_{pe} t = 16000$  with  $|S|/\sqrt{4\pi n_0 T_e} = 8.10^{-3}$  and  $k_p \lambda_{De} = 0.09$  and the dashed line corresponds to the initial electron distribution.




Figure 5: The left plot corresponds to the Langmuir wave in real space and the right plot corresponds to the low frequency density in real space




Figure 6: (A) corresponds to the Langmuir envelope spectra and (B) corresponds to the ion-acoustic spectra at the beginning of nonlinear processes.






Figure 7: (A) corresponds to the Langmuir envelope spectra and (B) corresponds to the ion-acoustic spectra at the final time  $\omega_{pe}t = 16000$ .

We will give more general numerical results and physical interpretation in a further work.

### Acknowledgment

The authors are grateful to V. Tikhonchuk for helpful discussions and suggestions concerning the results of this paper.

### References

- [1] H. Added and S. Added. *Equation of Langmuir turbulence and nonlinear Schrödinger equation : smoothness and approximation*. J. Funct. Anal., Vol. 79, (1988), 183-210.
- [2] B. Bidégaray. *On a nonlocal Zakharov equation*. Nonlinear Anal., Vol. 25 (3), (1995), 247-278.
- [3] T. Colin, G. Metivier. *Ill-Posedness for a degenerate Zakharov System*. In preparation.
- [4] M. Colin, T. Colin. *On a quasilinear Zakharov System describing laser-plasma interactions*. Differential and Integral Equations, 17 (2004), no. 3-4, 297-330.
- [5] J-L. Delcroix and A. Bers. “Physique des plasmas 1, 2”. Inter Editions-Editions du CNRS, (1994).
- [6] J. Ginibre, Y. Tsutsumi and G. Velo. *On the Cauchy problem for the Zakharov system*. J. Funct. Anal., Vol. 151, (1997), 384-436.
- [7] L. Glangetas and F. Merle. *Existence of self-similar blow-up solutions for Zakharov equation in dimension two. I*. Comm. Math. Phys., Vol. 160 (1), (1994), 173-215.
- [8] L. Glangetas and F. Merle. *Concentration properties of blow up solutions and instability results for Zakharov equation in dimension two. II*. Comm. Math. Phys., Vol. 160 (2), (1994), 349-389.

- [9] R. T. Glassey *Convergence of an energy-preserving scheme for the Zakharov equations in one space dimension* Math. Comp., Vol. 58 , (1992), 83-102.
- [10] N. Hayashi and T. Ozawa. *Global, small radially symmetric solutions to nonlinear Schrödinger equations and a gauge transformation.* Diff. and Int. Eqs., Vol. 8, (1995), 1061-1072.
- [11] C.E. Kenig, G. Ponce and L. Vega. *Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations.* Invent. Math., Vol. 134 (3), (1998), 489-545.
- [12] D.E Knuth, *The art of computer programming.* Addison-Wesley Publishing Company, Menlo Park, California, (1969).
- [13] F. Linares, G. Ponce, J.C. Saut. *On a Degenerate Zakharov System.* In preparation.
- [14] T. Ozawa and Y. Tsutsumi. *Existence and smoothing effect of solution for the Zakharov equations.* Publ. Res. Inst. Math. Sci, Vol. 28 (3), (1992), 329-361.
- [15] G.L. Payne, D.R. Nicholson and R.M. Downie, *Numerical Solution of the Zakharov Equations,* J. Compt. Phys., **50**, 482-498 (1983)
- [16] G. Riazuelo. *Etude théorique et numérique de l'influence du lissage optique sur la filamentation des faisceaux lasers dans les plasmas sous-critiques de fusion inertielle.* Thèse de l'Université Paris XI.
- [17] D.A. Russel, D.F. Dubois and H.A. Rose. *Nonlinear saturation of simulated Raman scattering in laser hot spots.* Physics of Plasmas, Vol. 6 (4), (1999), 1294-1317.
- [18] K. Y. Sanbomatsu , *Competition between Langmuir wave-wave and wave-particle interactions,* Thesis of University of Colorado, Department of Astrophysical, 1997.
- [19] S. Schochet and M. Weinstein. *The nonlinear Schrödinger limit of the Zakharov equations governing Langmuir turbulence.* Comm. Math. Phys., Vol. 106, (1986), 569-580.
- [20] C. Sulem and P-L. Sulem. "The nonlinear Schrödinger Equation. Self-Focusing and Wave Collapse." Applied Mathematical Sciences 139, Springer, (1999).
- [21] C. Sulem and P-L. Sulem. *Quelques résultats de régularité pour les équations de la turbulence de Langmuir.* C. R. Acad. Sci. Paris Sér. A-B, Vol. 289 (3), (1979), 173-176.
- [22] B. Texier. *Derivation of the Zakharov equations.* In preparation.
- [23] V.E. Zakharov, S.L. Musher and A.M. Rubenchik. *Hamiltonian approach to the description of nonlinear plasma phenomena.* Phys. Reports, Vol. 129, (1985), 285-366.